
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Laine, E.-M.; Breuer, H.-P.; Piilo, J.

Nonlocal memory effects allow perfect teleportation with mixed states

Published in:
Scientific Reports

DOI:
[10.1038/srep04620](https://doi.org/10.1038/srep04620)

Published: 01/01/2014

Document Version
Publisher's PDF, also known as Version of record

Published under the following license:
CC BY-NC-ND

Please cite the original version:
Laine, E.-M., Breuer, H.-P., & Piilo, J. (2014). Nonlocal memory effects allow perfect teleportation with mixed states. *Scientific Reports*, 4, 1-5. Article 4620. <https://doi.org/10.1038/srep04620>

This material is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.



OPEN

Nonlocal memory effects allow perfect teleportation with mixed states

SUBJECT AREAS:

QUANTUM
INFORMATION

QUBITS

Elsi-Mari Laine^{1,2}, Heinz-Peter Breuer³ & Jyrki Piilo¹

¹Turku Centre for Quantum Physics, Department of Physics and Astronomy, University of Turku, FI-20014 Turun yliopisto, Finland, ²QCD Labs, COMP Centre of Excellence, Department of Applied Physics, Aalto University, P.O. Box 13500, FI-00076 AALTO, Finland, ³Physikalisches Institut, Universität Freiburg, Hermann-Herder-Strasse 3, D-79104 Freiburg, Germany.

Received
29 November 2013Accepted
19 March 2014Published
9 April 2014

Correspondence and
requests for materials
should be addressed to
E.-M.L. (elsi.laine@
aalto.fi)

One of the most striking consequences of quantum physics is quantum teleportation – the possibility to transfer quantum states over arbitrary distances. Since its theoretical introduction, teleportation has been demonstrated experimentally up to the distance of 143 km. In the original proposal two parties share a maximally entangled quantum state acting as a resource for the teleportation task. If, however, the state is influenced by decoherence, perfect teleportation can no longer be accomplished. Therefore, one of the current major challenges in accomplishing teleportation over long distances is to overcome the limitations imposed by decoherence and the subsequent mixedness of the resource state. Here we show that, in the presence of nonlocal memory effects, perfect quantum teleportation can be achieved even with mixed photon polarisation states. Our results imply that memory effects can be exploited in harnessing noisy quantum systems for quantum communication and that non-Markovianity is a resource for quantum information tasks.

Quantum teleportation is the best known example of how entanglement serves as a valuable resource for information processing. Since the original discovery of quantum teleportation by Bennett *et al.* in 1993¹, remarkable technological advances have been made and teleportation with higher than classically achievable fidelities has been demonstrated experimentally^{2–7} up to the distance of 143 km⁸. In practice, perfect teleportation is not possible since, instead of maximally entangled states, decoherence effects force us to deal with mixed states. It has been shown that with classical communication and local operations perfect teleportation with mixed states is impossible^{1,9} and therefore, one of the current major challenges in accomplishing teleportation over long distances is to overcome the limitations imposed by decoherence and the subsequent mixedness of the resource state.

When an open quantum system, due to its coupling with the external environment, continuously loses information to its surroundings, the noise induced dynamics is called Markovian^{10–12}. Non-Markovian quantum dynamics with memory effects arise when the system does not only lose information, but temporarily recovers some of it from the environment at a later time^{13,14}. Recently, a significant progress in developing a general theory of non-Markovian quantum dynamics^{13–20} as well as in the experimental detection and control of memory effects^{21–23} has been made.

Previously, environment-assisted quantum processes have been studied in the context of e.g. quantum transport^{24,25} and communication networks²⁶, but protocols for quantum information tasks utilising the non-Markovian nature of otherwise harmful noise have been so far missing. In this paper we show how quantum memory effects can be harnessed to give an advantage in mixed state quantum teleportation in the presence of dephasing noise. We consider photonics realisations due to their dominant role in the experimental implementations of teleportation⁸. The key element of the scheme introduced here are nonlocal memory effects where the local exposure of the bipartite quantum system to Markovian noise can create strong global memory effects²⁷; a scheme which has also been recently experimentally demonstrated²³. The fundamental source for these effects are the initial correlations between the local environments of the bipartite open system. For entangled photon polarisation states, nonlocal memory effects needed for the protocol arise naturally since the frequency distributions of the photons, which act as environments, are unavoidably correlated after a downconversion process²³. We also analyse how the amount of initial correlations between the environments affects the fidelity of the teleportation, and find that even in the absence of maximal correlations between the environments, the protocol will substantially outdo the performance of the known optimal protocol without memory effects⁹.

In the standard quantum teleportation protocol Alice has a qubit whose state $|\phi\rangle_1 = \alpha|+\rangle + \beta|-\rangle$ she wants to teleport to Bob. Alice and Bob share an entangled pair of qubits 2 and 3 in the Bell-state:



$$|\phi^+\rangle_{23} = \frac{1}{\sqrt{2}}(|++\rangle + |--\rangle). \tag{1}$$

Alice performs a Bell-state measurement on the particles 1 and 2, which projects particle 3 in Bob's hands into one of four states depending on which Bell-state Alice gains as an outcome. Alice further communicates her measurement outcome to Bob who performs a unitary operation on particle 3 depending on the outcome of Alice. The final state of particle 3 is the original state $|\phi\rangle_3 = \alpha|+\rangle + \beta|-\rangle$. During the Bell-state measurement particle 1 becomes entangled with particle 2 and the state $|\phi\rangle_1$ is destroyed on Alice's side during the protocol.

However, if the entangled pair of particles that Alice and Bob share is disturbed by noise, the fidelity of the standard teleportation goes down radically. Even if one chooses, instead of the standard teleportation scheme, an optimised protocol, perfect teleportation can no longer be achieved⁹. Let us now demonstrate how perfect teleportation can be achieved with photon polarisation states, even in the presence of dephasing noise, if the pair shared by Alice and Bob is influenced by nonlocal memory effects, see Fig. 1.

Results

Let us assume that the pair of entangled photons is created in a spontaneous parametric downconversion process after which the two particles are sent to remote locations to Alice and Bob. However, the particle sent to Alice is not perfectly isolated, but is interacting with its local environment giving rise to local decoherence, which destroys the entanglement between the two parties. As a physical implementation of the local dephasing noise, we consider photon traveling through a quartz plate, where the polarisation degree of freedom (system) and the frequency degree of freedom (environment) interact^{22,23,27}. The quartz plate exposes the polarisation state of the photon to a random external field giving rise to a dephasing evolution²⁸.

Now, Alice wishes to teleport the state $|\phi\rangle_1 = \alpha|H\rangle + \beta|V\rangle$ to Bob. Here, H (V) refers to horizontal (vertical) polarisation state of the photon. Initially Alice and Bob share the Bell-state $|\phi^+\rangle_{23} = \frac{1}{\sqrt{2}}(|HH\rangle + |VV\rangle)$ and the total initial state (system and environment) of the photons 2 and 3 is

$$|\phi(0)\rangle = |\phi^+\rangle_{23} \otimes \int d\omega_2 d\omega_3 g(\omega_2, \omega_3) |\omega_2\rangle |\omega_3\rangle, \tag{2}$$

where $g(\omega_2, \omega_3)$ gives the joint frequency amplitude of the photons 2 and 3 and $\int d\omega_2 d\omega_3 |g(\omega_2, \omega_3)|^2 = 1$. The Hamiltonian for the local dephasing due to the quartz plate takes the form

$$H_i = - \int d\omega_i \omega_i [n_V^i |V\rangle\langle V| + n_H^i |H\rangle\langle H|] \otimes |\omega_i\rangle\langle\omega_i|, \tag{3}$$

where, e.g., $|V\rangle \otimes |\omega_i\rangle$ denotes the state of photon i ($i = 2, 3$) with polarisation V and frequency ω_i , and n_H^i (n_V^i) is the index of refraction for polarisation component H (V). The time dependent interaction Hamiltonian describing the evolution of the two-photon state is given by

$$H_i(t) = \chi_2(t)H_2 + \chi_3(t)H_3, \tag{4}$$

where

$$\chi_i(t) = \begin{cases} 1 & \text{if } t_{in}^i \leq t \leq t_{out}^i \\ 0 & \text{otherwise} \end{cases} \tag{5}$$

and t_{in}^i denotes the time photon i enters the quartz plate and t_{out}^i the time photon i exits the quartz plate. Now the time evolution of the total system is given by $|\psi(t)\rangle = \exp[-i\int_0^t dt' H_I(t')] |\psi(0)\rangle$. Let us also write $t_i = \int_0^\infty \chi_i(t) dt = t_{out}^i - t_{in}^i$.

In terms of our protocol, first photon 2 interacts with its quartz plate followed by the Bell-state measurement on photons 1 and 2. Then Alice communicates her measurement outcome to Bob and the interaction of photon 3 with its local environment in Bob's side finishes the procedure. The combined polarisation state of the systems 2 and 3, after photon 2 has interacted with its quartz plate on Alice's side, is

$$\rho_{23}(t_2) = \frac{1}{2} (|HH\rangle\langle HH| + \kappa_2(t_2)|HH\rangle\langle VV| + \kappa_2^*(t_2)|VV\rangle\langle HH| + |VV\rangle\langle VV|), \tag{6}$$

where the decoherence function κ_2 is

$$\kappa_2(t_2) = \int d\omega_2 d\omega_3 |g(\omega_2, \omega_3)|^2 e^{-i\Delta n_2 \omega_2 t_2}, \tag{7}$$

$\Delta n_2 = n_V^2 - n_H^2$ is the birefringence, and t_2 the interaction time of photon 2. Now, if Alice and Bob were to perform the standard teleportation with the shared state $\rho_{23}(t_2)$, the fidelity of the teleportation would decrease linearly with respect to κ_2 , see Fig. 2. Let us now describe how they can outperform this fidelity with nonlocal memory effects.

After the local interaction of photon 2 the total state for the three photons is

$$|\Psi(t_2)\rangle = \frac{1}{\sqrt{2}} |\phi\rangle_1 (|HH\rangle |\psi_{HH}(t_2)\rangle + |VV\rangle |\psi_{VV}(t_2)\rangle), \tag{8}$$

where

$$\begin{aligned} |\psi_{HH}(t_2)\rangle &= \int d\omega_2 d\omega_3 g(\omega_2, \omega_3) e^{in_H^2 \omega_2 t_2} |\omega_2\rangle |\omega_3\rangle, \\ |\psi_{VV}(t_2)\rangle &= \int d\omega_2 d\omega_3 g(\omega_2, \omega_3) e^{in_V^2 \omega_2 t_2} |\omega_2\rangle |\omega_3\rangle. \end{aligned} \tag{9}$$

This can be written in the form




Figure 1 | A schematic picture of the teleportation protocol with nonlocal memory effects. EPR refers to a source producing states of the form of equation (1) and BSM refers to a Bell-state measurement. The local noise before the BSM makes the EPR pair mixed while the noise in the last step of the protocol allows to recover the teleported state.




Figure 2 | The fidelities of the standard, optimal and non-Markovian protocols. The fidelities of different teleportation protocols for the mixed state of equation (6) as a function of the decoherence function $|\kappa_2|$. The red dashed line represents the fidelity of the standard protocol F_w^M [equation (20)], the red dot dashed line the fidelity of the optimal protocol F^{opt} [equation (21)], the red solid lines the fidelity of the protocol with memory effects F_w^{NM} with $\delta K \in [0, 0.1]$ [equation (19)] and the light red solid lines with $\delta K \in [0.1, 0.5]$.

$$\begin{aligned}
 |\Psi(t_2)\rangle = & \frac{1}{2} |\phi^+\rangle_{12} (\alpha|H\rangle_3 |\psi_{HH}(t_2)\rangle + \beta|V\rangle_3 |\psi_{VV}(t_2)\rangle) \\
 & + \frac{1}{2} |\phi^-\rangle_{12} (\alpha|H\rangle_3 |\psi_{HH}(t_2)\rangle - \beta|V\rangle_3 |\psi_{VV}(t_2)\rangle) \\
 & + \frac{1}{2} |\psi^+\rangle_{12} (\beta|H\rangle_3 |\psi_{HH}(t_2)\rangle + \alpha|V\rangle_3 |\psi_{VV}(t_2)\rangle) \\
 & + \frac{1}{2} |\psi^-\rangle_{12} (\alpha|V\rangle_3 |\psi_{VV}(t_2)\rangle - \beta|H\rangle_3 |\psi_{HH}(t_2)\rangle),
 \end{aligned} \quad (10)$$

where

$$\begin{aligned}
 |\phi^+\rangle = & \frac{1}{\sqrt{2}} (|HH\rangle + |VV\rangle), \quad |\phi^-\rangle = \frac{1}{\sqrt{2}} (|HH\rangle - |VV\rangle), \\
 |\psi^+\rangle = & \frac{1}{\sqrt{2}} (|HV\rangle + |VH\rangle), \quad |\psi^-\rangle = \frac{1}{\sqrt{2}} (|HV\rangle - |VH\rangle)
 \end{aligned} \quad (11)$$

are the Bell states. Alice then performs the Bell state measurement and communicates her results to Bob. Bob further applies a unitary operation on his particle depending on Alice's measurement outcome. Thus, so far, Alice and Bob have performed the standard teleportation scheme with the mixed state of equation (6). Now, in order to improve the protocol, Bob needs to harness the nonlocal memory effects. He can use the information sent by Alice and subject his particle to conditional noise depending on Alice's measurement outcome. By adding the conditional noise to his system, Bob actually cancels out the effect of the noise which earlier acted on Alice's system. He chooses the following unitary operations and the birefringence of his quartz plate to produce noise according to Alice's outcomes:

$$\begin{aligned}
 |\varphi^+\rangle & \Rightarrow \mathbb{I}, \quad \Delta n_3 = \Delta n_2 \\
 |\varphi^-\rangle & \Rightarrow \sigma_z, \quad \Delta n_3 = \Delta n_2 \\
 |\psi^+\rangle & \Rightarrow \sigma_x, \quad \Delta n_3 = -\Delta n_2 \\
 |\psi^-\rangle & \Rightarrow i\sigma_y, \quad \Delta n_3 = -\Delta n_2.
 \end{aligned} \quad (12)$$

Let us assume that Alice's outcome of the Bell measurement is the state $|\phi^+\rangle_{12}$. Now the total state of Bob's photon 3 is

$$\begin{aligned}
 & |\alpha|^2 |H\rangle\langle H| \otimes \rho_{HH} + \alpha\beta^* |H\rangle\langle V| \otimes \rho_{HV} \\
 & + \alpha^* \beta |V\rangle\langle H| \otimes \rho_{VH} + |\beta|^2 |V\rangle\langle V| \otimes \rho_{VV},
 \end{aligned} \quad (13)$$

where

$$\rho_{HH} = \int d\omega_3 d\omega'_3 d\omega_2 g(\omega_2, \omega_3) g^*(\omega_2, \omega'_3) |\omega_3\rangle\langle \omega'_3| = \rho_{VV}, \quad (14)$$

$$\rho_{HV} = \int d\omega_3 d\omega'_3 \tilde{g}(\omega_3, \omega'_3) |\omega_3\rangle\langle \omega'_3| = \rho_{VH}^\dagger,$$

and $\tilde{g}(\omega_3, \omega'_3) = \int d\omega_2 g(\omega_2, \omega_3) g^*(\omega_2, \omega'_3) e^{-i\Delta n_2 \omega_2 t_2}$. When Bob subjects his photon to noise by putting his photon through a quartz plate with $\Delta n_3 = \Delta n_2$ his final state can be written as

$$\begin{aligned}
 \rho_F(t_2, t_3) = & |\alpha|^2 |H\rangle\langle H| + \alpha\beta^* \kappa(t_2, t_3) |H\rangle\langle V| \\
 & + \alpha^* \beta \kappa^*(t_2, t_3) |V\rangle\langle H| + |\beta|^2 |V\rangle\langle V|,
 \end{aligned} \quad (15)$$

where the decoherence function is

$$\kappa(t_2, t_3) = \int d\omega_2 d\omega_3 |g(\omega_2, \omega_3)|^2 e^{-i\Delta n_2 (\omega_2 t_2 + \omega_3 t_3)}, \quad (16)$$

and t_3 is the interaction time in Bob's quartz plate.

In the description of the downconversion process, the frequency distribution can be taken to be a joint Gaussian distribution^{23,27}

$$|g(\omega_2, \omega_3)|^2 = \frac{1}{2\pi\sqrt{\det C}} e^{-\frac{1}{2}(\vec{\omega} - \langle \vec{\omega} \rangle)^T C^{-1} (\vec{\omega} - \langle \vec{\omega} \rangle)}, \quad (17)$$

where $C = (C_{ij})$ is the covariance matrix with elements $C_{ij} = \langle \omega_i \omega_j \rangle - \langle \omega_i \rangle \langle \omega_j \rangle$. We assume that both the means and the variances of ω_2 and ω_3 are equal, i.e., $\langle \omega_2 \rangle = \langle \omega_3 \rangle = \omega_0/2$ and $C_{11} = C_{22} = \langle \omega_i^2 \rangle - \langle \omega_i \rangle^2$. To quantify the frequency correlations we use the correlation coefficient $K = C_{12}/\sqrt{C_{11}C_{22}} = C_{12}/C_{11}$ satisfying $|K| \leq 1$.

If Bob chooses $t_3 = t_2$ and the frequency distribution is of the form of equation (17) with the correlation coefficient $K = -1$, the polarisation state of photon 3 is given by equation (15) with $\kappa(t_2, t_3) = e^{i\omega_0 \Delta n_2 t_2}$, i.e., the magnitude of the decoherence function has returned to its original value equal to 1. If Bob now performs a phase gate with phase $-\omega_0 \Delta n_2 t_2$, perfect teleportation has been completed. The other measurement outcomes of Alice give the same result if Bob applies the operations given in equation (12). It is important to notice that all the steps of this protocol can be applied locally, and that the applied noise is locally Markovian and globally non-Markovian within the bipartite system²⁷.

Discussion

We have demonstrated how Alice and Bob can recover perfect teleportation in the presence of dephasing noise if Bob after the standard teleportation protocol subjects his particle to noise. The key element in the protocol are the nonlocal memory effects which arise when the local decoherence is present on Bob's side. The source of the nonlocal memory effects lies in the initial correlations between the local environments described by the correlation coefficient K , which in turn is determined by the width of the original pump pulse in the downconversion process. In the case of perfect anticorrelation $K = -1$ (delta peak pump), perfect teleportation is achieved and for an uncorrelated distribution $K = 0$ no advantage with respect to the standard protocol can be gained. It is also important to note that no initial entanglement is necessary between the local environments, but classical correlations suffice^{23,27}.

Thus, we have shown that when the initial joint frequency distribution has perfect anticorrelation $K = -1$, this leads to perfect teleportation with mixed polarisation states. If, however, the initial



pump pulse does not have a delta peak distribution, the fidelity of the protocol decreases. In the following we study how a finite width of the pump pulse will affect the fidelity of the teleportation protocol. Let us take the correlation coefficient to be $K = -1 + \delta K$, i.e., δK measures the deviation from the ideal case. The fidelity $F = \langle \phi | \rho_3 | \phi \rangle_1$ between the original state $\alpha |H\rangle + \beta |V\rangle$ and the teleported state becomes

$$F = 1 - 2|\alpha|^2|\beta|^2 \left(1 - |\kappa_2(t_2)|^{2\delta K}\right). \quad (18)$$

Considering the worst case scenario, i.e. $|\alpha|^2 = |\beta|^2 = 1/2$, we obtain

$$F_w^{\text{NM}} = 1 - \frac{1}{2} \left(1 - |\kappa_2(t_2)|^{2\delta K}\right). \quad (19)$$

If the teleportation was performed with the decohered state in equation (6), using the standard scheme without taking advantage of the memory effects, we would have the fidelity

$$F_w^{\text{M}} = 1 - \frac{1}{2} (1 - |\kappa_2(t_2)|). \quad (20)$$

If instead of the standard teleportation scheme, one would use an optimal teleportation scheme for the decohered state at hand⁹, one would get the fidelity

$$F_w^{\text{opt}} = \frac{|\kappa_2(t_2)| + 2}{3} \quad (21)$$

which gives the value $2/3$ for a classical state. The fidelities of the different teleportation protocols are plotted in Fig. 2 as a function of the decoherence function $|\kappa_2(t_2)|$. We see that the fidelity of the protocol with memory effects for sufficiently large values of the correlation coefficient $|K|$ is still much larger than the fidelity of the standard protocol or the optimal protocol in the absence of memory effects. For sufficiently large correlations ($\delta K \leq 0.1$, i.e. $-1.0 \leq K \leq -0.9$) the fidelity of the protocol with memory effects exceeds the fidelity of the optimal protocol without memory effects all the way to an almost fully decohered state. For a smaller value of correlations ($0.1 \leq \delta K \leq 0.5$) the fidelity still exceeds the one of the standard protocol. It is also important to note that the high value of the correlation coefficient $K = -0.9$ (or $\delta K = 0.1$) is experimentally realisable²³.

In Fig. 3, we plot the fidelity of the proposed protocol as a function of the non-Markovianity measure \mathcal{N} introduced in¹³. Assuming $t_3 = t_2$, $t_{\text{in}}^2 = 0$, and $t_{\text{in}}^3 = t_{\text{out}}^2$, the dynamical map describing the teleportation




Figure 3 | The fidelity of the proposed protocol as a function of the non-Markovianity measure \mathcal{N} . In the inset the different values of the decoherence function $|\kappa_2(t_2)|$ are given.

protocol for Alice's initial unknown state $|\phi\rangle = \alpha |H\rangle + \beta |V\rangle$ can be written as

$$\Phi_t[|\phi(0)\rangle] = \begin{cases} \rho_F(t,0) & \text{if } t \leq t_2 \\ \rho_F(t_2, t-t_2) & \text{if } t_2 \leq t \leq 2t_2. \end{cases} \quad (22)$$

Here, ρ_F is given by Eq. (15), $\rho_F(t, 0)$ denotes the case where the teleportation is performed at time t with no quartz plate on Bob's side, and $\rho_F(t_2, t-t_2)$ the case of teleportation at time t_2 with subsequent evolution with Bob's quartz plate. The non-Markovianity measure \mathcal{N} can now be written analytically²³:

$$\mathcal{N} = |\kappa_2(t_2)|^{1-K^2} - |\kappa_2(t_2)|. \quad (23)$$

Thus, the fidelity of the protocol in equation (19) as a function of the non-Markovianity is

$$F_w^{\text{NM}}(\mathcal{N}) = 1 - \frac{1}{2} \left(1 - |\kappa_2(t_2)|^2 \left(1 - \sqrt{1 - \frac{\ln(\mathcal{N} + |\kappa_2(t_2)|)}{\ln|\kappa_2(t_2)|}}\right)\right) \quad (24)$$

As can be seen in Fig. 3, the fidelity is a monotonically increasing function of the non-Markovianity and thus the memory effects act as a resource for quantum teleportation.

Summarising, we have found that nonlocal memory effects can substantially increase the fidelity of mixed state quantum teleportation in the presence of dephasing noise. In the protocol Alice and Bob act on their particles locally and the nonlocal memory effects occur due to initial correlations between the local environments of the photons. In order for Bob to harness the memory effects, he needs to subject his photon to local noise after the standard teleportation protocol. We have shown that one can perform perfect mixed state teleportation of photon polarisation states if the environments of the two photons share maximal initial correlations. The protocol presented here demonstrates how to overcome dephasing noise induced by random external fields in a quantum communication setup by exploiting memory effects. The result serves as an encouraging example of how non-Markovianity can serve as a resource for quantum information tasks but it still remains to be shown whether the same technique could be applied for environments giving rise to also relaxation and dissipation.

- Bennett, C. H. *et al.* Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels. *Phys. Rev. Lett.* **70**, 1895 (1993).
- Bouwmeester, D. *et al.* Experimental quantum teleportation. *Nature* **390**, 575 (1997).
- Boschi, D., Branca, S., De Martini, F., Hardy, L. & Popescu, S. Experimental realization of teleporting an unknown pure quantum state via dual classical and Einstein-Podolsky-Rosen channels. *Phys. Rev. Lett.* **80**, 1121 (1998).
- Kim, Y.-H., Kulik, S. P. & Shih, Y. Quantum teleportation of a polarization state with a complete Bell state measurement. *Phys. Rev. Lett.* **86**, 1370 (2001).
- Marcikic, I., de Riedmatten, H., Tittel, W., Zbinden, H. & Gisin, N. Long-distance teleportation of qubits at telecommunication wavelengths. *Nature* **421**, 509 (2003).
- Riebe, M. *et al.* Deterministic quantum teleportation with atoms. *Nature* **429**, 734 (2004).
- Barrett, M. D. *et al.* Deterministic quantum teleportation of atomic qubits. *Nature* **429**, 737 (2004).
- Ma, X.-S. *et al.* Quantum teleportation over 143 kilometres using active feed-forward. *Nature* **489**, 269 (2012).
- Verstraete, F. & Verschelde, H. Optimal teleportation with a mixed state of two qubits. *Phys. Rev. Lett.* **90**, 097901 (2003).
- Lindblad, G. On the generators of quantum dynamical semigroups. *Commun. Math. Phys.* **48**, 119 (1976).
- Gorini, V., Kossakowski, A. & Sudarshan, E. C. G. Completely positive dynamical semigroups of N-level systems. *J. Math. Phys.* **17**, 821 (1976).
- Breuer, H.-P. & Petruccione, F. *The theory of open quantum systems* (Oxford University Press, Oxford, 2007).
- Breuer, H.-P., Laine, E.-M. & Piilo, J. Measure for the degree of non-Markovian behavior of quantum processes in open systems. *Phys. Rev. Lett.* **103**, 210401 (2009).
- Laine, E.-M., Piilo, J. & Breuer, H.-P. Measure for the non-Markovianity of quantum processes. *Phys. Rev. A* **81**, 062115 (2010).



15. Piilo, J., Maniscalco, S., Härkönen, K. & Suominen, K.-A. Non-Markovian quantum jumps. *Phys. Rev. Lett.* **100**, 180402 (2008).
16. Breuer, H.-P. & Vacchini, B. Quantum semi-Markov processes. *Phys. Rev. Lett.* **101**, 140402 (2008).
17. Wolf, M. M., Eisert, J., Cubitt, T. S. & Cirac, J. I. Assessing non-Markovian quantum dynamics. *Phys. Rev. Lett.* **101**, 150402 (2008).
18. Rivas, Á., Huelga, S. F. & Plenio, M. B. Entanglement and non-Markovianity of quantum evolutions. *Phys. Rev. Lett.* **105**, 050403 (2010).
19. Chruściński, D. & Kossakowski, A. Non-Markovian quantum dynamics: local versus nonlocal. *Phys. Rev. Lett.* **104**, 070406 (2010).
20. Vacchini, B., Smirne, A., Laine, E.-M., Piilo, J. & Breuer, H.-P. Markovian and non-Markovian dynamics in quantum and classical systems. *New J. Phys.* **13**, 093004 (2011).
21. Tang, J.-S. *et al.* Measuring non-Markovianity of processes with controllable system-environment interaction. *Europhys. Lett.* **97**, 10002 (2012).
22. Liu, B.-H. *et al.* Experimental control of the transition from Markovian to non-Markovian dynamics of open quantum systems. *Nature Phys.* **7**, 931-934 (2011).
23. Liu, B.-H. *et al.* Photonic realization of nonlocal memory effects and non-Markovian quantum probes. *Sci. Rep.* **3**, 1781 (2013).
24. Mohseni, M. *et al.* Environment-assisted quantum walks in photosynthetic energy transfer. *J. Chem. Phys.* **129**, 174106 (2008).
25. Reberstrost, P. *et al.* Environment-assisted quantum transport. *New J. Phys.* **11**, 033003 (2009).
26. Caruso, F., Huelga, S. F. & Plenio, M. B. Noise-Enhanced Classical and Quantum Capacities in Communication Networks. *Phys. Rev. Lett.* **105**, 190501 (2010).
27. Laine, E.-M., Breuer, H.-P., Piilo, J., Li, C.-F. & Guo, G.-C. Nonlocal memory effects in the dynamics of open quantum systems. *Phys. Rev. Lett.* **108**, 210402 (2012).
28. Alicki, R. & Lendi, K. *Quantum Dynamical Semigroups and Applications* (Springer, Berlin Heidelberg, 2007).

Acknowledgments

This work was supported by Academy of Finland (259827, mobility from Finland), Jenny and Antti Wihuri Foundation, the Graduate School of Modern Optics and Photonics, and the German Academic Exchange Service (DAAD).

Author contributions

E.-M.L. carried out the theoretical analysis. E.-M.L., H.-P.B. and J.P. contributed to the development of the interpretation, writing the paper and discussing the contents.

Additional information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Laine, E.-M., Breuer, H.-P. & Piilo, J. Nonlocal memory effects allow perfect teleportation with mixed states. *Sci. Rep.* **4**, 4620; DOI:10.1038/srep04620 (2014).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. The images in this article are included in the article's Creative Commons license, unless indicated otherwise in the image credit; if the image is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the image. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>