
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Volovik, G.E.; Kopnin, N.B.

Comment on "Scaling of the quasiparticle spectrum for d-wave superconductors"

Published in:
Physical Review Letters

DOI:
[10.1103/PhysRevLett.78.5028](https://doi.org/10.1103/PhysRevLett.78.5028)

Published: 01/01/1997

Document Version
Publisher's PDF, also known as Version of record

Please cite the original version:
Volovik, G. E., & Kopnin, N. B. (1997). Comment on "Scaling of the quasiparticle spectrum for d-wave superconductors". *Physical Review Letters*, 78(26), 5028. <https://doi.org/10.1103/PhysRevLett.78.5028>

This material is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Comment on “Scaling of the Quasiparticle Spectrum for d -wave Superconductors”

In a recent Letter, Simon and Lee (SL) [1] suggested a scaling law for thermodynamic and kinetic properties of superconductors with lines of gap nodes. For example, the heat capacity as a function of temperature and magnetic field for $T \ll T_c$ and $H \ll H_{c2}$ is

$$C(T, H) = aT^2 G(x), \quad x = \alpha T/H^{1/2}, \quad (1)$$

where $G(x)$ is a dimensionless function of the dimensionless parameter x . Equation (1) is in agreement with our calculations [2–4] if the scaling parameter is $x_{KV} = (H_{c2}/H)^{1/2}(T/T_c)$ (apart from a logarithmic factor). Indeed, according to [2], the function $G(x) \sim 1 + (1/x^2)$ for large values of x ; the first term is the bulk heat capacity $C(T, H) \propto T^2$, while the second term results in a temperature-independent vortex contribution $C(T, H) \propto H$. For small x , the function $G(x) \sim 1/x$ which gives [3,4] $C(T, H) \propto T\sqrt{H}$. The crossover value between these two regimes is $x_{KV} \sim 1$. However, SL have obtained the crossover parameter $x_{SL} \sim (H_{c2}/H)^{1/2}(T/T_c)\sqrt{E_F/T_c}$. The difference between our x and that obtained by SL is thus by the large factor $\sqrt{E_F/T_c}$. We discuss the origin of this disagreement.

Let us introduce the anisotropic Fermi momentum $p_F(\theta)$ which depends on the angle θ in the a - b plane. If $T \ll T_c$ the quasiparticles which are close to the gap node $\theta \ll 1$ are important. Their spectrum is

$$E(\mathbf{p}) = \sqrt{v_F^2 [p - p_F(\theta)]^2 + (\Delta')^2 \theta^2}, \quad (2)$$

where v_F is the Fermi velocity and Δ' is the angular derivative of the gap, $\Delta(\theta) \approx \theta\Delta'$; both are in a vicinity of the node. Equation (2) was the starting point in [2,4].

In contrast, SL used the linearized spectrum

$$E(\mathbf{p}) = \sqrt{c_{\parallel}^2 (\delta p_x)^2 + c_{\perp}^2 p_y^2}, \quad (3)$$

where $\mathbf{p} = p_y \hat{y} + (p_F + \delta p_x) \hat{x}$, $c_{\parallel} = v_F$, and $c_{\perp} = \Delta'/p_F$. This is justified when the nonlinear contributions to

$$\begin{aligned} \epsilon(\mathbf{p}) - E_F &= p_x^2/2m_x + p_y^2/2m_y - E_F \\ &\approx v_F \delta p_x + p_y^2/2m_y \end{aligned}$$

can be neglected, i.e., when $p_y^2/2m_y \ll c_{\perp} p_y$ where $p_y = p_F \theta$. This requires much stronger restrictions both on the angle $\theta \ll T_c/E_F$ and on the energy and temperature $T \ll T_c^2/E_F$.

At first glance, one might expect that the temperature of the order of T_c^2/E_F marks the boundary between our scaling and that by SL. However, this is not the case. Our quasiclassical approach is valid down to the temperature

at which a discreteness of fermion bound states in the vortex background becomes important. For s -wave superconductors, the interlevel spacing of core fermions is of the order of T_c^2/E_F [5]; thus the quantum limit is reached at $T \sim T_c^2/E_F$. In d -wave superconductors, in a vicinity of the gap nodes, the interlevel distance is smaller; it is determined by a large dimension of the wave function which, for low energies, is limited by the intervortex distance R [2,3]. Thus the discreteness of the levels becomes important at lower temperatures $T \sim (T_c^2/E_F)(\xi/R) \sim (T_c^2/E_F)\sqrt{B/B_{c2}}$.

Therefore, one expects two changes of the regime with the crossover parameters as follows: (1) At $x_{KV} \sim 1$ [i.e., at $(H_{c2}/H)^{1/2}(T/T_c) \sim 1$] the single-vortex contribution to the thermodynamic quantity is comparable with the bulk contribution per one vortex. (2) At $x_{KV} \sim T_c/E_F$ [i.e., at $(H_{c2}/H)^{1/2}(T/T_c)(E_F/T_c) \sim 1$] the quasiclassical regime changes to the quantum one. However, there is no change in the regime at the SL scale, $x_{SL} \sim 1$ [i.e., at $(H_{c2}/H)^{1/2}(T/T_c)\sqrt{E_F/T_c} \sim 1$]. This is because the high anisotropy of the conical spectrum in Eq. (3) was not taken properly into account in [1]: The rescaling of coordinates to get an isotropic spectrum of fermions with the average “speed of light” $c = \sqrt{c_{\parallel}c_{\perp}}$ [1] leads to a high deformation of the potential well produced by vortices.

G.E. Volovik^{1,2} and N.B. Kopnin^{2,3}

¹Low Temperature Laboratory
Helsinki University of Technology
P.O. Box 2200
FIN-02015 HUT
Espoo, Finland

²Landau Institute for Theoretical Physics
117334 Moscow, Russia

³Laboratoire de Physique des Solides
Université Paris-Sud
Bât 510
91405 Orsay, France

Received 11 March 1997 [S0031-9007(97)03414-5]
PACS numbers: 74.25.Fy, 74.25.Jb, 74.72.-h

- [1] S.H. Simon and P.A. Lee, Phys. Rev. Lett. **78**, 1548 (1997).
- [2] N.B. Kopnin and G.E. Volovik, Pis'ma Zh. Eksp. Teor. Fiz. **64**, 641 (1996) [JETP Lett. **64**, 690 (1996)].
- [3] G.E. Volovik, Pis'ma Zh. Eksp. Teor. Fiz. **58**, 457 (1993) [JETP Lett. **58**, 469 (1993)].
- [4] G.E. Volovik, Pis'ma Zh. Eksp. Teor. Fiz. **65**, 465 (1997); Report No. cond-mat/9702093.
- [5] C. Caroli, P.G. de Gennes, and J. Matricon, Phys. Lett. **9**, 307 (1964).