



This is an electronic reprint of the original article. This reprint may differ from the original in pagination and typographic detail.

Sippola, Hannu; Taskinen, Pekka

Activity of Supercooled Water on the Ice Curve and Other Thermodynamic Properties of Liquid Water up to the Boiling Point at Standard Pressure

Published in: Journal of Chemical and Engineering Data

DOI: 10.1021/acs.jced.8b00251

Published: 09/08/2018

Document Version Publisher's PDF, also known as Version of record

Published under the following license: CC BY

Please cite the original version:

Sippola, H., & Taskinen, P. (2018). Activity of Supercooled Water on the Ice Curve and Other Thermodynamic Properties of Liquid Water up to the Boiling Point at Standard Pressure. *Journal of Chemical and Engineering Data*, 63(8), 2986-2998. https://doi.org/10.1021/acs.jced.8b00251

This material is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

This is an open access article published under a Creative Commons Attribution (CC-BY) License, which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.



Article pubs.acs.org/jced

Activity of Supercooled Water on the Ice Curve and Other Thermodynamic Properties of Liquid Water up to the Boiling **Point at Standard Pressure**

Hannu Sippola^{*,#} and Pekka Taskinen

Aalto University School of Chemical Engineering, Department of Chemical Engineering and Metallurgy Metallurgical Thermodynamics and Modelling P.O. Box 16200, Aalto, FI-00076, Finland

ABSTRACT: A simple model for thermodynamic properties of water from subzero temperatures up to 373 K was derived at ambient pressure. The heat capacity of supercooled water was assessed as lambda transition. The obtained properties for supercooled water such as heat capacity, vapor pressure, density and thermal expansion are in excellent agreement with literature data. Activity of water on ice curve, independent of used electrolyte and Debye-Hückel constant applied in modeling, is also calculated. Thus, the ice curve activity of supercooled water can be used as a universal basis for thermodynamic modeling of aqueous solutions, precipitating hydrated and anhydrous solids. A simple model for heat capacity, density and thermal expansion of ice are also derived from 170 K up to melting point.



INTRODUCTION

The peculiar thermodynamic properties of supercooled water and its practical importance for modeling aqueous solution at subzero temperatures (e.g., ref¹) have been gaining more and more attention. The thermodynamic properties of liquid water are well-known over 273.15 K at standard pressure but are still ambiguous below 273.15 K mainly due to anomalies in its thermophysical properties caused by hydrogen bonds.²⁻

The thermodynamic properties of supercooled liquid water in equilibrium with ice Ih, that is, the ice curve, are generally obtained by the assessment of experimental ice curve data for one or more electrolytes. However, the activity of water on the ice curve is independent of the studied electrolyte system and it depends only on the thermodynamic properties of ice and pure metastable supercooled water.^{7,8} Similar results are obtained also for homogeneous⁸ and heterogeneous ice nucleation.⁵

Freezing point depression is used among other Gibbs energy related experimental data when fitting activity coefficient parameters. Usually in evaluation of the activity of water at subzero temperatures, the heat capacity difference between supercooled liquid water and solid ice (ΔC_p) is assumed either insignificant, that is zero, or independent of temperature. However, the heat capacity of supercooled water starts to increase rapidly below -10 °C while the heat capacity of ice decreases steadily, so the assumption of constant heat capacity difference below -10 °C is not usable in accurate thermodynamic modeling. Moreover, the calculated activity of water in subzero aqueous solutions will also depend on the equation used for the Debye-Hückel constant. Both items reflect also to the estimated vapor pressure of water solution at subzero temperatures, which is important in meteorological and climate models.

The purpose of this study is to generate a practical thermodynamic expression for the supercooled liquid water for modeling purposes of aqueous electrolyte systems in 1 atm total pressure. The aim is to simplify the universal approach in the T-P domain (e.g., ref 10) for tools and thermodynamic descriptions available in most software used in various engineering approaches of thermodynamic modeling and aqueous simulation.

THEORY

In aqueous solution the chemical potential of the solvent, that is water, is defined as

$$\mu_{w}(T, p) = \mu_{w}^{o}(T, p) + RT \ln(a_{w}^{s})$$
⁽¹⁾

where the standard state for a solution is pure liquid water at temperature and pressure of the solution and a_w is the activity of liquid water. Superscript s indicates a solution, R is the gas constant, and T is temperature in Kelvin.

The chemical potential of pure liquid water at any temperature and pressure is

$$\mu_{\rm w}^{*}(T, p) = \mu_{\rm w}^{\rm o}(T) + RT \ln(a_{\rm w}^{*})$$
⁽²⁾

where the standard state of pure liquid is a pure liquid at standard pressure, p° . The star refers to pure liquid and a_{w} is the activity of pure liquid water. Standard pressure used in this article is 101.325 kPa. As far as condensed phases are considered the thermodynamic properties are practically equal at 101.325 kPa (1 atm) and 100 kPa (1 bar) pressure.

Received: March 28, 2018 Accepted: July 11, 2018 Published: July 27, 2018

Activity is always related to chosen standard state so we obtain for activity of pure liquid water:

$$RT \ln(a_{w}^{*}) = \int_{p^{o}}^{p} v_{w}^{*} dp$$
(3)

where v_w is the molar volume of liquid water.

So, for the chemical potential of liquid water in solution we obtain

$$\mu_{w}(T, p) = \mu_{w}^{o}(T, P) + RT \ln(a_{w}^{s})$$
$$= \left[\mu_{w}^{o}(T) + \int_{p^{o}}^{p} v_{w}^{*} dp\right] + RT \ln(a_{w}^{s})$$
(4)

Similarly for the chemical potential of ice:

$$\mu_{\rm ice}^{*}(T, p) = \mu_{\rm ice}^{\rm o}(T) + \int_{p^{\rm o}}^{p} \nu_{\rm ice}^{*} \, \mathrm{d}p \tag{5}$$

Equilibrium between Liquid Water in a Solution and Ice. When supercooled liquid water is in equilibrium with pure solid ice the chemical potentials of water are equal:

$$\mu_{w}(T, p) = \mu_{ice}^{*}(T, p) \Leftrightarrow \mu_{ice}^{o}(T) + \int_{p^{o}}^{P} \nu_{ice}^{*} dp$$
$$= \mu_{w}^{o}(T) + \int_{p^{o}}^{P} \nu_{w}^{*} dp + RT \ln(a_{w}^{s})$$
(6)

Thus, solving activity for supercooled water in a solution in equilibrium with ice, yields

$$RT \ln(a_{w}^{s}) = \mu_{ice}^{o}(kT) + \int_{p^{o}}^{p} v_{ice}^{*} dp - \mu_{w}^{o}(T) - \int_{p^{o}}^{p} v_{w}^{*} dp$$
$$= -\Delta G^{o} - \int_{p^{o}}^{p} \Delta v dp$$
(7)

where

$$\Delta G^{\circ} = \mu_{\rm w}^{\rm o}(T) - \mu_{\rm ice}^{\rm o}(T) \tag{8}$$

$$\Delta v^* = v_{\rm w}^* - v_{\rm ice}^* \tag{9}$$

At standard pressure p° the integral term is zero so

$$RT \ n(a_w^s) = -\Delta G^\circ = RT \ \ln K \Rightarrow a_w^s = K$$
(10)

 Table 1. Heat Capacity Data for Liquid Water Used in the

 Assessment

authors	temp range (K)	no. expts included	excluded	weight
NIST ²²	273.15- 373.756	100	T > 374 K	1
Wagner and Pruss ²³	274-372	50	$T \geq 374 \; K$	1
Speedy ¹⁵	236.05- 257.05	8	236.05 K	0.5
Angel et al., ^{24a}	236-270	9	236 K	0.5
Angel et al. ^{19,20,24b}	236-290	0	all values	0
Archer and Carter ¹²	235-285	0	all values	0
$a_{\rm C}$ (11)	d · 77 11 1	by 1 1		F 1 1 1

"Suggested values in their Table 1. "Measured values in their Table 1 (ref 24).

Table 2. Fitted Parameters for Heat Capacity of Water C_p

Table 3. Quality of the Fit for Heat Capacity of Water C_{ν}

authors	no. expts included	RMSE J/Kmol	AARD %
NIST ²²	100	0.011	0.008
Wagner and Pruss ²³	50	0.011	0.008
Speedy ¹⁵	8	0.53	0.50
Angel et al. ²⁴	9	0.57	0.54

Thus the activity of supercooled water in equilibrium with ice at standard pressure can be calculated if the Gibbs energy change or the equilibrium constant for reaction $H_2O(ice) = H_2O(l)$ is known. Moreover, it is independent of the electrolytes in the solution.

The Gibbs energy change can be calculated from $\Delta H_{\text{fus}}^{\circ}$ and $\Delta C_{\text{p}}^{\circ}$ where $\Delta H_{\text{fus}}^{\circ}$ is enthalpy of fusion and $\Delta C_{\text{p}}^{\circ}$ is the heat capacity change between liquid water and solid ice $\Delta C_{p}^{\circ} = C_{p,\text{w}}^{\circ} - C_{p,\text{ice}}^{\circ}$.

$$\Delta G^{\circ} = \Delta H^{\circ} - T \Delta S^{\circ} \tag{11}$$

where

$$\Delta H^{\circ}(T) = \Delta H^{\circ}(T_{\text{fus}}) + \int_{T_{\text{fus}}}^{T} \Delta C_{p}^{o} \, \mathrm{d}T$$
(12)

and

$$\Delta S^{\circ}(T) = \frac{\Delta H^{\circ}(T_{\text{fus}})}{T_{\text{fus}}} + \int_{T_{\text{fus}}}^{T} \frac{\Delta C_{p}^{\circ}}{T} \, \mathrm{d}T$$
(13)

Below 273.15 K at ambient pressure, the heat capacity of ice is well-known but the heat capacity of supercooled water is more problematic, due to the metastability of liquid water. The heat capacity of supercooled water is showing λ -transition-like behavior so we have applied the approach used in a higher temperature system, that is, equation developed by Hillert and Jarl¹¹

$$C_{p,w}^{o} = 2K_{\rm A}R\left(\tau^{-5} + \frac{\tau^{-15}}{3} + \frac{\tau^{-25}}{5}\right)$$
(14)

where $\tau = 1 - T/T_c$ and T_c is the critical temperature and K_A is a parameter specific to the system.

Pressure Ratio between Pure Supercooled Liquid Water and Ice. The chemical potential of pure water vapor is

$$\mu_{g}^{*}(T, p) = \mu_{g}^{\circ}(T) + RT \ln(p^{*}/p^{\circ}) + RT \ln(\phi^{*})$$
(15)

where the standard state for vapor is ideal gas, p is vapor pressure and ϕ refers to fugacity coefficient, which can be evaluated in ambient pressure with the second virial coefficient B:

$$RT \ln(\phi^*) = B^*(T)p \tag{16}$$

In equilibrium between ice and vapor the chemical potentials are equal and we obtain from eqs 5, 15, and 16:

$$\mu_{ice}^{o}(T) + \int_{p^{o}}^{p_{ice}^{*}} \nu_{ice}^{*} dp$$

= $\mu_{g}^{o}(T) + RT \ln(p_{ice}^{*}/p^{o}) + B^{*}(T)p_{ice}^{*}$ (17)

range	temp range (K)	K_{A}	A_i	$B_i (\mathrm{K}^{-1})$	$D_i \left(\mathrm{K}^2 \right)$
1	237-262.15	5.4943	-255.07	1.07493	
2	262.15-298.15		134.4	-0.385856	6.29422×10^{-4}
3	298.15-373.15		89.8098	-0.09426775	1.53047×10^{-4}

2987



Figure 1. Assessed heat capacity C_p of supercooled water (line) as a function of temperature compared with experimental data^{12,15,24,25} (dots) at subzero temperatures. Dashed line presents extrapolated values. Equation used in freezing point modeling by Thomsen et al.²⁶ is also shown in the graph.

Similarly for pure water and vapor from eqs 2, 3, 15, and 16

$$\mu_{w}^{o}(T) + \int_{p^{o}}^{p_{w}^{*}} v_{w}^{*} dp$$

= $\mu_{g}^{o}(T) + RT \ln(p_{w}^{*}/p^{o}) + B^{*}(T)p_{w}^{*}$ (18)

Combining eqs 17 and 18 yields

$$\mu_{w}^{o}(T) - \mu_{ice}^{o}(T) + \int_{p^{o}}^{p_{w}^{*}} v_{w}^{*} dp - \int_{p^{o}}^{p_{ice}^{*}} v_{ice}^{*} dp$$
$$= RT \ln(p_{w}^{*}/p_{ice}^{*}) + B^{*}(T)(p_{w}^{*} - p_{ice}^{*})$$
(19)

Up to moderate pressure, molar volumes of condensed phases are insensitive to pressure so

$$\mu_{w}^{o}(T) - \mu_{ice}^{o}(T) + \nu_{w}^{*}p_{w}^{*} - \nu_{ice}^{*}p_{ice}^{*} - p^{o}(\nu_{w}^{*} - \nu_{ice}^{*})$$
$$= RT \ln(p_{w}^{*}/p_{ice}^{*}) + B^{*}(T)(p_{w}^{*} - p_{ice}^{*})$$
(20)

Defining $\Delta p = (p_w - p_{ice})$ yields

$$\mu_{w}^{o}(T) - \mu_{ice}^{o}(T) + (p_{ice}^{*} - p^{o})\Delta v^{*} + v_{ice}^{*}\Delta p^{*} + \Delta v^{*}\Delta p^{*}$$
$$= RT \ln(p_{w}^{*}/p_{ice}^{*}) + B^{*}(T)\Delta p^{*}$$
(21)

and furthermore

$$\frac{\Delta G^{o}}{RT} + (p_{ice}^{*} - p^{o})\frac{\Delta v^{*}}{RT} + (v_{ice}^{*} - B^{*}(T))\frac{\Delta p^{*}}{RT} + \frac{\Delta v^{*} \Delta p^{*}}{RT} = \ln(p_{w}^{*}/p_{ice}^{*})$$
(22)

Solving p_w/p_{ice} ratio yields

$$(p_{\rm w}^*/p_{\rm ice}^*) = \frac{Q_1^*Q_2^*Q_3}{K}$$
(23)

$$Q_{1} = \exp\left((p_{ice}^{*} - p^{\circ})\frac{\Delta\nu^{*}}{RT}\right),$$

$$Q_{2} = \exp\left((\nu_{ice}^{*} - B^{*}(T))\frac{\Delta p^{*}}{RT}\right), Q_{3} = \exp\left(\frac{\Delta\nu^{*}\Delta p^{*}}{RT}\right)$$
(24)

Assuming all Q_i 's equal 1 yields for the pressure of liquid water:

$$p_{\rm w}^* \approx p_{\rm ice}^*/K \tag{25}$$

Thus, the vapor pressure of pure supercooled water can be estimated from vapor pressure of pure ice, if the value of equilibrium constant is known as a function of temperature.

COMPUTATIONAL METHODS AND RESULTS

Heat Capacity of Liquid Water. The heat capacity of liquid water in the temperature range -35 to +100 °C was modeled from experimental data using three temperature ranges. Heat capacity for supercooled water was modeled using the equation of λ transition by Hillert and Jarl.¹¹ During the assessment, it was found that the following equations will describe the heat capacity of liquid water when a baseline, $A_1 + B_1T$, to lambda transition is added.

$$C_{p,w}^{o} = 2K_{A}R\left(\tau^{-5} + \frac{\tau^{-15}}{3} + \frac{\tau^{-25}}{5}\right) + A_{1} + B_{1}T$$

for 237 K < T ≤ 262.15 K (26)

$$C_{p,w}^{o} = A_2 + B_2 T + D_2 T^2$$

for 262.15 K < T \leq 298.15 K (27)

$$C_{p,w}^{o} = A_3 + B_3 T + D_3 T^2$$
 for 298.15 K < T \leq 373.15 K
(28)

Moreover, it was found that the heat capacity data of Archer and $Carter^{12}$ was not at lower temperatures in agreement with the

where

Journal of Chemical & Engineering Data

other literature data so it was not included in the assessment. A similar conclusion was made by Holten et al. 13 in 2014.

The value of critical temperature, T_c , in the literature varies from 227 K to 228 K.^{13–15} The value of 228 K is chosen

Table 4. Heat Capacity Data of Ice and the ObtainedCorrelation Coefficient R

authors	temp range/K	no. expts	weight	R
Gurvich et al. 1989 ²⁷	170-270	12	1	0.99998
Feistel and Wagner 1996 ²⁸	170-273	12	1	0.99994
Giauque and Stout 1936 ²⁹	236.05-257.05	11	1	0.99999

to retain consistency with the HKF model.¹⁶ The experimental data used are shown in Table 1 and the obtained parameters are in Table 2. The C_p data by Anisimov et al.¹⁷ were not used in the assessment because of their narrow temperature interval below 273.15 K. Also, the early data of Rasmussen and MacKenzie,¹⁸ Angell and Tucker,¹⁹ Angell et al.,²⁰ and Oguni and Angell²¹ were not used.

The quality of the assessments for each experimental data set is estimated by standard deviations (SD), which describes the absolute deviation, also known as root-mean-square-error (RMSE), defined as



Figure 2. Assessed heat capacity C_p of solid ice (black line) as a function of temperature compared with experimental data^{28,29} (dots) and equations in the literature^{12,30} (red and blue lines).



Figure 3. Assessed heat capacity change, ΔC_{ρ}° , as a function of temperature compared with the equation of Thurmond and Brass.⁷

Table 5. Values for Equilibrium Constant *K* for the Reaction H_2O (ice) = H_2O (l) at Temperatures -40-0 °C

<i>t</i> (°C)	this work	Spencer et al. ³²	difference
-40	0.6799	0.6805	-0.00057
-35	0.7120	0.7123	-0.0003
-30	0.7467	0.7469	-0.0002
-25	0.7837	0.7840	-0.0003
-20	0.8229	0.8232	-0.0003
-15	0.8641	0.8643	-0.0002
-10	0.9073	0.9074	-0.0001
-5	0.9526	0.9525	0.0001
0	1.0000	1.0002	-0.0002

 Table 6. Density Data for Liquid Water Used in the

 Assessment

authors	temp. range	no. expts included	excluded	weight
Hare and Sorensen ³³	(−34−0) °C	35		1
Wagner and Pruss ²³	(273.15– 373.124) K	18	T≥ 374 K	1
RMSE =	$\sqrt{\sum_{i=1}^{N} \frac{(C_i - E_i)^2}{N}}$			(29)

where *i* goes over all included experimental points (*N*), and C_i and E_i are the calculated and experimental values, respectively. Mean absolute percentage error (MAPE), also known as absolute average relative deviation (AARD %), is used to describe the relative deviation:

AARD% =
$$\frac{100}{N} \left(\sum_{i=1}^{N} \frac{|C_i - E_i|}{E_i} \right)$$
 (30)

The RMSE and AARD% values of different authors used in the assessment are listed in Table 3.

A comparison with the fitted and literature values at subzero temperatures is shown in Figure 1.

Heat Capacity of Ice. The heat capacity of solid ice in the temperature range 170-273.15 K was fitted using the relationship $a + b \cdot (T/K)$ from the experimental data listed in Table 4.

The obtained equation for heat capacity of ice in the temperature range 170-270 K is

$$C_{p,\text{ice}}(J/\text{Kmol}) = 2.1128 + 0.130484 \cdot (T/\text{K})$$
 (31)

and it is compared with experimental data and the other available C_p equations in Figure 2.

Heat Capacity Change. The heat capacity change ΔC_p° can now be evaluated and assessed in the temperature range (237–273.15) K as

$$\Delta C_{\rm p}^{\circ}(\rm J/\rm Kmol) = -19656.303 + 98.468097(T/\rm K) + 2.3432088^{\circ}10^{8} \cdot (T/\rm K)^{-2} - 0.1386227 \cdot (T^{2})$$
(32)

The fitted ΔC_p° compared with the equation by Thurmond and Brass⁷ is presented in Figure 3.

 ΔG° and *K* for the reaction H₂O(ice) = H₂O(I). The value of 6009.5 J/mol is used for heat of fusion of ice at reference temperature 273.15 K.³¹

From eqs 12 and 32 we can evaluate for heat of fusion of ice as a function of temperature:

$$\Delta H^{\circ}(J/mol) = 3501286.79 - 19656.303(T/K) + 49.2340485(T/K)^{2} - 234320890(T/K)^{-2} - 0.046207566(T/K)^{3}$$
(33)



Figure 4. Assessed density ρ of liquid water compared with experimental data.^{15,23,33,34} Dotted lines indicate extrapolated values. Data by Speedy¹⁵ and Lind and Trusler³⁴ were not included in the assessment.



Figure 5. Calculated molar volume ν of liquid water compared with literature data.^{15,23,33}





Thus, the following equations will be obtained from eqs 11-13, and 32:

$$\Delta G^{\circ} = 3501286.89 - 109795.687T + 19656.303T \ln(T)$$

- 49.2340485T² + 0.023103783T³ - 117160445/T (34)
$$\ln(\mathbf{K}) = -421105.608/T + 13205.3106$$

- 2364.09638 ln(T) + 5.92146122T
- 0.0027787306T² + 14091079/T² (35)

A comparison between the equilibrium constant of this work (K) with the equilibrium constant obtained in a thermodynamic modeling of the Na-K-Ca-Mg-Cl-SO₄-H₂O system at temperatures below 25 °C according to Spencer et al.³² in 1990, is displayed in Table 5.

As can be seen from Table 5, the difference in equilibrium constant values is less than -0.0006.

Volumetric Properties. *Liquid Water.* The density for supercooled water was assessed from the recent data of Hare and Sorensen,³³ and Wagner and Pruss.²³ Instead of a sixth degree



Figure 7. Assessed density ρ of solid ice compared with the model of Feistel and Wagner.²⁸





polynomial equation used by Hare and Sorensen,³³ a similar equation by Speedy¹⁵ with one additional parameter *A* was used so that density could be fitted in the entire temperature range -34 to +100 °C.

Thus, the equations for density ρ , molar volume ν , and thermal expansion coefficient α of water are

$$\rho = \rho_o \exp\{-T_c(A + B\varepsilon + 2C\varepsilon^{1/2})\}$$
(36)

$$\nu = \frac{M_{\text{H2O}}}{\rho} = \frac{M_{\text{H2O}}}{\rho_{\text{o}}} \exp\{T_{\text{c}}(A + B\varepsilon + 2C\varepsilon^{1/2})\}$$
(37)

$$\alpha = -\frac{1}{\rho} (\mathrm{d}\rho/\mathrm{d}T)_p = B + C\varepsilon^{-1/2}$$
(38)

where ρ_{o} , A, B, and C are parameters, T is temperature in Kelvin. $M_{\rm H2O}$ is molecular weight of water, and ε is defined as $T/T_{\rm c} - 1$. The data used in the assessment are shown in Table 6.



Figure 9. Vapor pressure of ice p_{ice} according to various authors.^{28,36–38}

Table 7. Volume Related Properties of Supercooled Water and Ice at Subzero Temperatures T^a

T (°C)	T(K)	$\nu_{\rm w} ({\rm cm^3/mol})$	$\nu_{\rm ice}~({\rm cm^3/mol})$	Δu (cm ³ /mol)
0	273.15	18.015	19.651	-1.636
-5	268.15	18.027	19.637	-1.610
-10	263.15	18.049	19.622	-1.573
-15	258.15	18.084	19.607	-1.523
-20	253.15	18.135	19.593	-1.457
-25	248.15	18.208	19.578	-1.370
-30	243.15	18.312	19.563	-1.251
-35	238.15	18.464	19.549	-1.085
-40	233.15	18.710	19.534	-0.824
-45	228.15	19.396	19.520	-0.123

 ${}^{a}\nu_{\rm w}$ is molar volume of liquid water, $\nu_{\rm ice}$ is molar volume of ice and $\Delta\nu = \nu_{\rm w} - \nu_{\rm ice}$.

Obtained parameters for density for liquid water in the temperature range -34 to 100 °C are

$$\rho_{\rm o} = 1.007853 ({\rm g/cm}^3), \quad A = 3.9744 \cdot 10^{-04} (1/{\rm K}),$$

 $B = 1.6785 \cdot 10^{-03} (1/{\rm K}) \quad \text{and}$
 $C = -7.816510^{-04} (1/{\rm K})$

RMSE and AARD values for Hare and Sorensen's³³ data were 0.0001 g/cm³ and 0.010%, respectively. For the data by Wagner and Pruss,²³ the corresponding values were 0.0002 g/cm³ and 0.020%, respectively.

The assessed density of liquid water compared with the experimental data^{15,23,33,34} is presented in Figure 4, and the calculated molar volume is in Figure 5.

After completing the density assessment it was noticed that the calculated thermal expansion coefficient are in good agreement up to 100 °C when compared to thermal expansion coefficient data by Hare and Sorensen³³ and Wagner and Pruss.²³ The thermal expansion coefficient from the data by Wagner and

Table 8. Pressure Related Properties at SubzeroTemperatures T^a

$T(^{\circ}C)$	T(K)	K	$p_{\rm w}$ (Pa)	$p_{\rm ice}$ (Pa)	$\Delta p(Pa)$	$B (cm^3/mol)$
0	273.15	1.000	611.15	611.15	0.00	-1116
-5	268.15	0.953	401.76	421.74	19.98	-1226
-10	263.15	0.907	259.89	286.44	26.55	-1345
-15	258.15	0.864	165.29	191.29	26.00	-1472
-20	253.15	0.823	103.25	125.48	22.23	-1608
-25	248.15	0.784	63.28	80.75	17.46	-1752
-30	243.15	0.747	38.01	50.90	12.89	-1904
-35	238.15	0.712	22.35	31.39	9.04	-2064
-40	233.15	0.680	12.84	18.89	6.05	-2233
-45	228.15	0.651	7.21	11.07	3.87	-2410

"Notation: K, equilibrium constant; p_{w} , vapour pressure of supercooled water; p_{ice} , vapour pressure of ice. $\Delta p = p_w - p_{ice}$ and B is the second virial coefficient of water vapour. Vapour pressure of supercooled water is calculated using equation 25.

Pruss²³ was calculated using eq 38 in which the partial derivate of density at standard pressure was calculated by FluidCal.³⁵

As can been seen from Figures 4–6, the assessed density and thermal expansion coefficient derived from it, agree well with the literature data and the obtained equations have good extrapolation capabilities especially to lower temperatures.

Ice lh. The density of ice was fitted from the data of Feistel and Wagner²⁸ in the temperature range (230-273.15) K. A linear fit was found satisfactory as can been seen in Figure 7, with linear correlation coefficient 0.9998 obtained. Thus, the following equations are obtained for ice:

$$\rho_{\rm ice} = A + BT \tag{39}$$

$$\nu_{\rm ice} = \frac{M_{\rm H2O}}{\rho_{\rm ice}} = \frac{M_{\rm H2O}}{A + BT} \tag{40}$$

where A = 0.954205 and B = -0.0001371 (K⁻¹).

Table 9. Values of Q Parameters and Correlated Pressure (p_w) of Supercooled Water Calculated by Equation 23^{*a*}

T (°C)	T (K)	O_1	O_2	<i>O</i> ₃	$p_{w corr}$ (Pa)	Δp (Pa)
0	272.15	1 0001	1 0000	1 0000	611 20	0.04
0	2/3.13	1.0001	1.0000	1.0000	011.20	0.04
-5	268.15	1.0001	1.0000	1.0000	421.71	0.04
-10	263.15	1.0001	1.0000	1.0000	286.38	0.03
-15	258.15	1.0001	1.0000	1.0000	191.23	0.02
-20	253.15	1.0001	1.0000	1.0000	125.42	0.01
-25	248.15	1.0001	1.0000	1.0000	80.70	0.01
-30	243.15	1.0001	1.0000	1.0000	50.86	0.00
-35	238.15	1.0001	1.0000	1.0000	31.36	0.00
-40	233.15	1.0000	1.0000	1.0000	18.87	0.00
-45	228.15	1.0000	1.0000	1.0000	11.06	0.00
az 7		c	11.	. 1	1	1.0

^aVapour pressure of supercooled water in calculation of Q_2 and Q_3 parameters is obtained by using equation 25.

Volume Difference between Supercooled Water and Solid lce. Molar volume difference between supercooled water and solid ice can now be calculated from equation $\Delta \nu = \nu_{\rm w} - \nu_{\rm ice}$. Obtained values are shown in Figure 8. It is interesting to find out that the volume difference disappears when temperature approaches the critical temperature 228 K.

VAPOR PRESSURE

Murphy and Koop³⁶ have derived the following equation for the vapor pressure of ice when temperature is over 110 K:

$$\ln(p_{\rm ice}^*/Pa) = b_0 + b_1/T + b_2 \ln(T/K) + b_3T$$
(41)

where $b_0 = 9.550426$, $b_1 = -5723.265$ K, $b_2 = 3.53068$, and $b_3 = -0.00728332$ K⁻¹. Analytical eq 41 with a comparison with the literature data is presented in Figure 9.

As can been seen, the equation for the fitted vapor pressure of ice by Murphy and Koop³⁶ is in good agreement with the

literature data. Assuming all Q_i 's are equal to 1 in eq 23 we end up with eq 25 as

$$p_{\rm w}^* \approx p_{\rm ice}^* / K \tag{25}$$

To test the validity of eq 25, the second virial coefficient at subzero temperatures is needed. No experimental data is available at subzero temperatures; so the second virial coefficient B was extrapolated from the values by Wagner and Pruss²³ in the temperature range 273.15–323.15 K. The obtained equation with the correlation coefficient value of 0.997 is

$$B(cm3/mol) = -19362.33 + 112.334 \cdot (T/K) - 0.1667 \cdot (T/K)2$$
(42)

Volume related properties of supercooled water and ice in the temperature range -45-0 °C are listed in Table 7, pressure related properties in Table 8, and Q parameters with corrected vapor pressure of supercooled water in Table 9.

As can be seen in Table 9, the pressure correction does not exceed 0.04 Pa and so eq 25 is an excellent approximation for modeling purposes for aqueous solutions at subzero temperatures.

The calculated vapor pressure of pure supercooled water with the literature data is shown in Figure 10 and the pressure difference between pure supercooled water and ice in Figure 11. The calculated vapor pressure of pure supercooled water compared to the values obtained with the equation by Murphy and Koop³⁶ is presented in Figure 12. The difference with the equation by Murphy and Koop is less than 0.07 Pa using eq 25 and 0.03 Pa using eq 23.

Activity of Supercooled Water on Ice Curve. The calculated water activity along the ice-curve is presented in Table 10 and compared with the literature data in Figure 13.



Figure 10. Calculated vapor pressure of pure supercooled water p_w compared with selected literature data.^{37,39-41} Not all data by Kraus and Greer⁴¹ is shown on the graph.

ΔP



Figure 11. Calculated vapor pressure difference $\Delta p = p_w - p_{ice}$ between pure supercooled water and ice compared with the literature data.^{36,37}



Figure 12. Difference in calculated vapor pressure of pure supercooled water, $\Delta p = p_{\text{thiswork}} - p_{\text{Koop}}$, compared with the equation presented by Murphy and Koop.³⁶ The diamonds (blue) were calculated with eq 25 and squares (red) with eq 23.

As can been seen from Figure 13, the activity of water on ice curve is predicted within ± 0.001 down to 237 K if values by Murphy and Koop are neglected.

DISCUSSION AND CONCLUSIONS

A simple model for thermodynamic properties of water from subzero temperatures up to 373 K was derived at ambient

pressure. The heat capacity of supercooled water was assessed as lambda transition. The obtained properties for supercooled water such as heat capacity, vapor pressure, density, and thermal expansion are in excellent agreement with literature data.

Thermodynamic description for the ice-curve in electrolyte solutions is generally obtained in modeling the solubility of ice in the studied electrolyte solution. Thus, the thermodynamic

T (K)	a _w	Т (К)	a _w	Т (К)	a _w	T (K)	a _w
273.15	1.0000	263.15	0.9073	253.15	0.8229	243.15	0.7467
272.15	0.9904	262.15	0.8985	252.15	0.8149	242.15	0.7396
271.15	0.9808	261.15	0.8898	251.15	0.8070	241.15	0.7326
270.15	0.9713	260.15	0.8811	250.15	0.7991	240.15	0.7256
269.15	0.9619	259.15	0.8726	249.15	0.7914	239.15	0.7188
268.15	0.9526	258.15	0.8641	248.15	0.7837	238.15	0.7121
267.15	0.9434	257.15	0.8557	247.15	0.7762	237.15	0.7054
266.15	0.9342	256.15	0.8473	246.15	0.7687	236.15	0.6989
265.15	0.9252	255.15	0.8391	245.15	0.7613	235.15	0.6925
264.15	0.9162	254.15	0.8309	244.15	0.7540	234.15	0.6862





Temperature K

Figure 13. Calculated activity of supercooled water on ice curve compared to activity using eq 10 to equilibrium constants by Murphy and Koop³⁶ and Spencer et al.³² as well as polynomial equation with six terms by Carslaw et al.⁴² and Toner et al.¹

Table 11. Values of Debye Hückel Constants. An Outlying Value Obtained by Extrapolating the Equation Adopted by Møller et al.⁴³ at -10° C Is Underlined

t °C	MTDATA ⁴⁴	FactSage ⁴⁵	Møller et al. 1988 ⁴³	Archer and Wang 1990 ⁴⁶	Spencer et al. 1990 ³²
-30	0.3639	0.3640	0.3636	0.3535	0.3687
-25	0.3657	0.3658	0.3653	0.3599	0.3698
-20	0.3677	0.3678	0.3671	0.3644	0.3711
-15	0.3697	0.3698	0.3689	0.3679	0.3725
-10	0.3719	0.3720	0.3867	0.3709	0.3742
-5	0.3742	0.3743	0.3744	0.3737	0.376
0	0.3767	0.3768	0.3767	0.3764	0.3781
5	0.3793	0.3795	0.3793	0.3792	0.3804

properties of the ice curve are dependent on the studied electrolyte systems and their number, the excess model used for nonideal behavior of aqueous solution, and the Debye–Hückel constant used. The values of the Debye–Hückel (DH) equation used varies significantly at subzero temperatures from each other, as can been in Table 11. Sometimes the terms used in the DH equation such as 1/(T - 263 K) and 1/(628 K - T) produce ambiguous values, as noted by Spencer et al.³²

As can be seen from Table 11, there is a variation between the different DH equations and it increases at lower temperatures. Variation can be expected to increase when first and second derivatives are calculated. Thus, modeled heat capacity of pure supercooled water will depend on the DH equation used if it is obtained by thermodynamic modeling of the apparent heat capacity data.

We suggest a converse procedure. Our equations for heat capacity of supercooled water eqs 26-28 and heat capacity change eq 32 as well as activity of water on ice curve (see Table 10) should be used as the basis for thermodynamic modeling at subzero temperatures. So, the activity of water on the ice curve will form a uniform basis for all electrolyte solutions.

Table 12. Heat Capacity of Supercooled and Ordinary Water up to 373.15 K (100 °C) in 1 atm Pressure. C_p° (J/Kmol) = $A + BT + CT^{-2} + D/T^2$

range	temp range (K)	A_i	B_i (K ⁻¹)	$C_i (\mathrm{K}^{-2})$	$D_i \left(\mathrm{K}^2 ight)$
1	237-262.15	-19654.2	98.5986	-0.138623	2.343209×10^{8}
2	262.15-298.15	134.4	-0.385856	6.29422×10^{-4}	
3	298.15-373.15	89.8098	-0.09426775	1.53047×10^{-4}	

Article

Journal of Chemical & Engineering Data

Several substances form hydrates at subzero temperatures so the activity of water will be included in the solubility product of the precipitated hydrate. Thus, the solubility is connected with the thermodynamic properties of ice and supercooled water via the used activity coefficient model. If the heat capacity of precipitated hydrate is measured from subzero temperature up to 298.15 K, a link between the critically evaluated thermodynamic data and thermodynamic properties of water is formed which enables a critical evaluation of thermodynamic properties of the hydrate.

Generally, heat capacity in thermodynamic and engineering software are expressed as a polynomial equation. Combining heat capacity data of ice and heat capacity change data between ice and supercooled water, these parameters for polynomial equations are obtained (Table 12).

AUTHOR INFORMATION

Corresponding Author

*E-mail: hannu.sippola@aalto.fi.

ORCID 0

Hannu Sippola: 0000-0001-8137-4677

Present Address

[#]H.S.: FCG Design and Engineering, Osmontie 34, FI-00601 Helsinki, Finland.

Funding

A research grant received from K.H. Renlund Foundation is kindly acknowledged.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Fruitful cooperation with Prof. Mika Järvinen of Aalto University, School of Engineering, in the framework of steel slag recycling is also kindly appreciated.

REFERENCES

(1) Toner, J. D.; Catling, D. C.; Light, B. A revised Pitzer model for low-temperature soluble salt assemblages at the Phoenix site, Mars. *Geochim. Cosmochim. Acta* **2015**, *166*, 327–343.

(2) Holten, V.; Bertrand, C. E.; Anisimov, M. A.; Sengers, J. V. Thermodynamics of supercooled water. *J. Chem. Phys.* **2012**, *136*, 094507.

(3) Debenedetti, P.; Stanley, E. Supercooled and glassy water. *Phys. Today* **2003**, *56*, 40–46.

(4) Mallamace, F.; Corsaro, C.; Stanley, H. E. A singular thermodynamically consistent temperature at the origin of the anomalous behavior of liquid water. *Sci. Rep.* **2012**, *2*, 993.

(5) Nilsson, A.; Pettersson, L. G. M. The structural origin of anomalous properties of liquid water. *Nat. Commun.* **2015**, *6*, 8998.

(6) Stanley, H. E.; Buldyrev, S. V.; Mishima, O.; Sadr-Lahijany, M. R.; Scala, A.; Starr, F. W. Unsolved mysteries of water in its liquid and glassy phases. *J. Phys.: Condens. Matter* **2000**, *12*, A403.

(7) Thurmond, V. L.; Garrett, B. W. Geochemistry of freezing brines Low-temperature properties of sodium chloride. *CRREL Report* **1987**, 87-13, 1-16.

(8) Koop, T.; Luo, B.; Tsias, A.; Thomas, P. Water activity as the determinant for homogeneous ice nucleation in aqueous solutions. *Nature* **2000**, *406*, 611–614.

(9) Zobrist, B.; Marcolli, C.; Pete, T.; Koop, T. Heterogeneous Ice Nucleation in Aqueous Solutions: the Role of Water Activity. *J. Phys. Chem. A* **2008**, *112*, 3965–3975.

(10) Pátek, J.; Hrubý, J.; Klomfar, J.; Součková, M.; Harvey, A. H. Reference Correlations for Thermophysical Properties of Liquid Water at 0.1 MPa. *J. Phys. Chem. Ref. Data* **2009**, *38*, 21–29.

(11) Hillert, M.; Jarl, M. A model for alloying in ferromagnetic metals. *CALPHAD: Comput. Coupling Phase Diagrams Thermochem.* **1978**, *2*, 227–238.

(12) Archer, D. G.; Carter, R. W. Thermodynamic Properties of the NaCl + H2O System. 4. Heat Capacities of H2O and NaCl(aq) in Cold-Stable and Supercooled States. *J. Phys. Chem. B* **2000**, *104*, 8563–8584.

(13) Holten, V.; Sengers, J. V.; Anisimov, M. A. Equation of State for Supercooled Water at Pressures up to 400 MPa. *J. Phys. Chem. Ref. Data* **2014**, *43*, 043101.

(14) Speedy, R. J.; Angell, C. A. Isothermal compressibility of supercooled water and evidence for a thermodynamic singularity at -45° C. J. Chem. Phys. **1976**, 65, 851–858.

(15) Speedy, R. J. Thermodynamic properties of supercooled water at 1 atm. J. Phys. Chem. **1987**, *91*, 3354–3358.

(16) Shock, E. L.; Helgeson, H. C. Calculation of the thermodynamic and transport properties of aqueous species at high pressures and temperatures: Correlation algorithms for ionic species and equation of state predictions to 5 kb and 1000°C. *Geochim. Cosmochim. Acta* **1988**, *52*, 2009–2036.

(17) Anisimov, M. A.; Voronel', A. V.; Zaugol'Nikova, N. S.; Ovodov, G. I. Specific Heat of Water Near the Melting Point and Ornstein-Zernike Fluctuation Corrections. *Journal of Experimental and Theoretical Physics Letters* **1972**, *15*, 317–319.

(18) Angell, C. A.; Tucker, J. C. Anomalous heat capacities of supercooled water and heavy water. *Science* **1973**, *181*, 342–344.

(19) Angell, C. A.; Tucker, J. C. Anomalous Heat Capacities of Supercooled Water and Heavy Water. *Science* **1973**, *181*, 342.

(20) Angell, C. A.; Shuppert, J.; Tucker, J. C. Anomalous properties of supercooled water. Heat capacity, expansivity, and proton magnetic resonance chemical shift from 0 to -38%. *J. Phys. Chem.* **1973**, *77*, 3092–3099.

(21) Oguni, M.; Angell, C. A. Heat capacities of H2O+H2O2, and H2O+N2H4, binary solutions: Isolation of a singular component for Cp of supercooled water. *J. Chem. Phys.* **1980**, *73*, 1948–1954.

(22) Thermophysical Properties of Fluid Systems. http://webbook. nist.gov/chemistry/fluid/ (accessed 11.11.2017).

(23) Wagner, W.; Pruss, A. The IAPWS Formulation 1995 for the Thermodynamic Properties of Ordinary Water Substance for General and Scientific Use. J. Phys. Chem. Ref. Data 2002, 31, 387–535.

(24) Angell, C. A.; Oguni, M.; Sichina, W. J. Heat capacity of water at extremes of supercooling and superheating. *J. Phys. Chem.* **1982**, *86*, 998–1002.

(25) Tombari, E.; Ferrari, C.; Salvetti, G. Heat capacity anomaly in a large sample of supercooled water. *Chem. Phys. Lett.* **1999**, 300, 749–751.

(26) Arshad, M. W.; Fosbøl, P. L.; von Solms, N.; Thomsen, K. Freezing Point Depressions of Phase Change CO2 Solvents. *J. Chem. Eng. Data* **2013**, *58*, 1918–1926.

(27) Gurvich, L. V.; Veyts, I. V.; Alcock, C. B. Thermodynamic properties of individual substances 1. Elements O, H(D, T), F, Cl, Br, I, He, Ne, Ar, Kr, Xe, Rn, S, N, P, and their compounds. Part 2 Tables; Hemispere Publishing Corporation: New York, 1989; pp 340.

(28) Feistel, R.; Wagner, W. A New Equation of State for H2O Ice Ih. *J. Phys. Chem. Ref. Data* **2006**, 35, 1021–1047.

(29) Giauque, W. F.; Stout, J. W. The Entropy of Water and the Third Law of Thermodynamics. The Heat Capacity of Ice from 15 to 273 K. J. Am. Chem. Soc. **1936**, *58*, 1144–1150.

(30) Fukusako, S. Thermophysical properties of ice, snow, and sea ice. *Int. J. Thermophys.* **1990**, *11*, 353–372.

(31) Osborne, N. S. Heat of fusion of ice. A revision. J. Res. Natl. Bur. Stand. **1939**, 23, 643–646.

(32) Spencer, R. J.; Moller, N.; Weare, J. H. The prediction of mineral solubilities in natural waters: A chemical equilibrium model for the Na-K-Ca-Mg-Cl-SO4-H2O system at temperatures below 25°C. *Geochim. Cosmochim. Acta* **1990**, *54*, 575–590.

(33) Hare, D. E.; Sorensen, C. M. The density of supercooled water. II. Bulk samples cooled to the homogeneous nucleation limit. *J. Chem. Phys.* **1987**, *87*, 4840–4845.

Journal of Chemical & Engineering Data

(34) Lin, C. -W.; Trusler, J. P. M. The speed of sound and derived thermodynamic properties of pure water at temperatures between (253 and 473) K and at pressures up to 400 MPa. *J. Chem. Phys.* **2012**, *136*, 094511.

(35) Wagner, W. Software for the Calculation of Thermodynamic Properties for a Great Number of Substances, FLUIDCAL. http:// www.thermo.ruhr-uni-bochum.de/en/prof-w-wagner/software/ fluidcal.html (accessed 18.7.2013).

(36) Murphy, D. M.; Koop, T. Review of the vapour pressures of ice and supercooled water for atmospheric applications. *Q. J. R. Meteorol. Soc.* **2005**, *131*, 1539–1565.

(37) Scheel, K.; Heuse, W. Bestimmung des Sättigungsdruckes von Wasserdampf unter 0°. *Ann. Phys.* **1909**, *334*, 723–737.

(38) Fernicola, V.; Rosso, L.; Giovannini, M. Investigation of the Ice-Water Vapor Equilibrium Along the Sublimation Line. *Int. J. Thermophys.* **2012**, *33*, 1363–1373.

(39) Fukuta, N.; Gramada, C. M. Vapor Pressure Measurement of Supercooled Water. J. Atmos. Sci. 2003, 60, 1871–1875.

(40) Beltramino, G.; Rosso, L.; Smorgon, D.; Fernicola, V. Vapor pressure measurements over supercooled water in the temperature range from -10^{1} °C to $+10^{-2}$ °C. *J. Chem. Thermodyn.* **2017**, *105*, 159–164.

(41) Kraus, G. F.; Greer, S. C. Vapor pressures of supercooled water and deuterium oxide. *J. Phys. Chem.* **1984**, *88*, 4781–4785.

(42) Carslaw, K. S.; Clegg, S. L.; Brimblecombe, P. A thermodynamic model of the system HCI-HNO3-H2SO4-H2O, including solubilities of HBr, from< 200 to 328 K. J. Phys. Chem. **1995**, *99*, 11557–11574.

(43) Møller, N. The prediction of mineral solubilities in natural waters: A chemical equilibrium model for the Na-Ca-Cl-SO4-H2O system, to high temperature and concentration. *Geochim. Cosmochim. Acta* **1988**, *52*, 821–837.

(44) Davies, R. H.; Dinsdale, A. T.; Gisby, J. A.; Robinson, J. A. J.; Martin, S. M. MTDATA - thermodynamic and phase equilibrium software from the national physical laboratory. *CALPHAD: Comput. Coupling Phase Diagrams Thermochem.* **2002**, *26*, 229–271.

(45) Bale, C. W.; Bélisle, E.; Chartrand, P.; Decterov, S. A.; Eriksson, G.; Gheribi, A. E.; Hack, K.; Jung, I. -H.; Kang, Y. -B.; Melançon, J.; Pelton, A. D.; Petersen, S.; Robelin, C.; Sangster, J.; Spencer, P.; Van Ende, M. FactSage thermochemical software and databases, 2010–2016. *CALPHAD: Comput. Coupling Phase Diagrams Thermochem.* 2016, 54, 35–53.

(46) Archer, D. G.; Wang, P. The Dielectric Constant of Water and Debye-Hückel Limiting Law Slopes. *J. Phys. Chem. Ref. Data* **1990**, *19*, 371–411.

Article