
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Nissinen, J.; Volovik, G. E.

Elasticity tetrads, mixed axial-gravitational anomalies, and (3+1)-d quantum Hall effect

Published in:
PHYSICAL REVIEW RESEARCH

DOI:
[10.1103/PhysRevResearch.1.023007](https://doi.org/10.1103/PhysRevResearch.1.023007)

Published: 06/09/2019

Document Version
Publisher's PDF, also known as Version of record

Published under the following license:
CC BY

Please cite the original version:
Nissinen, J., & Volovik, G. E. (2019). Elasticity tetrads, mixed axial-gravitational anomalies, and (3+1)-d quantum Hall effect. *PHYSICAL REVIEW RESEARCH*, 1(2), Article 023007.
<https://doi.org/10.1103/PhysRevResearch.1.023007>

This material is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Elasticity tetrads, mixed axial-gravitational anomalies, and (3+1)-d quantum Hall effect

J. Nissinen^{1,*} and G. E. Volovik^{1,2}¹Low Temperature Laboratory, Aalto University, P.O. Box 15100, FI-00076 Aalto, Finland²Landau Institute for Theoretical Physics, Acad. Semyonov Av., 1a, 142432, Chernogolovka, Russia

(Received 4 June 2019; revised manuscript received 5 August 2019; published 6 September 2019)

For two-dimensional topological insulators, the integer and intrinsic (without external magnetic field) quantum Hall effect is described by the gauge anomalous (2+1)-dimensional [(2+1)-d] Chern-Simons (CS) response for the background gauge potential of the electromagnetic U(1) field. The Hall conductance is given by the quantized prefactor of the CS term, which is a momentum-space topological invariant. Here, we show that three-dimensional crystalline topological insulators with no other symmetries are described by a topological (3+1)-dimensional [(3+1)-d] mixed CS term. In addition to the electromagnetic U(1) gauge field, this term contains elasticity tetrad fields $E_\mu^a(\mathbf{r}, t) = \partial_\mu X^a(\mathbf{r}, t)$ which are gradients of crystalline U(1) phase fields $X^a(\mathbf{r}, t)$ and describe the deformations of the crystal. For a crystal in three spatial dimensions $a = 1, 2, 3$ and the mixed axial-gravitational response contains three parameters protected by crystalline symmetries: the weak momentum-space topological invariants. The response of the Hall conductance to the deformations of the crystal is quantized in terms of these invariants. In the presence of dislocations, the anomalous (3+1)-d CS term describes the Callan-Harvey anomaly inflow mechanism. The response can be extended to all odd spatial dimensions. The elasticity tetrads, being the gradients of the lattice U(1) fields, have canonical dimension of inverse length. Similarly, if such tetrad fields enter general relativity, the metric becomes dimensionful, but the physical parameters, such as Newton's constant, the cosmological constant, and masses of particles, become dimensionless.

DOI: [10.1103/PhysRevResearch.1.023007](https://doi.org/10.1103/PhysRevResearch.1.023007)

I. INTRODUCTION

The effective Chern-Simons (CS) description of the integer (and fractional [1,2]) quantum Hall effect (IQHE) and the ensuing topological quantization of Hall conductivity has been originally considered in even spatial dimensions [3]. Similarly, in (2+1)-dimensional [(2+1)-d] topological insulators [4], in the generalized QED₃ [5–7] or in thin films of topological superfluids and superconductors [8,9], the intrinsic or anomalous quantum Hall effect (AQHE) in the absence of magnetic flux is also described by a CS term. For both IQHE and AQHE, the prefactor of this term is expressed in terms of momentum-space topological invariants: the Chern number(s). The same mechanism works for gapped systems in all even space dimensions [10–12].

Here, we show that the IQHE/AQHE in (3+1)-dimensional [(3+1)-d] crystalline quantum Hall systems or topological insulators [13] is also described by CS term with mixed field content. Namely, the (3+1)-d CS term features elasticity tetrads [14–17], which describe the geometry of elasticity theory, including crystals with dislocation defects. The density of dislocations corresponds to spatial torsion of the geometry, more familiar in gravitational theories (see, e.g., Ref. [18]). In contrast to the gravitational

tetrads, the elasticity tetrads have canonical dimensions of inverse length. As a result, the CS term is dimensionless (in units $\hbar = 1$), and as in the case of even space dimensions, the prefactor is given by integer momentum-space topological invariants. The CS term leads to the analog of mixed (axial and gravitational/elastic) anomaly inflow in (3+1)-d (see, e.g., Refs. [19–23]). The structure of the CS term reflects the Callan-Harvey mechanism of anomaly cancellation [23], provided here by the fermion zero modes living on dislocations/sample boundaries [24–26]. In this way, the CS response related to the (3+1)-d QHE is valid in the presence of deformations and satisfies the consistency conditions of gauge invariance and anomaly inflow. The same mechanism works for gapped crystalline systems in all odd spatial dimensions.

The rest of this paper is organized in the following way. Before introducing the elasticity tetrads and the ensuing effective response in (3+1)-d in Secs. III and IV, we first review the simpler (2+1)-d QH case. Section V describes the anomaly inflow mechanism of the action in (3+1)-d. In Sec. VI, we briefly describe the extension to arbitrary even space-time dimensions. The coupling of the elasticity tetrads to space-time geometry in QHE is discussed in Sec. VII along with the speculative possibility to have quantized and dimensionless gravitational couplings if the gravitational space-time metric is identified with the metric of elasticity tetrads. We conclude in Sec. VIII.

II. (2+1)-d TOPOLOGICAL ACTION FOR QHE

The topological Chern-Simons action for the IQHE and for the anomalous, intrinsic (i.e., without external magnetic field)

*Corresponding author: jaakko.nissinen@aalto.fi

AQHE in the $D = (2+1)$ -dimensional crystalline insulator is given by [4,6–8]

$$S_{(2+1)\text{-d}}[A_\mu] = \frac{1}{4\pi} N \int d^2x dt \epsilon^{\nu\alpha\beta} A_\nu \partial_\alpha A_\beta, \quad (1)$$

where $|e| = \hbar = 1$ and the electromagnetic U(1) gauge field A_μ has dimensions of momentum. The integer prefactor N in the response is expressed in terms of momentum-space topological invariant [3,6–8]

$$N = \frac{1}{8\pi^2} \epsilon_{ij} \int_{-\infty}^{\infty} d\omega \int_{\text{BZ}} dS \times \text{Tr}[(G\partial_\omega G^{-1})(G\partial_{k_i} G^{-1})(G\partial_{k_j} G^{-1})], \quad (2)$$

where the spatial momentum integral is over the 2-d torus of the two-dimensional Brillouin zone (BZ). The integer N is a topological invariant of the system and in particular remains locally well defined under smooth deformations of the lattice. Under sufficiently strong deformations or disorder one can have regions of different $N(x)$ with the associated chiral edge modes. In that case, the global invariant, if any, is defined by the topological charge of the dominating cluster which percolates through the system [27].

A. (2+1)-d bulk Chern-Simons and consistent boundary anomaly

The topologically protected physics of the quantum Hall effect arises due to the Callan-Harvey [23] anomaly inflow of the Chern-Simons action from the bulk to the boundary [28–30], which we now review. In fact, the boundary current J_{bdry}^μ realizes the (consistent) (1+1)-d chiral anomaly

$$\partial_\mu J_{\text{bdry}}^\mu = \frac{N}{8\pi} \epsilon^{\mu\nu} F_{\mu\nu}^{\text{bdry}}, \quad (3)$$

where $J_{\text{bdry}}^{0,\parallel}, F_{0\parallel}^{\text{bdry}} = \partial_t A_\parallel - \partial_\parallel A_t$ are along the boundary. In this way the protected edge modes arise from the cancellation of bulk and boundary gauge anomalies [7,23,28,29]. In more detail, the (2+1)-d Chern-Simons term (1) can be written as

$$S_{\text{CS}}[A] = \frac{1}{4\pi} \int d^3x \epsilon^{\mu\nu\lambda} \sigma(x) A_\mu \partial_\nu A_\lambda, \quad (4)$$

where $\sigma(x) = N\Theta(x^1)$ has a step function domain wall at $x^1 = 0$. This is equivalent to a space-time manifold with a boundary. The current is

$$j_{\text{H}}^\mu = \frac{\delta S_{\text{CS}}[A]}{\delta A_\mu} = \frac{\sigma}{2\pi} \epsilon^{\mu\nu\lambda} \partial_\nu A_\lambda - \frac{1}{4\pi} \epsilon^{\mu\nu\lambda} (\partial_\nu \sigma) A_\lambda \quad (5)$$

with bulk and boundary contributions. We see that the Hall conductivity is $\sigma_{\text{H}} = Ne^2/h$. The current has the divergence

$$\partial_\mu j_{\text{H}}^\mu = \frac{1}{4\pi} \epsilon^{\mu\nu\lambda} \partial_\mu \sigma \partial_\nu A_\lambda = \frac{N}{8\pi} \delta(x^1) \epsilon^{1\nu\lambda} F_{\nu\lambda} = \frac{N}{4\pi} E^\parallel, \quad (6)$$

where we have specialized to the case when $F_{02} = -E^\parallel$ is an electric field on the boundary. This arises because of the *consistent* gauge anomaly under gauge transformations

$$\delta_\lambda A_\mu = \partial_\mu \lambda:$$

$$\begin{aligned} \delta_\lambda S_{\text{CS}}[A] &= -\frac{1}{4\pi} \int d^3x \lambda \epsilon^{\mu\nu\lambda} \partial_\mu \sigma \partial_\nu A_\lambda \\ &= \int d^3x \lambda \partial_\mu j_{\text{H}}^\mu = -\delta_\lambda S_{\text{bdry}}[A]. \end{aligned} \quad (7)$$

The anomalous divergence is compensated by protected edge modes on the boundary described by S_{bdry} .

III. ELASTICITY TETRAD FIELDS

The pure (3+1)-d CS term cannot be defined in even space-time dimensions, therefore, the QH response requires additional fields and constitutes a mixed response which is only weakly topologically protected. These fields are due to the (weak) crystalline symmetries of the system.

Specifically, these are the elastic deformations of the crystal lattice described in terms of elasticity tetrads $E_\mu^a(x)$, which represent the hydrodynamic variables of elasticity theory [14,15]. In the absence of dislocations, the tetrads $E^a = E_\mu^a dx^\mu$ are exact differentials. They can be expressed in general form in terms of a system of three deformed crystallographic coordinate planes, surfaces of constant phase $X^a(x) = 2\pi n^a$, $n^a \in \mathbb{Z}$, with $a = 1, 2, 3$, in three dimensions. The intersections of the three constant surfaces

$$X^1(\mathbf{r}, t) = 2\pi n^1, \quad X^2(\mathbf{r}, t) = 2\pi n^2, \quad X^3(\mathbf{r}, t) = 2\pi n^3 \quad (8)$$

are points of the (possibly deformed) crystal lattice

$$L = \{\mathbf{r} = \mathbf{R}(n_1, n_2, n_3) | \mathbf{r} \in \mathbb{R}^3, n^a \in \mathbb{Z}^3\}. \quad (9)$$

The elasticity tetrads are gradients of the three U(1) phase fields X^a , $a = 1, 2, 3$,

$$E_\mu^a(x) = \partial_\mu X^a(x) \quad (10)$$

and have units of crystal momentum. By the inverse function theorem, we can define the inverse vectors

$$E_\mu^a(x) E_a^\nu(x) = \delta_\mu^\nu. \quad (11)$$

In the simplest undeformed case, $X^a(\mathbf{r}, t) = \mathbf{K}^a \cdot \mathbf{r}$, where $E_i^{(0)a} \equiv \mathbf{K}^a$ are the (primitive) reciprocal lattice vectors \mathbf{K}^a . In the general case, they depend on space and time and are quantized in terms of the lattice L in Eq. (9).

In the absence of dislocations, when $X^a(x)$ are globally well defined, the tetrads $E_\mu^a(x)$ are pure gauge and satisfy the integrability condition (in differential form notation)

$$T^a = dE^a = \frac{1}{2} [\partial_\mu E_\nu^a(x) - \partial_\nu E_\mu^a(x)] dx^\mu \wedge dx^\nu = 0. \quad (12)$$

In the presence of dislocations, $T^a \neq 0$, and $X^a(x)$ are multi-valued.

We can also consider a metric associated with these tetrads

$$g_{\mu\nu} = E_\mu^a E_\nu^b \eta_{ab}, \quad (13)$$

where η is the metric associated to the background lattice, say the spatial Euclidean or Minkowski metric. The important difference is that $dn^2 = g_{\mu\nu} dx^\mu dx^\nu$ in terms of the elasticity tetrads is dimensionless and therefore counts the space-time distances in terms of the lattice points of L [16]. The tetrad fields E_μ^a , but not the metric, enter the CS action for a (3+1)-d quantum Hall effect of the lattice QH systems and topological insulators as we now discuss.

IV. (3+1)-d TOPOLOGICAL ACTION FOR QHE

Using the elasticity tetrads, the generalization of the (2+1)-d QH response to (3+1)-d is in principle straightforward. This extends the electromagnetic (2+1)-d CS response to (3+1)-d with the following topological terms, featuring U(1) fields in combination with the elasticity tetrads $E_\mu^a(x)$:

$$S_{(3+1)\text{-d}}[A_\mu] = \frac{1}{8\pi^2} \sum_{a=1}^3 N_a \int d^4x E_\mu^a \epsilon^{\mu\nu\alpha\beta} A_\nu \partial_\alpha A_\beta, \quad (14)$$

where $a = 1, 2, 3$ labels the spatial lattice directions in three space dimensions. The derivation of this formula using semiclassical expansion is in the next section. The tetrads $E_\mu^a(x)$ in Eq. (10) have the dimension of the momentum, and thus the integrals in Eq. (14) are dimensionless ($\hbar = 1$). It follows, as in the (2+1)-d case, that the prefactors are dimensionless and are also expressed in terms of integer topological momentum-space invariants. The integer coefficients N_a are antisymmetric integrals of the Green's functions:

$$N_a = \frac{1}{8\pi^2} \epsilon_{ijk} \int_{-\infty}^{\infty} d\omega \int_{\text{BZ}} dS_a^i \times \text{Tr}[(G\partial_\omega G^{-1})(G\partial_{k_i} G^{-1})(G\partial_{k_j} G^{-1})], \quad (15)$$

where the momentum integral is now over the restricted 2-d BZ torus determined by the area dS_a^i normal to E_i^a .

A similar expression for the (3+1)-d QH was proposed in Ref. [31] without the elasticity tetrads and space-time dependence. In the deformed crystalline systems in (3+1)-d, the tetrads $E_\mu^a(x)$ entering the (3+1)-d Chern-Simons action (14) depend slowly on space and time. For arbitrary background fields, the space-time dependence violates the gauge invariance of the action. However, in the absence of dislocations the latter does not happen for the elasticity tetrads: under deformations Eq. (14) remains gauge invariant due to the condition $dE^a = 0$ in Eq. (12). The variation $\delta_\phi S$ under $\delta A_\mu = \partial_\mu \phi$ is identically zero modulo the bulk/boundary QH currents of the sample. The anomaly cancellation in the presence of dislocations is discussed in detail below.

Here is the main difference between the topological insulator and a gapless system in (3+1)-d, e.g., a Weyl semimetal. Integrating out the fermions leads to the chiral anomaly in the bulk and the ensuing gauge anomaly has to be consistently canceled. In that case, instead of the elasticity tetrads the four-momentum separation of the Weyl points appears in the counterterm [21,32–37]. In the deformed system, this parameter depends on the coordinates, but the integrability condition is now absent, and the term cannot arise without the bulk chiral gauge anomaly.

A. Semiclassical expansion

The response in Eq. (8) can be obtained via the semiclassical Green function formalism [6,8,38]. The assumptions are gauge invariance and semiclassical expansion to lowest order. The effective action for a slowly varying background field $A_\mu(x, t)$ is

$$\log \frac{Z[A]}{Z_0} = iS_{\text{eff}}[A] = i \text{tr} \ln \frac{G[A]}{G_0}. \quad (16)$$

Now, we want to obtain the effective action to first order in gradients of $A_\mu(x)$. We use the fact that the coordinate dependence of $A_\mu(x)$ is semiclassical, i.e., the field varies slowly on the scale of the lattice momenta. In the semiclassical phase-space analysis [6,8,11,38], with $G \equiv G[A] \equiv G(\mathbf{p}, \omega; A_\mu(\mathbf{x}, t))$, we obtain to the lowest order in the gradients of $A_\mu(x)$

$$S_{\text{eff}}[A] = -\frac{i}{4} \int \frac{d^3\mathbf{p} d\omega}{(2\pi)^4} \int d^3\mathbf{x} dt \text{tr} [G\partial_{x^\mu} G^{-1} G\partial_{k_\mu} G^{-1} - G\partial_{k_\mu} G^{-1} G\partial_{x^\mu} G^{-1} (G\partial_{k_\nu} G^{-1})|_{A=0}] A_\nu, \quad (17)$$

where $x^\mu = (t, \mathbf{x})$ and $k_\mu = (\omega, -\mathbf{p})$ and

$$G\partial_{x^\mu} G^{-1} = G\partial_{k_\nu} G_{|A=0}^{-1} \partial_{x^\mu} A_\nu. \quad (18)$$

The trace in the matrix product of Green functions is antisymmetric in the indices of k_μ derivatives. With the convention $\epsilon^{txyz} = -1$, we arrive to

$$S_{\text{eff}}[A] = \frac{i}{12} \int d^3x dt \epsilon^{\alpha\beta\gamma\rho} A_\alpha \partial_\beta A_\gamma \int \frac{d^3\mathbf{p} d\omega}{(2\pi)^4} \epsilon_{\mu\nu\lambda\rho} \times \text{tr}[(G\partial_{k_\mu} G^{-1})(G\partial_{k_\nu} G^{-1})(G\partial_{k_\lambda} G^{-1})]_{A=0}. \quad (19)$$

The momentum-space prefactor can be separated to be of the form

$$i \int d^3p \delta_a^\rho \epsilon_{\mu\nu\lambda\rho} \int \frac{d^2\mathbf{p} d\omega}{24\pi^2} \times \text{tr}[(G\partial_{k_\mu} G^{-1})(G\partial_{k_\nu} G^{-1})(G\partial_{k_\lambda} G^{-1})]_{A=0} = \int d^3p N_a(p^a) = K^a N_a, \quad (20)$$

where $N_a = N_a(p_a)$ is the (2+1)-d integer-valued invariant,

$$N_a(p^a) = \epsilon_{\mu\nu\lambda a} \int \frac{d^2\mathbf{p} d\omega}{24\pi^2} \times \text{tr}[(G\partial_{k_\mu} G^{-1})(G\partial_{k_\nu} G^{-1})(G\partial_{k_\lambda} G^{-1})]_{A=0} \quad (21)$$

evaluated in imaginary time over the cross-sectional reciprocal space, perpendicular to the reciprocal lattice direction K^a . The elementary cell can be taken to be triclinic. Topologically, the integration is over a (pinched) 3-torus and $N_a(p^a)$ is an element of $\pi_3(GL(n, \mathbb{C})) = \mathbb{Z}$ [6].

We conclude that

$$S_{\text{eff}}[A] = \frac{1}{8\pi^2} N_a \int d^3\mathbf{x} dt \epsilon^{\mu\nu\lambda\rho} E_\mu^{(0)a} A_\nu \partial_\lambda A_\rho, \quad (22)$$

where $E_\mu^{(0)a} = (K^a)_i \delta_\mu^i = \int d^3p^a$, $i = x, y, z$, are the reciprocal lattice vectors normal to the different lattice planes with invariants N_a . This is the statement that the (3+1)-d QHE is described by the weak vector invariant $N_a E_\mu^{(0)a}$ protected by the crystalline symmetry.

It is well known that the topological winding number N_a is stable against small variations $\delta G(k_x, k_y, k_z)$ of the Green's function that do not close the gap [6,8,11]. We can consider

small lattice deformations $x^\mu \rightarrow x^\mu + \xi^\mu(x)$, $\xi \ll 1$. Under these, the reciprocal vectors change as

$$E_\mu^{(0)a} = \int dp^a \rightarrow E_\mu^a(x) \equiv \int dp^a(x) \approx \frac{2\pi}{d} (\delta_\mu^a - \partial_\mu \xi^a), \quad (23)$$

where $E_\mu^a(x)$ is a semiclassical, slowly varying field at the lattice scale d in reciprocal space which defines the local normal direction of a set of lattice planes. Note that the topological invariants N_a are assumed to be constant and independent of deformations throughout. The final result is Eq. (8).

B. Hall current in terms of elasticity tetrads

The elasticity tetrads, i.e., elementary deformed reciprocal and direct lattice vectors $E_\mu^a(x) = \partial_\mu X^a(x)$ and the inverse $E_a^\mu(x)$ appear in the EM response, descending from the lattice field phase fields $N_a X^a(x)$ in the presence of deformations. As discussed, the phase fields satisfy the quantization condition $X^a(x) = 2\pi n^a$, $n^a \in \mathbb{Z}^3$, on the lattice points $x \in L$. The Hall conductance is [39]

$$\sigma_{ij} = \epsilon_{ijk} \frac{\sum_a N_a E_k^a(x)}{4\pi^2}, \quad (24)$$

i.e., the conductance is quantized in planes perpendicular to the layer normal $G_i = \sum_a N_a E_i^a(x)$. The reciprocal vector $\mathbf{G} = G_i$ is the weak Hall index of a weak (3+1)-d Chern insulator.

In the nondeformed crystal, where $E_k^{(0)a}$ are primitive reciprocal lattice vectors, this equation transforms to the well-known equation with (3+1)-d quantized Hall conductivity [13,32,36,39]:

$$\sigma_{ij} = \frac{e^2}{2\pi h} \epsilon_{ijk} G_k, \quad (25)$$

where G_k is a reciprocal lattice vector, which is expressed in terms of the topological invariants N_a and the primitive reciprocal lattice vectors $E^{(0)a} = \mathbf{K}^a$:

$$G_k = \sum_{a=1}^3 N_a E_k^{(0)a}. \quad (26)$$

In the deformed crystal, the conductivity tensor is space-time dependent, and thus is not universally quantized. However, the response of the conductivity to deformation is quantized:

$$\frac{d\sigma_{ij}}{dE_k^a} = \frac{e^2}{2\pi h} \epsilon_{ijk} N_a. \quad (27)$$

The Hall current is

$$J^\mu = \frac{-1}{4\pi^2} \sum_{a=1}^3 N_a \epsilon^{\mu\nu\alpha\beta} E_\nu^a \partial_\alpha A_\beta + \frac{1}{8\pi^2} \sum_{a=1}^3 N_a \epsilon^{\mu\nu\alpha\beta} (\partial_\alpha E_\beta^a) A_\nu \quad (28)$$

and has a bulk and a topological defect/boundary component [23,40]. In the absence of dislocations, $dE^a = 0$ in

Eq. (12), and the current represents a dissipationless, fully reversible current, which is conserved due to U(1) gauge invariance

$$\partial_\mu J^\mu = \partial_\mu J_{\text{bulk}}^\mu = 0. \quad (29)$$

C. Chiral magnetic effect

The action (14) and current (28) also describe the chiral magnetic effect (CME) [41,42]. In the presence of periodic directions varying in time, time dependence $X^a(\mathbf{r}, t)$ appears. In the CME an electric current along an applied magnetic field is induced:

$$\mathbf{J} = \frac{1}{4\pi^2} \sum_{a=1}^3 N_a E_t^a \mathbf{B}. \quad (30)$$

This current contains $E_t^a = \partial X^a / \partial t$ and thus it vanishes in equilibrium in agreement with Bloch theorem, according to which the total current is absent in the ground, or in general any equilibrium, state of the system (see, e.g., Ref. [43]). Here, we restrict to spatial lattices under deformations. The CME for a time-periodic insulator with timelike $E_\mu^0 = \omega_F \delta_\mu^0$ (with Floquet drive ω_F) and the temporal invariant $N_0 \neq 0$ was pointed out in Ref. [17]. This can be extended to spatial deformations as well.

V. ANOMALY CANCELLATION AND DISLOCATION ZERO MODES

A. Callan-Harvey effect on dislocations and mixed anomaly

Now, we describe the anomaly inflow. The constraint (12) is violated in the presence of topological defects: dislocations. The density of dislocations equals the torsion for the elasticity tetrads (with vanishing spin connection):

$$T^a = dE^a, \quad T_{kl}^a = (\partial_k E_l^a - \partial_l E_k^a), \quad k, l = x, y, z \quad (31)$$

similar to the role of space-time torsion in gravitational theories [14,18]. The nonzero dislocation density or torsion violates the conservation of the Hall current:

$$\partial_\mu J^\mu = \frac{-1}{8\pi^2} \frac{1}{4} \epsilon^{\mu\nu\alpha\beta} F_{\alpha\beta} \sum_{a=1}^3 N_a T_{\mu\nu}^a. \quad (32)$$

This mixed anomaly represents the Callan-Harvey mechanism of anomaly cancellation [23], which is provided here by the fermion zero modes living on dislocations [23–26,40]. The action remains well defined and gauge invariant in the presence of dislocations, i.e., 2π ambiguities in the phase field X^a , due to zero modes with (1+1)-d covariant anomaly along the dislocation string,

$$\partial_\mu J^\mu = \partial_\mu J_{\text{bulk}}^\mu + \partial_\mu J_{\text{dislocation}}^\mu = 0, \quad (33)$$

as shown in Refs. [23,40,44] for Dirac fermions in the presence of complex vortexlike axionic mass. The Dirac model can be taken as a topological model for the (3+1)-d QH system [11], which we discuss next.

B. Dirac fermion model

To verify the existence of dislocation zero modes on dislocations and overall gauge invariance, let us consider a simplified model with the same symmetries as the




FIG. 1. The four-dimensional gauge field vacuum polarization $\Pi^{\mu\nu}(q, k)$ in Eq. (38) with the axionic vertex $\partial_\lambda X$.

time-reversal and parity-breaking topological insulator with crystalline order, namely, a gapped and time-reversal and parity-breaking Dirac model with a complex mass induced by a scalar field X ,

$$\begin{aligned} S[\psi, \bar{\psi}, A_\mu, X] &= \int d^3\mathbf{x} dt \bar{\psi}(\gamma^\mu i\partial_\mu)\psi - m(x)\bar{\psi}e^{i\gamma^5 X}\psi - \bar{\psi}\gamma^\mu\psi A_\mu \\ &= \int d^3\mathbf{x} dt \bar{\psi}(\gamma^\mu i\partial_\mu - m)\psi \\ &\quad - m\bar{\psi}[e^{i\gamma^5 X} - 1]\psi - \bar{\psi}\gamma^\mu\psi A_\mu + \dots \end{aligned} \quad (34)$$

$$\begin{aligned} \Pi^{\mu\nu}(x, y) &= m\partial_\lambda X \int \frac{d^4q d^4p}{(2\pi)^8} d^4k \delta^{(4)}(k) e^{iq(y-x)} \frac{\partial}{\partial k_\lambda} \text{tr}[i\gamma^\mu G(p) i\gamma^\nu G(p+q) \gamma^5 G(p+q+k) + (\mu \leftrightarrow \nu, k \leftrightarrow q)] \\ &= +4i\epsilon^{\mu\nu\lambda\rho} \partial_\lambda X \int \frac{d^4q d^4p}{(2\pi)^8} e^{iq(y-x)} \frac{m^2 i q_\rho}{(p^2 - m^2)[(p+q)^2 - m^2]} \left[\frac{1}{p^2 - m^2} + \frac{1}{(p+q)^2 - m^2} \right]. \end{aligned} \quad (38)$$

Taking the limit $q \rightarrow 0$, we arrive to the *bulk* effective action [23,40]

$$S_{\text{eff}}[A, X] = \sum_a N_a \frac{1}{8\pi^2} \int d^4x \epsilon^{\mu\nu\lambda\rho} \partial_\mu X^a A_\nu \partial_\lambda A_\rho + \dots \quad (39)$$

after integrating out the massive fermions *far* from the string. The response follows from Eq. (38) by the replacement $X \rightarrow \sum_a N_a X^a$, i.e., instead of a single U(1) axion field, we have a flavored U(1)³ lattice phase field X^a with topological charges N_a which are *not* protected [38] without the existence of the slowly varying lattice phase field X^a .

Let us now extend the above effective action (39) to be valid *everywhere*, essentially by considering a vortex string with a delta-function core.

C. Induced anomaly current

The bulk effective action (39) was derived far from the string defect. Let us now check the induced current due to a topological configuration in $X(x) = N_a X^a(x)$. For simplicity, we will consider singularities only in the slowly varying continuum lattice field $X^a(x)$ and treat the topological numbers N_a as constants. A dislocation in the topological response is a $2\pi n^a N_a$ ambiguity in the field $X = N_a X^a(x)$, and n^a is the Burgers vector of the dislocation and $n = n^a N_a$.

The complex scalar $m(x)e^{i\gamma^5 X(x)}$ is equivalent to a slowly varying mass profile. A vortex line singularity (“axionic string” [23,40]) is the profile where $m(x) \rightarrow m$ far from the singularity and $m(x_c) = 0$, where x_c is the vortex location. The phase field $X(x)$ has a vortex singularity

$$\int_C dX = 2\pi n \quad (35)$$

around all contours around x_c and is such that $X = \text{const} + n\phi$ far from the line defect. In the present case, $X = N_a X^a$ and $n = n^a N_a$. The expansion in Eq. (34) applies far from the vortex. Let us compute the induced effective action *far* from the vortex string by starting from the expansion

$$m\bar{\psi}[e^{i\gamma^5 X} - 1]\psi(y) = im\partial_\mu X(y-x)^\mu \bar{\psi}\gamma^5\psi(x) + \dots \quad (36)$$

The response is found from the axial bosonic polarization vertex $\Pi^{\mu\nu}(x, y)$ [40] (or the equivalent Goldstone-Wilczek current [44]) of the effective action

$$S_{\text{eff}}[A] = \int d^4x d^4y A_\mu(x) \frac{1}{2} \Pi^{\mu\nu}(x, y) A_\nu(y) \quad (37)$$

depicted in Fig. 1 becomes, using $\text{tr}[\gamma^5 \gamma^\mu \gamma^\nu \gamma^\lambda \gamma^\rho] = +4i\epsilon^{\mu\nu\lambda\rho}$ with $\epsilon^{txyz} = -1$,

The induced current separates to bulk and boundary contributions

$$\begin{aligned} \langle J^\mu \rangle &= \delta S_{\text{eff}} / \delta A_\mu \\ &= \frac{-1}{4\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\nu X \partial_\lambda A_\rho + \frac{1}{8\pi^2} \epsilon^{\mu\nu\lambda\rho} (\partial_\lambda \partial_\nu X) A_\rho \end{aligned} \quad (40)$$

due to an electric field $F_{0z} = -E^z$ parallel along the string. This has contributions from the bulk, where $X(r, \phi, z) = n\phi$, and from the string where, from Eq. (35), we get

$$-\epsilon^{ijz} \partial_i \partial_j X = 2\pi n \delta^{(2)}(x_c), \quad i, j = x, y. \quad (41)$$

Therefore,

$$\begin{aligned} \langle J_{\text{bulk}}^\mu \rangle &= \frac{-1}{4\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\nu X \partial_\lambda A_\rho \\ &= \frac{1}{4\pi^2} \frac{n}{r} F_{0z} = -\frac{1}{4\pi^2} \frac{n}{r} E^z = \langle J_{\text{bulk}}^r \rangle, \end{aligned} \quad (42)$$

whence the current per unit time and length in the radial direction outward from the string is $-\frac{n}{2\pi} E^z$. In addition, there is the current localized on the string

$$\begin{aligned} \langle J_{\text{string}}^i \rangle &= \frac{1}{8\pi^2} \epsilon^{\mu\nu\lambda\rho} (\partial_\lambda \partial_\nu X) A_\rho \\ &= \frac{1}{8\pi^2} (2\pi n \delta^{(2)}(x_c)) \epsilon^{ij} A_j = \frac{n}{4\pi} \epsilon^{ij} A_j \delta^{(2)}(x_c), \end{aligned} \quad (43)$$

$i, j = t, z$, with the divergence $\partial_i \langle J_{\text{string}}^i \rangle = \frac{n}{8\pi} \epsilon^{ij} F_{ij} \delta^{(2)}(x_c) = \frac{n}{4\pi} E^z \delta^{(2)}(x_c)$. The divergence of the total current (40) on the string is anomalous

$$\begin{aligned} \partial_\mu \langle J^\mu \rangle &= -\frac{1}{8\pi^2} \epsilon^{\mu\nu\lambda\rho} (\partial_\mu \partial_\nu X) F_{\lambda\rho} \\ &= \frac{1}{4\pi^2} (2\pi n \delta^{(2)}(x_c)) F_{0z} = -\frac{n}{2\pi} E^z \delta^{(2)}(x_c). \end{aligned} \quad (44)$$

On the other hand, the chiral fermions on the string, where they are effectively massless, result in n chiral zero modes with chiral anomaly

$$\begin{aligned} \partial_i \langle J_{(1+1)\text{-d}}^i \rangle &= \partial_i \langle J_{\text{string}}^i \rangle + \partial_i \langle J_{(1+1)\text{-d}}^{i,\text{cons.}} \rangle \\ &= \pm \frac{1}{4\pi} \epsilon^{ij} F_{ij} = \mp \frac{F_{0z}}{2\pi} = \pm \frac{E^z}{2\pi}. \end{aligned} \quad (45)$$

This is the (1+1)-d *covariant* anomaly and is composed of the boundary consistent anomaly plus the contribution localized on the string from the bulk [21,40]. The anomaly due to the bulk-induced current and zero modes cancels.

In contrast, from the previous arguments in Sec. II A we see that the (2+1)-d QH anomaly is matched by the *consistent* anomaly of N -boundary (1+1)-d chiral fermions. This seems to coincide with the observation in Ref. [21]: The *covariant* anomaly is a Fermi-surface effect, whereas the *consistent* anomaly arises from the contribution of all the filled levels.

VI. EXTENSION TO EVEN $D = 2 + 2n$ SPACE-TIME DIMENSIONS

The anomaly equation (32) can be straightforwardly extended to arbitrary even $D = 2 + 2n$ space-time dimensions:

$$\partial_\mu J^\mu \propto \sum_{a=1}^{2n+1} N_a T^a \wedge \underbrace{F \wedge \dots \wedge F}_{n \text{ times}}. \quad (46)$$

In addition to the torsional field strength $T^a = dE^a$, it contains the n -fold antisymmetric tensor product of U(1) gauge field strengths $F = dA$, while the integer-valued topological invariants N_a are expressed in terms of $(2n + 1)$ -dimensional integrals in frequency-momentum space [7].

Equation (46) is valid even in case of $n = 0$, i.e., in one spatial dimension, producing the expected results. Consider a gapped one-dimensional chain of electrons with the action (see also [37])

$$S_{(1+1)\text{-d}}[A] = \frac{N_1}{2\pi} \int dx dt \epsilon^{\mu\nu} E_\mu^1 A_\nu, \quad (47)$$

where the index N_1 is defined via the Green's function $G(\omega, k_x)$,

$$N_1 = \frac{1}{2\pi i} \int d\omega \text{Tr} G(k_x, \omega) \partial_\omega G^{-1}(k_x, \omega). \quad (48)$$

The index N_1 is the same for any k_x without gap closings. Note that in (3+1)-d the index N_a in Eq. (15) is the same for any cross section \mathbf{S}_a of the three-dimensional Brillouin zone, while in (1+1)-d the cross section corresponds to one point k_x in the one-dimensional Brillouin zone.

Variation of Eq. (47) gives the electric current

$$J^\mu = \frac{N_1}{2\pi} \epsilon^{\mu\nu} E_\nu^1, \quad (49)$$

the conservation of which

$$\partial_\mu J^\mu = N_a dE^a = N_1 dE^1 = 0 \quad (50)$$

has a simple interpretation. The condition $dE^a = 0$ is equivalent to the conservation of the sites of the one-dimensional lattice, whereas the index N_1 corresponds to the number of the electrons per site, which is integer for band insulators. As a result, the number of the electrons is trivially conserved under adiabatic deformations.

Since any one-dimensional insulator is described by the topological invariant N_1 , which can only change when the gap closes, we may call any (1+1)-d insulator topological, although the topology of the filled states can only be detected by higher invariants: the Chern numbers. In fact, this is very similar to ordinary metals with Fermi surfaces [11]: the gapless Fermi surface represents a topological object in momentum space protected by an invariant similar to N_1 . Topology provides the stability of the Fermi surface with respect to interactions and explains why metals can be described by Landau Fermi-liquid theory. In this sense, metals can be considered as topological materials, making the zeroth-order invariant N_1 one of the most important topological invariants in the hierarchy of the topological invariants for fermionic systems. In particular, it gives rise to the Luttinger theorem: the number of states in the region between two Fermi points in (1+1)-d does not depend on interaction and arises through topology and adiabatic evolution. This can be generalized for any closed Fermi surface or insulator in higher dimensions [11,45,46].

VII. GRAVITATIONAL QH RESPONSE AND EMERGENT GRAVITY

Now, we briefly describe the anomalous coupling of the elasticity tetrads to the effective space-time metric in the quantum Hall system and speculate on the possibility of relating the elastic metric with the gravitational space-time metric in an effective low-energy effective model of quantum gravity.

The dimensional elasticity tetrads in the (3+1)-d QHE allow us to write the dimensional extension of the (2+1)-d gravitational framing anomaly as a mixed elastic-gravitational anomaly in (3+1)-d [23,47–50]:

$$\begin{aligned} S_{\text{eff,g}} &= \sum_{a=1}^3 \frac{\tilde{N}_a}{192\pi^2} \int d^3x dt E^a \wedge \left(\Gamma_\nu^\mu \wedge d\Gamma_\mu^\nu \right. \\ &\quad \left. + \frac{2}{3} \Gamma_\nu^\mu \wedge \Gamma_\rho^\nu \wedge \Gamma_\mu^\rho \right), \end{aligned} \quad (51)$$

where \tilde{N}_a are effective central charges, $\Gamma_\nu^\mu \equiv \Gamma_{\lambda\nu}^\mu dx^\lambda$ are the Christoffel symbol one-forms of the space-time metric, that arises for example through Luttinger's argument [51]. In particular, the mixed elastic-gravitational Chern-Simons term (51) implies the generalization of the (2+1)-d thermal Hall effect [49,51–54] to (3+1)-d quantum Hall systems and topological insulators with intrinsic Hall effect, although it is third order in derivatives and therefore beyond linear response.

Finally, let us speculate on the hypothetical connection of the elasticity tetrads E_μ^a to real space-time metric and gravity. While the microscopic structure of the relativistic quantum space-time vacuum is not known, phenomenological approaches and effective field theory can be used to describe the effects of the vacuum degrees of freedom. In one of these scenarios, it is assumed that the space-time vacuum has the properties of a (3+1)-d superplastic crystalline medium constructed from E_μ^a [55]. As in the condensed matter elasticity theory, dislocations and disclinations in this space-time crystal describe torsion and curvature of general relativity [14,56–58]. More specifically, in this superplastic model of gravity, the size of the elementary cell in the vacuum space-time crystal is not fixed but in principle can vary arbitrarily with no elementary Planck scale space-time lattice. As a result, the induced action for the gravitational field [55]

$$S[E_\mu^a] = \int_{M^{3,1}} d^4x |E| (KR + \Lambda) \quad (52)$$

contains only dimensionless quantities. Here, the metric $g_{\mu\nu} = \eta_{ab} E_\mu^a E_\nu^b$ is from Eq. (13) and the Ricci curvature scalar $R[E_\mu^a]$ depends on the elasticity tetrads in the standard way, thus making the gravitational constant K (inverse Newton constant) and the cosmological constant Λ dimensionless, $[K] = [R] = [\Lambda] = 1$. The same holds for higher derivative gravity and for the other (nongravitational) physical quantities, such as particle mass $[M] = 1$. Therefore, if gravity is related to fundamental elasticity tetrads, all the measurable physical quantities are dimensionless with respect to the space-time lattice.

VIII. CONCLUSION

In this paper, we have described the mixed topological response of (3+1)-d QH systems using elasticity tetrads. We

have shown how gauge invariance and anomaly inflow arise in the presence of deformations of the weak lattice symmetries protecting the state. The response is a mixed CS response featuring the elasticity tetrad fields of the lattice. This mixed and geometric nature of the (3+1)-d CS response is to be contrasted with the more familiar topological BF theory in four dimensions (see, e.g., Ref. [59]), which represents a mixed response with two one-form gauge fields and their the field strengths appearing in the action.

More generally, the dimensionful elasticity tetrads are the proper hydrodynamic variables related to the weakly protected IQHE/AQHE/CME on general even-dimensional crystalline backgrounds. They describe the topological QH response with elastic and geometric deformations in systems in odd spatial dimensions. In general, Eq. (32) and its extension to higher dimensions, Eq. (46), describe the mixed anomaly in terms of the gauge fields and elastic torsion. These equations do not contain any parameters, except for topological quantum numbers. This is because the elastic tetrads (torsion) have the canonical dimensions of $[I]^{-1}$ (respectively $[I]^{-2}$) instead of the conventional l^0 (respectively $[I]^{-1}$) for the gravitational space-time tetrads (torsion). This allows one to write many mixed “quasitopological Chern-Simons terms,” analogous to mixed axial-gravitational/elastic anomalies, in (3+1)-d quantum Hall systems with weak crystalline symmetries. In principle, we can envisage similar geometric extension of BF theory to (4 + 1)-d using elasticity tetrads.

ACKNOWLEDGMENT

This work has been supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (Grant Agreement No. 694248).

-
- [1] S.-C. Zhang, T. H. Hansson, and S. Kivelson, Effective-Field-Theory Model for the Fractional Quantum Hall Effect, *Phys. Rev. Lett.* **62**, 82 (1989).
 - [2] N. Read, Order Parameter and Ginzburg-Landau Theory for the Fractional Quantum Hall Effect, *Phys. Rev. Lett.* **62**, 86 (1989).
 - [3] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Quantized Hall Conductance in a Two-Dimensional Periodic Potential, *Phys. Rev. Lett.* **49**, 405 (1982).
 - [4] F. D. M. Haldane, Model for a Quantum Hall Effect without Landau Levels: Condensed-Matter Realization of the “Parity Anomaly”, *Phys. Rev. Lett.* **61**, 2015 (1988).
 - [5] T. Matsuyama, Quantization of conductivity induced by topological structure of energy-momentum space in generalized QED₃, *Prog. Theor. Phys.* **77**, 711 (1987).
 - [6] K. Ishikawa and T. Matsuyama, Magnetic field induced multi component QED in three-dimensions and quantum Hall effect, *Z. Phys. C* **33**, 41 (1986); A microscopic theory of the quantum Hall effect, *Nucl. Phys. B* **280**, 523 (1987).
 - [7] M. F. L. Golterman, K. Jansen, and D. B. Kaplan, Chern-Simons currents and chiral fermions on the lattice, *Phys. Lett. B* **301**, 219 (1993).
 - [8] G. E. Volovik, An analog of the quantum Hall effect in a superfluid ³He film, *Zh. Eksp. Teor. Fiz.* **94**, 123 (1988) [*JETP Lett.* **67**, 1804 (1988)].
 - [9] G. E. Volovik, Quantum Hall state and chiral edge state in thin ³He-A film, *Pis'ma Zh. Eksp. Teor. Fiz.* **55**, 363 (1992) [*JETP Lett.* **55**, 368 (1992)].
 - [10] S. C. Zhang and J. Hu, A Four-dimensional generalization of the quantum Hall effect, *Science* **294**, 823 (2001).
 - [11] G. E. Volovik, *The Universe in a Helium Droplet* (Clarendon, Oxford, 2003).
 - [12] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, Topological field theory of time-reversal invariant insulators, *Phys. Rev. B* **78**, 195424 (2008).
 - [13] B. I. Halperin, Possible states for a three-dimensional electron gas in a strong magnetic field, *Jpn. J. Appl. Phys. Suppl.* **26**, 1913 (1987).
 - [14] I. E. Dzyaloshinskii and G. E. Volovick, Poisson brackets in condensed matter, *Ann. Phys.* **125**, 67 (1980).
 - [15] G. E. Volovik and V. S. Dotsenko Jr., Poisson brackets and continual dynamics of the vortex lattice in rotating HeII, *Pisma Zh. Eksp. Teor. Fiz.* **29**, 630 (1979) [*JETP Lett.* **29**, 576 (1979)].

- [16] A. F. Andreev and M. Yu. Kagan, Hydrodynamics of a rotating superfluid liquid, *Zh. Eksp. Teor. Fiz.* **86**, 546 (1984) [*Sov. Phys. JETP* **59**, 318 (1984)].
- [17] J. Nissinen and G. E. Volovik, Tetrads in solids: from elasticity theory to topological quantum Hall systems and Weyl fermions, *Zh. Eksp. Teor. Fiz.* **154**, 1051 (2018) [*JETP* **127**, 948 (2018)].
- [18] H. Kleinert, *Gauge Fields in Condensed Matter: Vol. 2: Stresses and Defects (Differential Geometry, Crystal Melting)* (World Scientific, Singapore, 1989).
- [19] T. Eguchi and P. Freund, Quantum Gravity and World Topology, *Phys. Rev. Lett.* **37**, 1251 (1976).
- [20] L. Alvarez-Gaumé and E. Witten, Gravitational anomalies, *Nucl. Phys. B* **234**, 269 (1985).
- [21] K. Landsteiner, Notes on anomaly induced transport, *Acta Phys. Pol. B* **47**, 2617 (2016).
- [22] J. Gooth *et al.*, Experimental signatures of the mixed axial-gravitational anomaly in the Weyl semimetal NbP, *Nature (London)* **547**, 324 (2017).
- [23] C. G. Callan and J. A. Harvey, Anomalies and fermion zero modes on strings and domain walls, *Nucl. Phys. B* **250**, 427 (1985).
- [24] Y. Ran, Y. Zhang, and A. Viswanath, One-dimensional topologically protected modes in topological insulators with lattice dislocations, *Nat. Phys.* **5**, 298 (2009).
- [25] J. C. Y. Teo and C. L. Kane, Topological defects and gapless modes in insulators and superconductors, *Phys. Rev. B* **82**, 115120 (2010).
- [26] R.-J. Slager, A. Mesaros, V. Juricic, and J. Zaanen, Interplay between electronic topology and crystal symmetry: Dislocation-line modes in topological band insulators, *Phys. Rev. B* **90**, 241403(R) (2014).
- [27] G. E. Volovik, Topology of the $^3\text{He-A}$ film on corrugated graphene substrate, *JETP Lett.* **107**, 115 (2018).
- [28] X. G. Wen, Electrodynamical Properties of Gapless Edge Excitations in the Fractional Quantum Hall States, *Phys. Rev. Lett.* **64**, 2206 (1990).
- [29] R. Jackiw, Fractional charge and zero modes for planar systems in a magnetic field, *Phys. Rev. D* **29**, 2375 (1984).
- [30] T. L. Hughes, R. G. Leigh, and O. Parrikar, Torsional anomalies, Hall viscosity, and bulk-boundary correspondence in topological states, *Phys. Rev. D* **88**, 025040 (2013).
- [31] M. A. Zubkov, Wigner transformation, momentum space topology, and anomalous transport, *Ann. Phys.* **373**, 298 (2016); M. A. Zubkov and Xi Wu, Topological invariant in terms of the Green functions for the quantum Hall effect in the presence of varying magnetic field, [arXiv:1901.06661](https://arxiv.org/abs/1901.06661).
- [32] F. D. M. Haldane, Berry Curvature on the Fermi Surface: Anomalous Hall Effect as a Topological Fermi-Liquid Property, *Phys. Rev. Lett.* **93**, 206602 (2004); Attachment of surface “Fermi arcs” to the bulk Fermi surface: “Fermi-level plumbing” in topological metals, [arXiv:1401.0529](https://arxiv.org/abs/1401.0529).
- [33] A. Cortijo, Y. Ferreirós, K. Landsteiner, and M. A. H. Vozmediano, Elastic Gauge Fields in Weyl Semimetals, *Phys. Rev. Lett.* **115**, 177202 (2015).
- [34] A. G. Grushin, J. W. F. Venderbos, A. Vishwanath, and Roni Ilan, Inhomogeneous Weyl and Dirac Semimetals: Transport in Axial Magnetic Fields and Fermi Arc Surface States from Pseudo-Landau Levels, *Phys. Rev. X* **6**, 041046 (2016).
- [35] D. I. Pikulin, A. Chen, and M. Franz, Chiral Anomaly from Strain-Induced Gauge Fields in Dirac and Weyl Semimetals, *Phys. Rev. X* **6**, 041021 (2016).
- [36] F. Klinkhamer and G. Volovik, Emergent CPT violation from the splitting of Fermi points, *Int. J. Mod. Phys. A* **20**, 2795 (2005).
- [37] S. T. Ramamurthy and T. L. Hughes, Patterns of electromagnetic response in topological semimetals, *Phys. Rev. B* **92**, 085105 (2015).
- [38] J. I. Väyrynen and G. E. Volovik, Soft topological objects in topological media, *JETP Lett.* **93**, 344 (2011).
- [39] M. Kohmoto, B. I. Halperin, and Y.-S. Wu, Diophantine equation for the three-dimensional quantum Hall effect, *Phys. Rev. B* **45**, 13488 (1992).
- [40] S. G. Naculich, Axionic strings: Covariant anomalies and bosonization of chiral zero modes, *Nucl. Phys. B* **296**, 83 (1988).
- [41] K. Fukushima, D. E. Kharzeev, and H. J. Warringa, Chiral magnetic effect, *Phys. Rev. D* **78**, 074033 (2008).
- [42] Q. Li, D. E. Kharzeev, C. Chang, Y. Huang, I. Pletikoscic, A. V. Fedorov, R. D. Zhong, J. A. Schneeloch, G. D. Gu, and T. Valla, Chiral magnetic effect in ZrTe_5 , *Nat. Phys.* **12**, 550 (2016).
- [43] N. Yamamoto, Generalized Bloch theorem and chiral transport phenomena, *Phys. Rev. D* **92**, 085011 (2015).
- [44] J. Goldstone and F. Wilczek, Fractional Quantum Numbers on Solitons, *Phys. Rev. Lett.* **47**, 986 (1981).
- [45] M. Oshikawa, Topological Approach to Luttinger’s Theorem and the Fermi Surface of a Kondo Lattice, *Phys. Rev. Lett.* **84**, 3370 (2000).
- [46] J. T. Heath and K. S. Bedell, Necessary and sufficient conditions for the validity of Luttinger’s theorem, [arXiv:1906.00929](https://arxiv.org/abs/1906.00929).
- [47] E. Witten, Quantum field theory and the Jones polynomial, *Commun. Math. Phys.* **121**, 351 (1989).
- [48] G. E. Volovik, The gravitational-topological Chern-Simons term in a film of superfluid $^3\text{He-A}$, *Pis’ma Zh. Eksp. Teor. Fiz.* **51**, 111 (1990) [*JETP Lett.* **51**, 125 (1990)].
- [49] N. Read and D. Green, Paired states of fermions in two dimensions with breaking of parity and time reversal symmetries and the fractional quantum Hall effect, *Phys. Rev. B* **61**, 10267 (2000).
- [50] A. G. Abanov and A. Gromov, Electromagnetic and gravitational responses of two-dimensional noninteracting electrons in a background magnetic field, *Phys. Rev. B* **90**, 014435 (2014).
- [51] J. M. Luttinger, Theory of thermal transport coefficients, *Phys. Rev.* **135**, A1505 (1964).
- [52] M. Stone, Gravitational anomalies and thermal Hall effect in topological insulators, *Phys. Rev. B* **85**, 184503 (2012).
- [53] M. Stone and JiYoung Kim, Mixed anomalies: Chiral vortical effect and the Sommerfeld expansion, *Phys. Rev. D* **98**, 025012 (2018).
- [54] A. Gromov and A. G. Abanov, Thermal Hall Effect and Geometry with Torsion, *Phys. Rev. Lett.* **114**, 016802 (2015).
- [55] F. R. Klinkhamer and G. E. Volovik, Tetrads and q -theory, *Pis’ma Zh. Eksp. Teor. Fiz.* **109**, 369 (2019) [*JETP Lett.* **109**, 362 (2019)].
- [56] B. A. Bilby and E. Smith, Continuous distributions of dislocations: A new application of the methods of

- non-Riemannian geometry, *Proc. R. Soc. Sect. A* **231**, 263 (1955); Continuous distributions of dislocations. III, **236**, 481 (1956).
- [57] E. Kröner, Allgemeine Kontinuumstheorie der Versetzungen und Eigenspannungen, *Arch. Rational Mech. Anal.* **4**, 273 (1959).
- [58] H. Kleinert and J. Zaanen, World nematic crystal model of gravity explaining the absence of torsion, *Phys. Lett. A* **324**, 361 (2004).
- [59] G. Y. Cho and J. E. Moore, Topological BF field theory description of topological insulators, *Ann. Phys. (NY)* **326**, 1515 (2011).