



This is an electronic reprint of the original article. This reprint may differ from the original in pagination and typographic detail.

Niessen, Eva; Bracco, Martina; Mutanen, Tuomas P.; Robertson, Edwin M. An analytical approach to identify indirect multisensory cortical activations elicited by TMS?

Published in: Brain Stimulation

DOI: 10.1016/j.brs.2021.02.003

Published: 01/03/2021

Document Version Publisher's PDF, also known as Version of record

Published under the following license: CC BY-NC-ND

Please cite the original version:

Niessen, E., Bracco, M., Mutanen, T. P., & Robertson, E. M. (2021). An analytical approach to identify indirect multisensory cortical activations elicited by TMS? *Brain Stimulation*, *14*(2), 376-378. https://doi.org/10.1016/j.brs.2021.02.003

This material is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Brain Stimulation 14 (2021) 376-378

Contents lists available at ScienceDirect

Brain Stimulation

journal homepage: http://www.journals.elsevier.com/brain-stimulation

An analytical approach to identify indirect multisensory cortical activations elicited by TMS?



We analysed a pre-existing TMS-EEG dataset (58 participants, 41 females, 23 ± 4 years old (mean \pm std); right-handed (defined by Edinburgh Handedness Inventory); neurologically and psychiatrically normal participants, meeting the safety criteria for the use of TMS). Participants received 126 neuro-navigated biphasic single TMS-pulses to left primary motor cortex (M1), while EEG (a 62-channel system) was concurrently recorded. Stimulation intensity was set at 80% of participants' rMT ([7]; 54 \pm 8% MSO (mean \pm std), range from 31 to 77% MSO). Individually adjusted white noise (the volume was gradually increased until participants could not hear the coil click anymore or their threshold of discomfort was reached) was played during the stimulation via padded earplugs.

In our analysis, we concentrated on the most commonly assessed dimension of TMS-EEG signal (i.e., TEPs [3]; Fig. 1B). Initially, the signal was cleaned from the typical noise and TMSinduced artifacts, via the MATLAB-based toolbox EEGLAB and the TMS-EEG signal analyser plugin (TESA [8]). For evaluating the spatio-temporal effects of absolute stimulation intensity, we computed TEPs (as the average time-course over trials) for each channel and each subject separately (in the average reference). Then, at each resulting time point (331; i.e. 20 - 350 ms after the TMS pulse) and at each channel (62 channels) we used linear regression to test whether the independent variable, subjectspecific absolute intensity, can predict the subject-specific TEP amplitude (FieldTrip [9]). To test for possible significance, we ran independent-samples regression coefficient t-tests (two-tailed, α level p < 0.025). To control the multiple comparison problem (the familywise error rate [10]), clusters were built by assembling neighbouring significant spatio-temporal samples and tested against corresponding permutation statistics (5000 iterations, cluster α level p < 0.025 [10]).

We found that absolute stimulation intensity showed a significant regression against TEP amplitude in three different clusters. All three of the clusters occurred at later latencies. One of these was a negative cluster showing a negative TEP deflection (Fig. 1C, bottom row). The first was a positive cluster visible at 174–241 ms after the TMS pulse and mainly involved central

sound, stronger coil vibration, and stronger stimulation of

https://doi.org/10.1016/j.brs.2021.02.003

1935-861X/© 2021 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).



Dear Editor,



霐

BRAIN



Electroencephalography (EEG) is widely used for detecting

transcranial-magnetic stimulation (TMS) evoked responses in the

human brain. TMS-evoked potentials (TEPs) may contain both

direct activations due to the TMS-induced cortical electric field

and indirect cortical activations due to the subsequent multisen-

sory responses to TMS [1]. Distinguishing between direct and indi-

rect sources of cortical activation has largely been attempted using

careful experimental designs [2-5]. Here, we complement those

advances by providing a novel statistical approach. Our results

converge with the pattern of indirect activation identified by recent

experimental work [3], which provides encouraging support for the

with the indirect response of stimulation. One approach is to sup-

press the sources of multisensory stimulation and consequently minimize indirect cortical activations. This is powerfully illustrated

in a recent study, in which the sound of the coil was very efficiently masked by playing noise-masking via earplugs as well as using ear

defenders, and the vibrations of the coil were attenuated by using a

foam layer beneath the coil [3]. Whether this suppression approach

is completely effective is debated (e.g. [4,6]). An alternative exper-

imental approach is to use sham stimulation to mimic, and conse-

quently isolate multisensory sources of indirect cortical activation

[3–5]. Yet, this requires techniques to accurately reproduce the

multisensory response elicited by TMS [3,4], which is challenging

to achieve. Both of these approaches deal with indirect responses

to stimulation using experimental manipulations within-subjects. We offer a complementary approach using between-subjects anal-

ysis to indentify the indirect cortical activations due to stimulation.

tion. When stimulating the primary motor cortex (M1) this is

called the resting-motor threshold (rMT, [7]; relative intensity).

This corresponds to a percentage of stimulator output (absolute in-

tensity). For example, for one participant 80% of rMT may require

48% of the maximum stimulator output (MSO); while for another

participant 80% of rMT will be achieved with only 36% of MSO

(Fig. 1A). The administered relative intensity (e.g. 80% of rMT) rep-

resents a normalization implying that the magnitude of direct mo-

tor cortical activation should be comparable across subjects. By

contrast, the indirect activation of a TMS pulse may be closely

related to the amount of sensory input, which itself is higher

with higher absolute intensity (e.g. 36% vs 48% of MSO; i.e., louder

Commonly in TMS—EEG studies, participants receive a stimulation intensity tailored to their individual susceptibility to stimula-

Several different experimental strategies have been used to deal

potential utility of this novel approach.



Fig. 1. Experimental design and results (A) Despite receiving the same relative intensity (80% rMT), every participant (P1 vs. P2) received a different, absolute intensity (48% vs. 36%; for example), which determines the magnitude of multisensory stimulation (coil vibration, coil click and excitation of nerves in the skin; see table). **(B)** The grand average TEP (i.e. mean over C1, C3, CP1 and CP3), showing a typical response when targeting M1. **(C)** On the left are shown the TEPs and the different time windows when their amplitude is related significantly with absolute intensity. The three clusters likely represent non-specific multisensory contributions from multiple sources. TEPs are averaged for those channels included within the significant cluster (black dots on scalp distribution). On the right are shown the respective topographical plots for each of the three clusters. Black box: time-window excluded from the analysis due to TMS-artifacts (i.e., 0-20ms). Grey box: significant time window in the cluster-based analysis. TEPs shadowing: standard error of the mean. Black asterisk (*): p < 0.05. Black dots: significant channels in the cluster-based analysis. rMT: resting motor threshold. MSO: maximum stimulator output. TEP: TMS-evoked potential.

electrodes (corrected p = 0.025; Fig. 1C, top row). The second positive cluster occurred later at 264–337 ms and was located above the left sensorimotor cortex (corrected p = 0.034, Fig. 1C, middle row). Finally, the negative cluster was found at 259–303 ms and involved central electrodes as well as electrodes located above left-frontal regions (corrected p = 0.038, Fig. 1C, bottom row). Based on their spatio-temporal patterns, the three clusters may be due to indirect cortical activations elicited by the sensory stimulation caused by TMS (e.g. a combination of auditory and somatosensory processes [3]).

These results show that the amplitude of late (>170 ms) TEPs were related to the absolute stimulation intensity. This may identify indirect cortical activations, and suggests that in a between-subject design absolute intensity is linked to the multisensory stimulation elicited by delivering TMS. Despite the methodological differences, our results converge both temporally and spatially with the pattern of indirect activation identified by recent experimental work [3]. Hence, our analytical method makes the identification of indirect activations possible - unconstrained by experimental design allowing new questions to be posed using innovative designs, which were previously impossible. Yet, further work is needed to fully validate this between-subject analytical approach. Nonetheless, it may prove to be a simple yet effective tool to identify and potentially remove multisensory contributions from TMS-EEG data; without the need to include additional experimental conditions within a study.

Declaration of competing interest

We declare that the research was conducted free from any commercial or financial relationships that could have lead, or be seen to have lead to a conflict of interest.

Acknowledgements

We are grateful to the Air Force Office of Scientific Research (FA9550-16-1-0191; EMR) and the Academy of Finland (Grant No. 321631; TPM) for supporting this work.

References

- Ilmoniemi RJ, Kičić D. Methodology for combined TMS and EEG. Brain Topogr 2009;22:233–48.
- [2] Biabani M, Fornito A, Mutanen TP, Morrow J, Rogasch NC. Characterizing and minimizing the contribution of sensory inputs to TMS-evoked potentials. Brain Stimul 2019;12:1537–52.
- [3] Rocchi L, Di Santo A, Brown K, Ibáñez J, Casula E, Rawji V, Di Lazzaro V, Koch G, Rothwell J. Disentangling EEG responses to TMS due to cortical and peripheral activations. Brain Stimul 2021;14:4–18.
- [4] Conde V, Tomasevic L, Akopian I, Stanek K, Saturnino GB, Thielscher A, Bergmann TO, Siebner HR. The non-transcranial TMS-evoked potential is an

inherent source of ambiguity in TMS-EEG studies. Neuroimage 2019;185: 300-12.

- [5] Gordon PC, Desideri D, Belardinelli P, Zrenner C, Ziemann U. Comparison of cortical EEG responses to realistic sham versus real TMS of human motor cortex. Brain Stimul 2018;11:1322–30.
- [6] Belardinelli P, Biabani M, Blumberger DM, Bortoletto M, Casarotto S, David O, Desideri D, Etkin A, Ferrarelli F, Fitzgerald PB, et al. Reproducibility in TMS-EEG studies: a call for data sharing, standard procedures and effective experimental control. Brain Stimul 2019;12:787–90.
- [7] Rossini PM, Burke D, Chen R, Cohen LG, Daskalakis Z, Di Iorio R, Di Lazzaro V, Ferreri F, Fitzgerald PB, George MS, et al. Non-invasive electrical and magnetic stimulation of the brain, spinal cord, roots and peripheral nerves: basic principles and procedures for routine clinical and research application. An updated report from an I.F.C.N. Committee. Clin Neurophysiol 2015;126: 1071–107.
- [8] Rogasch NC, Sullivan C, Thomson RH, Rose NS, Bailey NW, Fitzgerald PB, Farzan F, Hernandez-Pavon JC. Analysing concurrent transcranial magnetic stimulation and electroencephalographic data: a review and introduction to the open-source TESA software. Neuroimage 2017;147:934–51.
- [9] Oostenveld R, Fries P, Maris E, Schoffelen J-M. FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell Neurosci 2011;2011:156869.
- [10] Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEGdata. J Neurosci Methods 2007;164:177–90.

Eva Niessen¹

Institute of Neuroscience & Psychology, Centre for Cognitive Neuroimaging, University of Glasgow, Glasgow, G12 8QB, UK

Individual Differences and Psychological Assessment, Faculty of Human Sciences, University of Cologne, Cologne, Germany

Martina Bracco^{1,**}

Institute of Neuroscience & Psychology, Centre for Cognitive Neuroimaging, University of Glasgow, Glasgow, G12 8QB, UK

Tuomas P. Mutanen

Institute of Neuroscience & Psychology, Centre for Cognitive Neuroimaging, University of Glasgow, Glasgow, G12 8QB, UK

Department of Neuroscience and Biomedical Engineering, Aalto University, Espoo, Finland

Edwin M. Robertson^{*} Institute of Neuroscience & Psychology, Centre for Cognitive Neuroimaging, University of Glasgow, Glasgow, G12 8QB, UK

** Corresponding author.

* Corresponding author. Institute of Neuroscience and Psychology, 62 Hillhead Street, University of Glasgow, Glasgow, G12 8QB, UK. *E-mail address:* martina.bracco@icm-institute.org (M. Bracco). *E-mail address:* edwin.robertson@glasgow.ac.uk (E.M. Robertson).

> 11 January 2021 Available online 10 February 2021

¹ Both authors contributed equally.

¹ Both authors contributed equally.