
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Wallach, Christoph; Geitner, Felix S.; Karttunen, Antti J.; Fässler, Thomas F.

Boranyl-Functionalized [Ge₉] Clusters : Providing the Idea of Intramolecular Ge/B Frustrated Lewis Pairs

Published in:
Angewandte Chemie - International Edition

DOI:
[10.1002/anie.202012336](https://doi.org/10.1002/anie.202012336)

Published: 01/02/2021

Document Version
Publisher's PDF, also known as Version of record

Published under the following license:
CC BY-NC-ND

Please cite the original version:
Wallach, C., Geitner, F. S., Karttunen, A. J., & Fässler, T. F. (2021). Boranyl-Functionalized [Ge₉] Clusters : Providing the Idea of Intramolecular Ge/B Frustrated Lewis Pairs. *Angewandte Chemie - International Edition*, 60(5), 2648-2653. <https://doi.org/10.1002/anie.202012336>

Frustrated Lewis Pairs

Boranyl-Functionalized [Ge₉] Clusters: Providing the Idea of Intramolecular Ge/B Frustrated Lewis Pairs

Christoph Wallach[†], Felix S. Geitner[†], Antti J. Karttunen, and Thomas F. Fässler*

Dedicated to Professor Herbert W. Roesky on the Occasion of his 85th Birthday

Abstract: The unique three-dimensional structure of spherical, homoatomic nine-atom germanium clusters opens various possibilities for the spatial arrangement of functional groups. Ligands comprising lone pairs have recently been introduced in the cluster sphere, and we now report the addition of a boranyl group to the cluster featuring a Ge–B *exo*-cluster bond. The reaction of the twofold-silylated cluster [Ge₉{Si(TMS)₃}₂]^{2–} (TMS = trimethylsilyl) with 2-chloro-1,3,2-diazaborolidines DAB^R-Cl leads to the first boranyl-functionalized [Ge₉] clusters [Ge₉{Si(TMS)₃}₂DAB^R][–] (R = methyl (**1a**), isopropyl (**2a**), *ortho*-tolyl (**3a**)). The anions **2a** and **3a** were structurally characterized as [NHC^{Dipp}Cu]⁺ complexes (NHC^{Dipp} = 1,3-di(2,6-diisopropylphenyl)imidazolyldine) through single crystal X-ray structure determination. Quantum-chemical calculations manifest the frustrated Lewis pair (FLP) character of the boranyl-functionalized cluster [Ge₉{Si(TMS)₃}₂BCy₂][–] (**4a**).

By fusion of stoichiometric amounts of the elements, the tetrel Zintl phase K₄Ge₉ containing soluble [Ge₉]^{4–} ions is easily accessible. These clusters have shown great potential to react with main group and transition metal compounds.^[1] The silylation of [Ge₉] through a heterogeneous reaction of K₄Ge₉ with chloro-tris(trimethylsilyl)silane in MeCN has proven to be a straightforward way to reduce the fourfold negative cluster charge thereby enhancing the stability and solubility of the clusters. By variation of the stoichiometry of the reaction, either bis-silylated [Ge₉{Si(TMS)₃}₂]^{2–} or tris-silylated [Ge₉{Si(TMS)₃}₃][–] (TMS: trimethylsilyl) species are accessible in high yields.^[2] Furthermore, various tris-silylated clusters have been prepared with silyl groups of different bulkiness.^[3] In

similar reactions the tris-stannylated cluster [Ge₉(SnⁱPr₃)₃][–] can also be obtained.^[4] Recently, also ligands with donor functions have been introduced to the sphere of the [Ge₉] clusters allowing for subsequent reactions with Lewis acids, leading to charge-neutral zwitterionic cluster compounds [Ge₉{Si(TMS)₃}₂(^tBu₂P)]M(NHC^{Dipp}) (M: Cu, Ag, Au).^[5] Up to three ligands with Lewis base groups can be introduced in form of the tris-phosphanyl-functionalized clusters [Ge₉(PRR¹)₃][–] (R: NⁱPr₂; R¹: NⁱPr₂, ^tBu).^[6] These clusters still possess six or seven unsubstituted cluster vertices with low oxidation state,^[7] and are thus prone to undergo subsequent reactions.

Owing to the shielding of the [Ge₉] core, reactions of the tris-silylated cluster are limited to the introduction of less bulky main group element ligands, yielding uncharged cluster compounds with Ge–C, Ge–P and Ge–Sn *exo*-cluster bonds such as [Ge₉{Si(TMS)₃}₃R] [R: Et,^[8] (CO)^tBu,^[9] ^tBu,^[9] {(CH₂)_nCH=CH₂} (n: 1, 3)^[7c], [Ge₉{Si(TMS)₃}₃(PRR¹)] (R = R¹: ⁱPr, Cy,^[5] R: ^tBu, R¹: {(CH₂)₃CH=CH₂}^[10]), and [Ge₉{Si(TMS)₃}₃SnR₃] (R: ^tBu,^[8] Ph^[11]). Except for the Sn-containing species, which (partially) show multicenter Ge–Sn bonds, all main group element fragments are attached to the [Ge₉] entity via 2-center-2-electron (2c-2e) bonds pointing radially away from the center of the cluster. The Lewis basic property of the [Ge₉] core is manifested in various isolated transition metal decorated clusters such as [Ge₉{Si(TMS)₃}₃TiCp₂-(MeCN)], [Cp₂Ti(η¹-Ge₉{Si(TMS)₃}₂)₂]^{3–},^[12] [Ge₉{Si(TMS)₃}₃-Cr(CO)₅][–]^[13] as well as [Zn(η¹-Ge₉{Si(TMS)₃}₂)₄]^{6–}.^[14]

Because of the decreased shielding of the [Ge₉] cluster core, the di-anion [Ge₉{Si(TMS)₃}₂]^{2–} is an attractive candidate for subsequent reactions. Upon addition of further halo-silanes, mixed-substituted [Ge₉] clusters of the general composition [Ge₉{Si(TMS)₃}₂R][–] [R: Si(TMS)₂SiPh₃,^[2a] Si(TMS)₂Si(ⁱPr)₃,^[3b] SiPh₂{(CH₂)_nCH=CH₂}^[16] (n = 1, 3)] are formed. Using a rigid bis-halo-silane, a connection of two clusters to the anion [[Si(SiMe₃)₃]₂Ge₉-SiMe₂-(C₆H₄)-SiMe₂-Ge₉{Si(SiMe₃)₃}₂]^{2–} is achieved.^[17] Comparable to the reactions reported for the tris-silylated species, alkyl-, germanyl- or phosphanyl-decorated anions [Ge₉{Si(TMS)₃}₃Et]^{2–} (one Si(TMS)₃ group is cleaved upon crystallization),^[18] [Ge₉{Si(TMS)₃}₂Ge(TMS)₃][–]^[19] or [Ge₉{Si(TMS)₃}₂(PRR¹)][–] (R, R¹: alkyl, aryl, aminoalkyl)^[5,10,20] can be synthesized, revealing the prosperous chemistry of [Ge₉{Si(TMS)₃}₂]^{2–} with halo-main group element compounds.

We have shown that ligands with free electron lone pairs such as phosphanyl groups (Scheme 1, A) allow for the attachment of Lewis acidic organometallic fragments,^[5,10,20] and now focused on the introduction of electron acceptor


[*] M. Sc. C. Wallach,^[‡] Dr. F. S. Geitner,^[‡] Prof. Dr. T. F. Fässler
Department Chemie, Technische Universität München
Lichtenbergstraße 4, 85747 Garching b. München (Germany)
E-mail: thomas.faessler@lrz.tum.de

Prof. Dr. A. J. Karttunen
Department of Chemistry and Materials Science, Aalto University
00076 Aalto (Finland)

[‡] These authors contributed equally to this work.

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202012336>.

© 2020 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.




Scheme 1. Different Lewis basic sites at $[\text{Ge}_9]$ occurring in reported **A**^[5,10,20] and boranyl-functionalized $[\text{Ge}_9]$ cluster **B** with FLP character according to quantum-chemical calculations. The lone pair of only one Ge atom is shown. **C** shows a potential mixed-functionalized $[\text{Ge}_9]$ core featuring both Lewis acidic and basic moieties.

functions to the sphere of the $[\text{Ge}_9]$ cluster (Scheme 1, **B**). Scheschkewitz and co-workers recently reported on the grafting of silicoindoles of general composition Si_6R_6 with $[\text{BH}_3]^-$ fragments,^[21] as well as p-doping such clusters by a formal exchange of one Si vertex by a boranyl group.^[22] These clusters exhibit unsubstituted vertices and thus are conceptually related to *Zintl* ions.^[21b] For several reasons we particularly suspected halo-boranes, 1,3,2-diazaborolidines (DABs), as promising candidates for this venture. Due to the straightforward preparation of 2-halo-DABs from BX_3 (X : Cl, Br) and diamines, the wingtip substituents can easily be tuned concerning their bulkiness. Furthermore, the electron deficiency of the boron center is compensated by π -donation of the lone pairs of adjacent N atoms into the boron p_z orbital, thereby enhancing the stability of the compounds. These species have been established as ligands in complex chemistry reported for the shell-structured ion $[\text{In}_{68}[\text{DAB}^{\text{Dipp}}]_{12}]^-$ ^[23] or in lanthanide boryl compounds.^[24] Furthermore, diazaboroles act as σ -donor ligands and have recently been used to prepare new types of low-valent group 14 element (Si ,^[25] Ge ,^[25,26] Sn ,^[25,26c,27] Pb ^[28]) compounds. Such species are prone to activate small molecules via oxidative addition (e.g. H_2 or further nonpolar $E\text{--H}$ bonds; E : B, Si, N, O), and also insertion reactions into the Ge--B bond (e.g. phenylacetylene and diphenylacetylene) have been reported.^[26c,e,27a] Within this work we investigated reactions of the negatively charged germanide cluster $[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2]^{2-}$ towards DABs, focusing on the generation of a hitherto

unreported boranyl-functionalized cluster species revealing unprotected Ge atoms in negative oxidation states.

Treatment of solid $\text{K}_2[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2]$ with THF or 1,4-dioxane solutions of $\text{DAB}^{\text{R}}\text{--Cl}$ (R : Me, ^iPr , *o*-tol, Mes = 2,4,6-trimethylbenzene) yielded deep red reaction mixtures, which were stirred for 3–5 h at room temperature. Subsequently, the mixtures were filtered, and the solvent was removed in vacuo. The remaining oily solids were treated with hexane, which was then removed in vacuo to eliminate traces of remaining solvents (this procedure was repeated twice). The brown (R : Me, Mes) or ochre (R : ^iPr , *o*-tol) residues were investigated by NMR spectroscopy.

^1H NMR examinations indicated the formation of $[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{DAB}^{\text{R}}]^-$ [R : Me (**1a**), ^iPr (**2a**) and *o*-tol (**3a**)] by the downfield shift of the cluster's silyl group protons and the 1:2 ratio of signals assigned to the boranyl moiety and the cluster's silyl protons. The formation of **1a–3a** (Scheme 2a) was confirmed by further NMR experiments (^{11}B , ^{13}C and ^{29}Si). In the ^{11}B NMR spectra a significant downfield shift of the product signal from approx. 27 ppm ($\text{thf-}d_8$) ($\text{DAB}^{\text{R}}\text{--Cl}$; R : Me, ^iPr , *o*-tol) to 43.31 ppm (**1a**), 41.90 ppm (**2a**) and 42.92 ppm (**3a**) was observed. Moreover, anions **1a–3a** were detected in ESI MS experiments at m/z 1246.6 (**1a**), m/z 1302.6 (**2a**) and m/z 1398.7 (**3a**) in the negative-ion mode (Figure 1).



Scheme 2. a) Overview on the reactivity of $\text{K}_2[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2]$ towards $\text{DAB}^{\text{R}}\text{--Cl}$ (R : Me, ^iPr , *o*-tol, Mes); b) formation of **2a-Cu** and **3a-Cu** upon addition of $\text{NHC}^{\text{Dipp}}\text{CuCl}$ to **2a** and **3a**, respectively.




Figure 1. Selected areas of ESI MS spectra of anions **1a–3a**. Spectra were recorded from diluted THF solutions of the samples in the negative-ion mode (3500 V, 300°C). Calculated isotope patterns are represented as black bars. a) molecule peak of **1a** $[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{DAB}^{\text{Me}}]^-$ at m/z 1246.6; b) molecule peak of **2a** $[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{DAB}^{\text{Pr}}]^-$ at m/z 1302.6; c) molecule peak of **3a** $[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{DAB}^{\text{o-tol}}]^-$ at m/z 1398.7. The respective overview spectra can be found in the Supporting Information.

For crystallization, the crude solids **1a-K** to **3a-K** containing the K^+ -salt of **1a** to **3a** were dissolved in toluene and filtered. Unfortunately, the purification of **1a-K** and **2a-K** has not been successful to date, but $[Ge_9\{Si(TMS)_3\}_2DAB^{o-tol}]K$ (**3a-K**) was obtained as an ochre, polycrystalline powder from concentrated toluene solutions. In analogous reactions of $K_2[Ge_9\{Si(TMS)_3\}_2]$ with $DAB^{Mes}-Cl$ no conversion of the reactants was observed (14 d at room temperature and at elevated temperature; Scheme 2a). Since the DAB^{Mes} ligand has a higher steric demand, we assume that there is a steric limitation for the introduction of boranyl moieties at $[Ge_9]$.

In previous studies we found that neutral species with a $[NHC^{Dipp}Cu]^+$ moiety as the fourth substituent crystallized much more easily than the respective anionic species.^[6,20] Since the isolation of single-crystalline material suitable for structure determination was not possible from the isolated solids, we carried out subsequent reactions of **1a-K** to **3a-K** with $NHC^{Dipp}CuCl$ (Scheme 2b). **1a-K** to **3a-K** were dissolved in MeCN and the resulting dark red solutions were treated dropwise with MeCN solutions of the Cu-NHC complex, immediately resulting in the precipitation of brown (**1a-Cu**) or reddish (**2a-Cu**, **3a-Cu**) solids. In order to assure complete conversion, the mixtures were stirred at room temperature for further 30 min, before the supernatant red solutions were filtered, and the residues were washed with MeCN. 1H NMR examinations of the solids indicated the addition of $[NHC^{Dipp}Cu]^+$ to **2a** and **3a** under formation of $NHC^{Dipp}Cu[Ge_9\{Si(TMS)_3\}_2DAB^R]$ [**R**: *Pr* (**2a-Cu**) and *o*-tol (**3a-Cu**)]. By contrast, the product resulting from the reaction of **1a** with $NHC^{Dipp}CuCl$ was identified as the previously published species $(NHC^{Dipp}Cu)_2[Ge_9\{Si(TMS)_3\}_2]$ comprising two $[NHC^{Dipp}Cu]^+$ moieties.^[29] The Ge–B bond cleavage upon the reaction with the Cu-NHC complex hints for an increased lability of DAB moieties with smaller wingtip substituents. Crude materials of **2a-Cu** and **3a-Cu** were dissolved in toluene and filtered to remove remaining solids. The samples were stored in a freezer at $-40^\circ C$ for crystallization yielding red block-shaped single crystals of **2a-Cu** and **3a-Cu** suitable for single crystal X-ray diffraction. The isolated materials were further characterized by elemental analysis and NMR spectroscopy (1H , ^{11}B , ^{13}C and ^{29}Si). Compounds **2a-Cu** and **3a-Cu** crystallize in the monoclinic space groups $P2_1/n$ and $I2/a$, respectively, and show functionalized clusters and $[NHC^{Dipp}Cu]^+$ moieties in a 1:1 ratio. The shapes of the $[Ge_9]$ cluster cores are best described as slightly distorted, tricapped trigonal prisms (their heights are emphasized by fragmented thick lines between the eclipsed triangles Ge1 to Ge3 and Ge7 to Ge9, Figure 2), with one prism height [$d(Ge1-Ge7) = 3.584(1)$ Å (**2a-Cu**) and $d(Ge1-Ge7) = 3.682(1)$ Å (**3a-Cu**)] being significantly longer compared to the others [$3.164(2)$ Å (Ge2–Ge8) and $3.283(1)$ Å (Ge3–Ge9) in **2a-Cu** and $3.147(1)$ Å (Ge3–Ge9) and $3.166(1)$ Å (Ge2–Ge8) in **3a-Cu**], resulting in C_{2v} -symmetry. The three substituents (two silyl groups and the DAB moiety) bind to three capping Ge atoms (Ge4, Ge5 and Ge6; Figure 2). All Ge–Ge and Ge–Si distances are in the typical range of known functionalized $[Ge_9]$ cluster species.^[2,30] The DAB ligands in **2a-Cu** and **3a-Cu** bind to the $[Ge_9]$ cluster through *exo*-cluster single bonds between one cluster vertex atom and the B atom, with Ge–B




Figure 2. Molecular structures of compounds **2a-Cu** (a,b) and **3a-Cu** (c,d). Ellipsoids are shown at a 50% probability level. a) Front view on compound **2a-Cu** with the two silyl groups and the DAB ligand bonding to the capping Ge atoms of the tricapped trigonal prism. b) Side view on **2a-Cu** showing C_{2v} symmetric shape of $[Ge_9]$ core (prism heights are indicated by thick fragmented lines) and coordination of $[NHC^{Dipp}Cu]^+$ moiety via one of the trigonal prism bases. c) and d) analogue pictures for compound **3a-Cu**. In all pictures protons as well as toluene solvent molecules are omitted. Moreover, TMS groups of silyl substituents (a–d) as well as the NHC^{Dipp} (a,c) ligand are omitted and carbon atoms of DAB ligands are shown as black (a,c) or grey (b,d) wire sticks for clarity. For the same reason the Dipp wingtips in b) and d) are indicated as grey spheres.

distances of $2.063(5)$ Å and $2.055(5)$ Å, respectively. The bond lengths are in the range of previously reported Ge–B bonds in germaborates,^[31] germacarboranes^[32] or germynes.^[25,26b,c,e,33] The $[NHC^{Dipp}Cu]^+$ fragment coordinates one of the trigonal bases of the prism in an η^3 -fashion with Ge–Cu distances known for analogous complexes.^[6,20,29,34]

The anions **1a–3a** and compounds **2a-Cu** and **3a-Cu** are the first examples of boranyl-functionalized $[Ge_9]$ clusters. In general, group 13 element decoration of silylated $[Ge_9]$ species is rather rare if compared to the plethora of examples for group 14 (C-, Si-, Ge-, Sn-based) and group 15 (P) element substituted $[Ge_9]$ clusters. $[Ge_9\{Si(TMS)_3\}_3Tl]$ is the only known example of a triel derivative, however, the Tl atom is included as an additional cluster vertex.^[8]

The most interesting aspect about the anions **1a–3a** is the targeted introduction of boranyl groups to the $[Ge_9]$ core, which is an important step on the way to the possible formation of intramolecular frustrated Lewis acid base pairs (FLPs). The FLP is formed by the combination of an electron-deficient electrophilic boranyl ligand and the $[Ge_9]$ cluster core comprising electron lone pairs at each ligand-free Ge vertex atom with Lewis base properties (Scheme 1, **B**).

In order to obtain a deeper understanding of the electronic situation we carried out quantum-chemical calculations at the DFT-PBE0/TZVP level of theory^[35] using the TURBOMOLE program package.^[36] As an initial structure model for the $[Ge_9]$ cluster the single crystal data of compounds **2a-Cu** and **3a-Cu** were used, the $[NHC-Cu]^+$ fragments were deleted, and a COSMO solvent field was applied to counter the anionic charge.^[37] As depicted in

Figure 3 the HOMO and LUMO of **2a** are located at the $[\text{Ge}_9]$ cluster, and it is not until LUMO + 4 that the LUMO becomes boron-centered (for more MO diagrams see Supporting Information). Similar findings were obtained for anion **3a**. The calculated HOMO–LUMO gaps are 3.8 eV for anions **1a–3a**, which is in a similar range as reported for 1,3,2,5-diazadiborinines (up to 3.7 eV), which are capable of activating small molecules like CO_2 , ketones and alkenes.^[38] In order to investigate the impact of the ligand design on the localization of the HOMO and LUMO, further theoretical studies were carried out, in which the diazaborolidine ligand is substituted by a $-\text{BCy}_2$ group ($[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{BCy}_2]^-$, **4a**). Indeed, we observed a shift in the localization of the LUMO, which as for a FLP required becomes boron-based, and the HOMO–LUMO gap decreases to 3.4 eV. This arrangement of orbitals could be favorable to facilitate interactions with small molecules in a frustrated Lewis acid–base type manner, with the cluster acting as the Lewis base (HOMO) and the boranyl ligand as the Lewis acid (LUMO). Interestingly, also the

HOMO-2 of **4a**, which is energetically only 0.5 eV lower in energy than the HOMO, allows for another spatial orientation to interact with a Lewis acid. Investigations on the targeted synthesis of **4a** and other derivatives are ongoing. We found for example that the sterically more demanding arylborane Mes_2BBR remained unreactive in dioxane solution.

At this point it has to be mentioned that Roesky et al. previously reported an intramolecular Ge/B Lewis pair which, however, showed Ge–B interactions in the ground state (no FLP).^[33b] Due to their unique reactivity, FLPs have recently gained a lot of attention with respect to the activation of small molecules (CO_2 , H_2), and have also been applied as metal-free hydrogenation catalysts for imines or enamines.^[39] Furthermore, phosphine-borane based FLPs have been reported to insert sterically demanding azides.^[40] These reports hold promise for further investigations at the here presented systems. The unique three-dimensional structure of the $[\text{Ge}_9]$ core opens further possibilities for the spatial arrangement of Lewis-acid Lewis-base pairs. In this context we are currently investigating the addition of a phosphanyl group adjacent to a boranyl group (Scheme 1, C).

In summary we report on the synthesis of the first boranyl functionalized $[\text{Ge}_9]$ clusters $[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{DAB}^{\text{R}}]^-$ [R: Me (**1a**), *i*Pr (**2a**), *o*-tol (**3a**)]. The addition of a DAB ligand to the $[\text{Ge}_9]$ core is limited by steric factors (no reaction of bulky $\text{DAB}^{\text{Mes}}\text{-Cl}$). Subsequent reactions with $\text{NHC}^{\text{Dipp}}\text{CuCl}$ led to the formation of $\text{NHC}^{\text{Dipp}}\text{Cu}[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{DAB}^{\text{R}}]^-$ [R: *i*Pr (**2a-Cu**) and *o*-tol (**3a-Cu**)]. The structural characterization is in accordance with the spectroscopic data and confirms the presence of a localized Ge–B *exo*-cluster bond. Moreover, Ge–B bond cleavage during the analogous reaction with **1a** reveals an increased lability for DAB ligands with small wingtips (cleavage of DAB^{Me}). Molecular orbital analyses for the anions **1a–3a** using DFT calculations show $[\text{Ge}_9]$ based HOMOs and LUMOs. The isolation of the anions **1a–3a** clearly accounts for the possibility of introducing boranyl moieties to the cluster core. Moreover, quantum-chemical calculations reveal the possibility of altering the position of the LUMO by changing the boranyl group to $-\text{BCy}_2$ (**4a**), which might open the way to $[\text{Ge}_9]$ cluster-based FLP systems. Cyclic-ether ring opening and FLP-type nitrile activation of halo-borane nonagermanide-cluster Lewis acid-base-pairs will be reported in a forthcoming paper.^[41]

Acknowledgements

We thank the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) (project number 245845833) within International Research Training Group IRTG 2022—Alberta Technical University of Munich School for Functional Hybrid Materials (ATUMS) and the Bavarian Ministry of Economic Affairs, Regional Development and Energy within the project “SolarTechnologies go Hybrid” for funding. Support within TUM IGSSE is greatly appreciated. C. W. also thanks the Studienstiftung des Deutschen Volkes for granting a PhD scholarship. A. J. K. acknowledges funding from the Academy of Finland (grant 317273) and computational resources from CSC—the Finnish IT Center for




Figure 3. Selected molecular orbitals of a) the anion $[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{DAB}^{\text{iPr}}]^-$ (**2a**) and b) the theoretical anion $[\text{Ge}_9\{\text{Si}(\text{TMS})_3\}_2\text{BCy}_2]^-$ (**4a**). The calculations were carried out on the DFT-PBE0/TZVP level of theory, using the single crystal structure data of **2a-Cu** and **3a-Cu** with deleted $[\text{NHC-Cu}]^+$ fragments as a starting point. Orbital figures are drawn so that 50% of the density is enclosed within the isosurface, corresponding to approximately 0.04 a.u. isovalue.

Science. Open access funding enabled and organized by Projekt DEAL.

Conflict of interest

The authors declare no conflict of interest.

Keywords: boranes · cluster compounds · germanium · Lewis base · Zintl cluster

- [1] a) S. C. Sevov, J. M. Goicoechea, *Organometallics* **2006**, *25*, 5678–5692; b) S. Scharfe, F. Kraus, S. Stegmaier, A. Schier, T. F. Fässler, *Angew. Chem. Int. Ed.* **2011**, *50*, 3630–3670; *Angew. Chem.* **2011**, *123*, 3712–3754; c) K. Mayer, J. Weßing, T. F. Fässler, R. A. Fischer, *Angew. Chem. Int. Ed.* **2018**, *57*, 14372–14393; *Angew. Chem.* **2018**, *130*, 14570–14593; d) B. Weinert, S. Mitzinger, S. Dehnen, *Chem. Eur. J.* **2018**, *24*, 8470–8490; e) C. Liu, Z.-M. Sun, *Coord. Chem. Rev.* **2019**, *382*, 32–56; f) R. J. Wilson, N. Lichtenberger, B. Weinert, S. Dehnen, *Chem. Rev.* **2019**, *119*, 8506–8554.
- [2] a) O. Kysliak, A. Schnepf, *Dalton Trans.* **2016**, *45*, 2404–2408; b) F. Li, S. C. Sevov, *Inorg. Chem.* **2012**, *51*, 2706–2708.
- [3] a) L. J. Schiegerl, F. S. Geitner, C. Fischer, W. Klein, T. F. Fässler, *Z. Anorg. Allg. Chem.* **2016**, *642*, 1419–1426; b) O. Kysliak, T. Kunz, A. Schnepf, *Eur. J. Inorg. Chem.* **2017**, 805–810; c) O. Kysliak, C. Schrenk, A. Schnepf, *Inorg. Chem.* **2015**, *54*, 7083–7088.
- [4] a) L. G. Perla, A. Muñoz-Castro, S. C. Sevov, *J. Am. Chem. Soc.* **2017**, *139*, 15176–15181; b) L. G. Perla, S. C. Sevov, *J. Am. Chem. Soc.* **2016**, *138*, 9795–9798.
- [5] F. S. Geitner, J. V. Dums, T. F. Fässler, *J. Am. Chem. Soc.* **2017**, *139*, 11933–11940.
- [6] F. S. Geitner, W. Klein, T. F. Fässler, *Angew. Chem. Int. Ed.* **2018**, *57*, 14509–14513; *Angew. Chem.* **2018**, *130*, 14717–14721.
- [7] a) F. S. Geitner, T. F. Fässler, *Chem. Commun.* **2017**, 53, 12974–12977; b) L. J. Schiegerl, A. J. Karttunen, W. Klein, T. F. Fässler, *Chem. Eur. J.* **2018**, *24*, 19171–19174; c) S. Frischhut, T. F. Fässler, *Dalton Trans.* **2018**, *47*, 3223–3226.
- [8] F. Li, S. C. Sevov, *J. Am. Chem. Soc.* **2014**, *136*, 12056–12063.
- [9] S. Frischhut, W. Klein, M. Drees, T. F. Fässler, *Chem. Eur. J.* **2018**, *24*, 9009–9014.
- [10] C. Wallach, F. S. Geitner, W. Klein, T. F. Fässler, *Chem. Eur. J.* **2019**, *25*, 12349–12356.
- [11] F. F. Li, A. Muñoz-Castro, S. C. Sevov, *Angew. Chem. Int. Ed.* **2012**, *51*, 8581–8584; *Angew. Chem.* **2012**, *124*, 8709–8712.
- [12] F. S. Geitner, W. Klein, O. Storcheva, T. D. Tilley, T. F. Fässler, *Inorg. Chem.* **2019**, *58*, 13293–13298.
- [13] C. Schenk, A. Schnepf, *Chem. Commun.* **2009**, 3208–3210.
- [14] K. Mayer, W. Klein, T. F. Fässler, *Chem. Commun.* **2019**, 55, 12156–12159.
- [15] N. C. Michenfelder, C. Gienger, A. Schnepf, A.-N. Unterreiner, *Dalton Trans.* **2019**, *48*, 15577–15582.
- [16] K. Mayer, L. J. Schiegerl, T. Kratky, S. Günther, T. F. Fässler, *Chem. Commun.* **2017**, 53, 11798–11801.
- [17] O. Kysliak, C. Schrenk, A. Schnepf, *Inorg. Chem.* **2017**, *56*, 9693–9697.
- [18] S. Frischhut, W. Klein, T. F. Fässler, *C. R. Chim.* **2018**, *21*, 932–937.
- [19] O. Kysliak, A. Schnepf, *Z. Anorg. Allg. Chem.* **2019**, *645*, 335–339.
- [20] F. S. Geitner, C. Wallach, T. F. Fässler, *Chem. Eur. J.* **2018**, *24*, 4103–4110.
- [21] a) Y. Heider, N. E. Poitiers, P. Willmes, K. I. Leszczyńska, V. Huch, D. Scheschkewitz, *Chem. Sci.* **2019**, *10*, 4523–4530; b) P. Willmes, K. Leszczyńska, Y. Heider, K. Abersfelder, M. Zimmer, V. Huch, D. Scheschkewitz, *Angew. Chem. Int. Ed.* **2016**, *55*, 2907–2910; *Angew. Chem.* **2016**, *128*, 2959–2963.
- [22] Y. Heider, P. Willmes, V. Huch, M. Zimmer, D. Scheschkewitz, *J. Am. Chem. Soc.* **2019**, *141*, 19498–19504.
- [23] A. V. Protchenko, J. Urbano, J. A. Abdalla, J. Campos, D. Vidovic, A. D. Schwarz, M. P. Blake, P. Mountford, C. Jones, S. Aldridge, *Angew. Chem. Int. Ed.* **2017**, *56*, 15098–15102; *Angew. Chem.* **2017**, *129*, 15294–15298.
- [24] a) Y. Segawa, Y. Suzuki, M. Yamashita, K. Nozaki, *J. Am. Chem. Soc.* **2008**, *130*, 16069–16079; b) L. M. Saleh, K. H. Birj Kumar, A. V. Protchenko, A. D. Schwarz, S. Aldridge, C. Jones, N. Kaltsoyannis, P. Mountford, *J. Am. Chem. Soc.* **2011**, *133*, 3836–3839.
- [25] A. V. Protchenko, K. H. Birj Kumar, D. Dange, A. D. Schwarz, D. Vidovic, C. Jones, N. Kaltsoyannis, P. Mountford, S. Aldridge, *J. Am. Chem. Soc.* **2012**, *134*, 6500–6503.
- [26] a) A. Rit, J. Campos, H. Niu, S. Aldridge, *Nat. Chem.* **2016**, *8*, 1022; b) M. Usher, A. V. Protchenko, A. Rit, J. Campos, E. L. Kolychev, R. Tirfoin, S. Aldridge, *Chem. Eur. J.* **2016**, *22*, 11685–11698; c) A. V. Protchenko, M. P. Blake, A. D. Schwarz, C. Jones, P. Mountford, S. Aldridge, *Organometallics* **2015**, *34*, 2126–2129; d) Z. Dong, H. H. Cramer, M. Schmidtmann, L. A. Paul, I. Siewert, T. Müller, *J. Am. Chem. Soc.* **2018**, *140*, 15419–15424; e) R. J. Mangan, A. Rit, C. P. Sindlinger, R. Tirfoin, J. Campos, J. Hicks, K. E. Christensen, H. Niu, S. Aldridge, *Chem. Eur. J.* **2020**, *26*, 306–315.
- [27] a) A. V. Protchenko, J. I. Bates, L. M. Saleh, M. P. Blake, A. D. Schwarz, E. L. Kolychev, A. L. Thompson, C. Jones, P. Mountford, S. Aldridge, *J. Am. Chem. Soc.* **2016**, *138*, 4555–4564; b) A. V. Protchenko, D. Dange, M. P. Blake, A. D. Schwarz, C. Jones, P. Mountford, S. Aldridge, *J. Am. Chem. Soc.* **2014**, *136*, 10902–10905.
- [28] A. V. Protchenko, D. Dange, A. D. Schwarz, C. Y. Tang, N. Phillips, P. Mountford, C. Jones, S. Aldridge, *Chem. Commun.* **2014**, 50, 3841–3844.
- [29] F. S. Geitner, M. A. Giebel, A. Pöthig, T. F. Fässler, *Molecules* **2017**, *22*, 1204.
- [30] A. Schnepf, *Angew. Chem. Int. Ed.* **2003**, *42*, 2624–2625; *Angew. Chem.* **2003**, *115*, 2728–2729.
- [31] T. Gädt, J.-A. Dimmer, S. Fleischhauer, A. Frank, C. Nickl, T. Wütz, K. Eichele, L. Wesemann, *Dalton Trans.* **2015**, *44*, 4726–4731.
- [32] N. S. Hosmane, J. Yang, K.-J. Lu, H. Zhang, U. Siriwardane, M. S. Islam, J. L. Thomas, J. A. Maguire, *Organometallics* **1998**, *17*, 2784–2796.
- [33] a) Y. Wang, M. Karni, S. Yao, Y. Apeloig, M. Driess, *J. Am. Chem. Soc.* **2018**, *140*, 1655–1664; b) J. Li, B. Li, R. Liu, L. Jiang, H. Zhu, H. W. Roesky, S. Dutta, D. Koley, W. Liu, Q. Ye, *Chem. Eur. J.* **2016**, *22*, 14499–14503.
- [34] F. S. Geitner, T. F. Fässler, *Eur. J. Inorg. Chem.* **2016**, 2688–2691.
- [35] a) C. Adamo, V. Barone, *J. Chem. Phys.* **1999**, *110*, 6158–6170; b) J. P. Perdew, K. Burke, M. Ernzerhof, *Phys. Rev. Lett.* **1996**, *77*, 3865; c) F. Weigend, R. Ahlrichs, *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297–3305.
- [36] TURBOMOLE V7.3 2018, a development of University of Karlsruhe and Forschungszentrum Karlsruhe GmbH, **1989–2007**, TURBOMOLE GmbH, since 2007; available from www.turbomole.com.
- [37] A. Klamt, G. Schüürmann, *J. Chem. Soc. Perkin Trans. 2* **1993**, *5*, 799–805.
- [38] a) Y. Su, Y. Li, R. Ganguly, R. Kinjo, *Angew. Chem. Int. Ed.* **2018**, *57*, 7846–7849; *Angew. Chem.* **2018**, *130*, 7972–7975; b) D. Wu, R. Wang, Y. Li, R. Ganguly, H. Hirao, R. Kinjo, *Chem* **2017**, *3*, 134–151; c) D. Wu, L. Kong, Y. Li, R. Ganguly, R. Kinjo, *Nat. Commun.* **2015**, *6*, 1–6; d) D. Wu, R. Ganguly, Y. Li, S. N. Hoo, H. Hirao, R. Kinjo, *Chem. Sci.* **2015**, *6*, 7150–7155.

- [39] a) D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.* **2010**, *49*, 46–76; *Angew. Chem.* **2010**, *122*, 50–81; b) D. W. Stephan, *J. Am. Chem. Soc.* **2015**, *137*, 10018–10032; c) D. W. Stephan, *Science* **2016**, *354*, aaf7229; d) Z. Huang, S. Wang, R. D. Dewhurst, N. V. Ignat'ev, M. Finze, H. Braunschweig, *Angew. Chem. Int. Ed.* **2020**, *59*, 8800–8816; *Angew. Chem.* **2020**, *132*, 8882–8900.
- [40] a) C. M. Mömming, G. Kehr, B. Wibbeling, R. Fröhlich, G. Erker, *Dalton Trans.* **2010**, *39*, 7556–7564; b) A. Stute, L. Heletta, R. Fröhlich, C. G. Daniliuc, G. Kehr, G. Erker, *Chem. Commun.* **2012**, *48*, 11739–11741; c) L. M. Elmer, G. Kehr, C. G. Daniliuc, M. Siedow, H. Eckert, M. Tesch, A. Studer, K. Williams, T. H. Warren, G. Erker, *Chem. Eur. J.* **2017**, *23*, 6056–6068; d) D. H. Boom, A. R. Jupp, M. Nieger, A. W. Ehlers, J. C. Sootweg, *Chem. Eur. J.* **2019**, *25*, 13299–13308.
- [41] F. S. Geitner, C. Wallach, T. F. Fässler, submitted for publication.

Manuscript received: September 10, 2020
Revised manuscript received: October 3, 2020
Accepted manuscript online: October 14, 2020
Version of record online: December 1, 2020