
This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Kedves, Orsolya; Kocsubé, Sándor; Bata, Teodóra; Andersson, Maria A.; Salo, Johanna M.; Mikkola, Raimo; Salonen, Heidi; Szűcs, Attila; Kedves, Alfonz; Kónya, Zoltán; Vágvolgyi, Csaba; Magyar, Donát; Kredics, László

Chaetomium and chaetomium-like species from European indoor environments include *Dichotomopilus finlandicus* sp. nov.

Published in:
Pathogens

DOI:
[10.3390/pathogens10091133](https://doi.org/10.3390/pathogens10091133)

Published: 01/09/2021

Document Version
Publisher's PDF, also known as Version of record

Published under the following license:
CC BY

Please cite the original version:
Kedves, O., Kocsubé, S., Bata, T., Andersson, M. A., Salo, J. M., Mikkola, R., Salonen, H., Szűcs, A., Kedves, A., Kónya, Z., Vágvolgyi, C., Magyar, D., & Kredics, L. (2021). Chaetomium and chaetomium-like species from European indoor environments include *Dichotomopilus finlandicus* sp. nov. *Pathogens*, 10(9), Article 1133. <https://doi.org/10.3390/pathogens10091133>

Article

Chaetomium and *Chaetomium*-like Species from European Indoor Environments Include *Dichotomopilus finlandicus* sp. nov.

Orsolya Kedves ¹, Sándor Kocsubé ¹, Teodóra Bata ¹, Maria A. Andersson ², Johanna M. Salo ², Raimo Mikkola ² , Heidi Salonen ², Attila Szűcs ¹ , Alfonz Kedves ³, Zoltán Kónya ³ , Csaba Vágvolgyi ¹ , Donát Magyar ⁴  and László Kredics ^{1,*} 

- ¹ Department of Microbiology, Faculty of Science and Informatics, University of Szeged, Közép fasor 52, H-6726 Szeged, Hungary; kedvesorsolya91@gmail.com (O.K.); kocsube@bio.u-szeged.hu (S.K.); bata.teodora1997@gmail.com (T.B.); a.szunyi@windowslive.com (A.S.); csaba@bio.u-szeged.hu (C.V.)
 - ² Department of Civil Engineering, Aalto University, P.O. Box 12100, FI-00076 Aalto, Finland; maria.a.andersson@helsinki.fi (M.A.A.); johanna.72salos@gmail.com (J.M.S.); raimo.mikkola@aalto.fi (R.M.); heidi.salonen@aalto.fi (H.S.)
 - ³ Department of Applied and Environmental Chemistry, Faculty of Science and Informatics, University of Szeged, Rerrich Béla tér 1, H-6720 Szeged, Hungary; kedvesalfonz@yahoo.com (A.K.); konya@chem.u-szeged.hu (Z.K.)
 - ⁴ National Public Health Center, Albert Flórián út 2-6, H-1097 Budapest, Hungary; magyar.donat@gmail.com
- * Correspondence: kredics@bio.u-szeged.hu; Tel.: +36-62544516



Citation: Kedves, O.; Kocsubé, S.; Bata, T.; Andersson, M.A.; Salo, J.M.; Mikkola, R.; Salonen, H.; Szűcs, A.; Kedves, A.; Kónya, Z.; et al. *Chaetomium* and *Chaetomium*-like Species from European Indoor Environments Include *Dichotomopilus finlandicus* sp. nov. *Pathogens* **2021**, *10*, 1133. <https://doi.org/10.3390/pathogens10091133>

Academic Editor: Lawrence S. Young

Received: 2 August 2021

Accepted: 1 September 2021

Published: 3 September 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The genus *Chaetomium* is a frequently occurring fungal taxon world-wide. *Chaetomium* and *Chaetomium*-like species occur in indoor environments, where they can degrade cellulose-based building materials, thereby causing structural damage. Furthermore, several species of this genus may also cause adverse effects on human health. The aims of this research were to identify *Chaetomium* and *Chaetomium*-like strains isolated from indoor environments in Hungary and Finland, two geographically distant regions of Europe with drier and wetter continental climates, respectively, and to study their morphological and physiological properties, as well as their extracellular enzyme activities, thereby comparing the *Chaetomium* and *Chaetomium*-like species isolated from these two different regions of Europe and their properties. *Chaetomium* and *Chaetomium*-like strains were isolated from flats and offices in Hungary, as well as from schools, flats, and offices in Finland. Fragments of the translation elongation factor 1 α (*tef1 α*), the second largest subunit of RNA polymerase II (*rpb2*) and β -tubulin (*tub2*) genes, as well as the internal transcribed spacer (ITS) region of the ribosomal RNA gene cluster were sequenced, and phylogenetic analysis of the sequences performed. Morphological examinations were performed by stereomicroscopy and scanning electron microscopy. Thirty-one *Chaetomium* sp. strains (15 from Hungary and 16 from Finland) were examined during the study. The most abundant species was *Ch. globosum* in both countries. In Hungary, 13 strains were identified as *Ch. globosum*, 1 as *Ch. cochliodes*, and 1 as *Ch. interruptum*. In Finland, 10 strains were *Ch. globosum*, 2 strains were *Ch. cochliodes*, 2 were *Ch. rectangulare*, and 2 isolates (SZMC 26527, SZMC 26529) proved to be representatives of a yet undescribed phylogenetic species from the closely related genus *Dichotomopilus*, which we formally describe here as the new species *Dichotomopilus finlandicus*. Growth of the isolates was examined at different temperatures (4, 15, 20, 25, 30, 37, 35, 40, and 45 °C), while their extracellular enzyme production was determined spectrophotometrically.

Keywords: *Chaetomium*; *Dichotomopilus*; indoor environment; extracellular enzymes

1. Introduction

Chaetomium Kunze (Ascomycota, Sordariales) is the largest genus of the family Chaetomiaceae, present in various substrates and geographical regions [1]. More than 400 *Chaetomium* species have been described. The type species is *Ch. globosum* Kunze [2].

As cellulose-degrading fungi they possess the ability to degrade wet cellulosic building materials such as wood and plywood and synthetic building materials such as plastics and drywall [3]. Wet building materials are dominated by colonization and forming a dense mycelium covering most of the building materials, thereby damaging the building structure [4,5]. In addition to colonizing building materials, members of the genus *Chaetomium* are also known to produce more than 500 bioactive metabolites [6]. *Chaetomium globosum*, producing various mycotoxins—such as chaetoglobosin, chaetomin, chaetomugilin, and chaetoviridine—is the most common species of the genus in indoor environments [5,7].

Several *Chaetomium* species have been described to cause onychomycosis [8–12], skin surface infections [13], and cerebral mycosis [14–16]. Among them, some are able to opportunistically cause systemic infections and trigger severe allergic reactions that increase the risk of developing asthma [7]. On the other hand, some species of the genus *Chaetomium* are plant endophytes [17–20], producing high levels of various enzymes [21–24], antioxidants [25,26], as well as antifungal [25,27–31], antibacterial [28,32–37], and nematocidal metabolites [38,39], possessing anticancer [26,28,30,40–51] and anti-inflammatory activities [39,52], and being able to biosynthesize several types of nanoparticles [44,53,54]. The genus includes psychrotolerant [55], mesophilic, thermotolerant, and thermophilic species [5], which are widely used in the medicinal and industrial field (e.g., food, textile, and fermentation industries), agriculture (as biocontrol agents and in agricultural waste degradation), and waste processing (composting) [56].

The genus *Chaetomium* is generally characterized by rounded, ovoid, or obovate ostiolate ascomata covered with characteristic hairs. The walls of the ascomata are usually *textura intricata* (tissue of interwoven irregularly disposed hyphae with distinct interhyphal spaces, the walls not united), but they occur in *textura angularis* (tissue of short polyhedral cells without intercellular spaces). Ascomatal hairs can be straight (seta-like), flexible, curved, wavy, circulating, spirally curled, or otherwise branched in various morphologies. Asci are clavate or fusiform with 8 biseriate or irregularly arranged ascospores, evanescent. Ascospores are limoniform to globose, or irregular in a few species, bilaterally flattened, usually more than 7 µm in length. Asexual morphs, if present, are *Acremonium*-like [7].

Chaetomium is an intensively studied fungal genus worldwide, as it causes one of the biggest problems in indoor environments with damaging effects both to buildings and human health [3,4,7,54,57–61]. The closely related genus *Dichotomopilus* was first described by Wang et al. [7]; until then, members of the genus *Dichotomopilus* belonged to the genus *Chaetomium*. The genus name *Dichotomopilus* refers to the shape of terminal hairs of the ascomata, which are usually dichotomously branched. So far, this genus has included 12 species.

To the best of our knowledge, the diversity of this group of fungi has not yet been studied in Hungary. This study aimed to identify *Chaetomium* and *Chaetomium*-like strains isolated from Hungary and Finland, two geographically distant regions of Europe with drier and wetter continental climates, respectively, to study their morphology, determine their physiological properties, measure their extracellular enzyme activities, and compare the properties of *Chaetomium* and *Chaetomium*-like strains isolated from different sites of the two regions.

2. Results

Based on *tef1α* sequences, the most abundant species in this study was *Ch. globosum* in both countries (Table 1, Figure 1). In Hungary, 13 strains were identified as *Ch. globosum*, 1 as *Ch. cochliodes*, and 1 as *Ch. interruptum*, while in Finland, 10 strains were *Ch. globosum*, 2 strains were *Ch. cochliodes*, 2 were *Ch. rectangulare*, and 2 isolates (SZMC 26527, SZMC 26529) proved to be representatives of a yet undescribed phylogenetic species from the closely related genus *Dichotomopilus* (Figure 1).

Table 1. *Chaetomium* and *Chaetomium*-like isolates, reference strains and their sequences involved in the study.

Species	Collection Number	Lab Code	Location of Isolation	<i>tefla</i>	GenBank Accession Number		
					ITS	<i>rpb2</i>	<i>tub2</i>
<i>Ac. globosum</i> *	CBS 332.67 T		Rhizosphere, Lucknow, India	KM655479			
<i>Ac. strumarium</i> *	CBS 333.67 T		Soil, Lucknow, India	KC503252			
<i>Ch. afropilosum</i> *	CBS 145.38 T		Unknown	KT214713			
<i>Ch. angustispirale</i> *	CBS 137.58 T		<i>Fraxinus</i> sp., Tellerman forest, Baleshev region, Russia	KF001734			
<i>Ch. ascotrichoides</i> *	CBS 113.83		<i>Gossypium humitectum</i> , Argentina	KF001742			
<i>Ch. capillare</i> *	CBS 128489 T		Animal hair, California, USA	KT214724	NR_144860	KT214686	KT214760
<i>Ch. cervicicola</i> *	CBS 128492 T		Neck of <i>Homo sapiens</i> Texas, USA	KT214697			
<i>Ch. citrinum</i> *	CBS 693.82 T		Rice field soil, Tochigi, Japan	KT214730	NR_144863	KT214691	KT214764
<i>Ch. coarctatum</i> *	CBS 162.62 T		Seed of <i>Cappanula medium</i> , St. Petersburg, Russia,	KF001712			
<i>Ch. cochliodes</i>	SZMC 22473	T364	Wall (swab), room, apartment Tatabánya, Hungary	MW556657			
<i>Ch. cochliodes</i>	SZMC 24764	OT7 **	Settled dust, office, Helsinki, Finland	MT498103			
<i>Ch. cochliodes</i>	SZMC 26528	OT7b **	Settled dust, office, Helsinki, Finland	MT498102			
<i>Ch. cochliodes</i> *	CBS 155.52 T		Animal dung, USA	KF001721	NR_151835	KF001811	KC109772
<i>Ch. contagiosum</i> *	CBS 128494 T		Cornea of <i>Homo sapiens</i> , Northeast USA	KT214694	NR_144846	KT214659	KT214732
<i>Ch. cucurmericola</i> *	CBS 378.71 T		Izmir, Turkey	KT214718			
<i>Ch. elatum</i> *	CBS 374.66 T		Decomposing leaf, Aptos, California, USA	KF001730			
<i>Ch. elatum</i> *	CBS 142034 T		Cardboard, Denmark		KX976612	KX976832	KX976954
<i>Ch. fimeti</i> *	CBS 139034 T		Soil, Germany	KT214698			
<i>Ch. globosporum</i> *	CBS 108.83 T		Green leaf of <i>Triticum aestivum</i> , Germany	KF001735			
<i>Ch. globosum</i>	SZMC 22474	T365	Air, hotel, Budapest, Hungary	MW556658			
<i>Ch. globosum</i>	SZMC 22478	T369	Air, shop, Szentendre, Hungary	MW556659	MW541923		
<i>Ch. globosum</i>	SZMC 22481	T372A	House dust, basement, flat, Budapest, Hungary	MW556660			
<i>Ch. globosum</i>	SZMC 22788	T428B	Ceiling (swab), apartment, Budapest, Hungary	MW556661			
<i>Ch. globosum</i>	SZMC 23266	T457D	Under ceramic tiles in a kitchen (swab), apartment, Csepel, Hungary	MW556662			
<i>Ch. globosum</i>	SZMC 23275	T459A	Air, from gypsum board wall of a children's room, house, Budapest, Hungary	MW556663			
<i>Ch. globosum</i>	SZMC 23688	T499	House dust (swab), living room, apartment, Budapest, Hungary	MW556664			
<i>Ch. globosum</i>	SZMC 24035	T536A	Wall, kitchen, apartment, Budapest, Hungary	MW556665			
<i>Ch. globosum</i>	SZMC 24451	C13/LM	Exhaust air filter, school, Vantaa, Finland	MW556666			
<i>Ch. globosum</i>	SZMC 24455	C22/LM **	Exhaust air filter, school, Vantaa, Finland	MT498109			
<i>Ch. globosum</i>	SZMC 24456	MH5 **	Settled dust, public building, Espoo, Finland	MT498108			
<i>Ch. globosum</i>	SZMC 24464	MÖ9 **	Settled dust, piggery, Orimattila, Finland	MT498106	MW541924		

Table 1. Cont.

Species	Collection Number	Lab Code	Location of Isolation	GenBank Accession Number			
				<i>tef1a</i>	ITS	<i>rpb2</i>	<i>tub2</i>
<i>Ch. globosum</i>	SZMC 24508	T582D	Garage wall (swab), apartment, Budapest, Hungary	MW556667			
<i>Ch. globosum</i>	SZMC 24765	2c/26	Settled dust, apartment, Vantaa, Finland	MW310244			
<i>Ch. globosum</i>	SZMC 24766	2b/26 **	Settled dust, apartment, Vantaa, Finland	MT498110			
<i>Ch. globosum</i>	SZMC 24768	C22	Settled dust, apartment Vantaa, Finland	MW556668			
<i>Ch. globosum</i>	SZMC 24769	MH52 **	Settled dust, public building, Espoo, Finland	MT498107			
<i>Ch. globosum</i>	SZMC 24938	626C	Wall (swab), living room, house, Kazincbarcika, Hungary	MW556670			
<i>Ch. globosum</i>	SZMC 26530	Ruk10 **	Settled dust, apartment, Vantaa, Finland	MT498101	MW541927		
<i>Ch. globosum</i>	SZMC 26534	MTAV35 **	Settled dust, University of Oulu, Finland				
<i>Ch. globosum</i>	SZMC 26539	3b/APP	Exhaust air filter, public building, Espoo, Finland	MW588207			
<i>Ch. globosum</i>	SZMC 26845	T706	Wall (swab), kitchen, apartment, Budapest, Hungary	MW556672			
<i>Ch. globosum</i>	SZMC 26857	T711	Wallpaper (swab), living room, apartment, Budapest, Hungary	MW556673			
<i>Ch. globosum</i>	SZMC 27052	T730A	On paper packaging, imported from Sri Lanka, swab factory, Pécs, Hungary	MW556674			
<i>Ch. globosum</i> *	CBS 160.62 T		Compost, Germany	KT214704	NR_144851	KT214666	KT214742
<i>Ch. globosum</i> *	MUCL 39526 T		Dead stem of <i>Juncus</i> sp., Hungary	KF001710			
<i>Ch. globosum</i> *	CBS 666.82		Unknown		KX976617	KX976833	KX976959
<i>Ch. graminiforme</i> *	CBS 506.84 T		<i>Acer</i> sp., Muskoka District, Ontario, Canada	KT214725			
<i>Ch. grande</i> *	CBS 126780 T		Leaf of <i>Triticum aestivum</i> , Naghadeh, Iran	KT214692			
<i>Ch. interruptum</i>	SZMC 23937	T531B	Under wallpaper (swab), office, Budapest, Hungary	MW588206	MW301425		
<i>Ch. interruptum</i> *	CBS 126660 T		Seed of <i>Triticum aestivum</i> , Hadishahr East Azerbaijan Province, Iran	KT214703	KT214564	KT214665	
<i>Ch. madrasense</i> *	CBS 315.74 T		Rhizosphere of <i>Pennisetum typhoides</i> , Chennai, Tamil Nadu, India	KF001741			
<i>Ch. novozelandicum</i> *	CBS 124555 T		Dead decaying twig, Otaki, New Zealand	KT214715			
<i>Ch. pilosum</i> *	CBS 335.67 T		Grain of <i>Triticum aestivum</i> , Perth, Western Australia	KT214729	NR_144862	FJ666387	KT214763
<i>Ch. pseudocochliodes</i> *	CGMCC 3.9441 T		Roots of <i>Panax notoginseng</i> , Wenshan, Yunnan Province, China	KF001726			
<i>Ch. pseudoglobosum</i> *	CBS 574.71 T		Unknown	KT214712			
<i>Ch. rectangulare</i>	SZMC 26531	MO13 **	Settled dust, piggery, Orimattila, Finland	MT498104	MW541928		
<i>Ch. rectangulare</i>	SZMC 26535	MO15 **	Settled dust, piggery, Orimattila, Finland	MT498105	MW541929		
<i>Ch. rectangulare</i> *	CBS 126778 T		Leaf of <i>Hordeum vulgare</i> , Salmas, West Azerbaijan province, Iran	KT214726	NR_144817	HM365285	KT214688
<i>Ch. spiculipilium</i> *	CBS 373.66 T		Decaying vegetable debris, California, USA	KF001719			
<i>Ch. spirochaete</i> *	CBS 730.84 T		Animal dung, Great Smokey Mountains, Tennessee, USA	KF001729	NR_144823	KF001819	JN256191

Table 1. Cont.

Species	Collection Number	Lab Code	Location of Isolation	<i>tef1a</i>	GenBank Accession Number		
					ITS	<i>rpb2</i>	<i>tub2</i>
<i>Ch. subaffine</i> *	CBS 637.91 T		Cereal, USSR	KF001727			
<i>Ch. subfimet</i> *	CBS 370.66 T		Paper and vegetable material, Cardiff, Wales	KT214701	NR_144850	FJ666385	KT214739
<i>Ch. subglobosum</i> *	CBS 149.60 T		Dead herbaceous stem, St. Petersburg, Russia	KF001718			
<i>Ch. telluricola</i> *	CBS 151.59 T		Soil, Suffolk, Lakenheath Warren, United Kingdom	KT214723			
<i>Ch. tenue</i> *	CBS 139.38 T		Unknown	KT214707			
<i>Ch. umbonatum</i> *	CBS 293.83 T		Soil, Nova Scotia, Canada	KT214714			
<i>Ch. undulatulum</i> *	CBS 126775 T		Leaf of <i>Hordeum vulgare</i> , Bonab, East Azerbaijan province, Iran	KT214720			
<i>Ch. unguicola</i> *	CBS 128446 T		Nail of <i>Homo sapiens</i> , Los Angeles, USA	KT214706			
<i>Ch. megalocarpum</i> *	CBS 149.59 T		Leaf of <i>Ficus carica</i> , Greece	KF001738			
<i>D. dolichotrichus</i> *	CBS 162.48 T		USA	KC485023	HM449049	KX976852	JF772462
<i>D. erectus</i> *	CBS 140.56 T		<i>Petroselinum sativum</i> , USA	KC485018	HM449044	KX976854	JF772458
<i>D. funicola</i> *	CBS 159.52 T		Germany	KC485013	GU563369	KX976856	JF772461
<i>D. funicola</i> *	CBS 136.38		Unknown			KX976857	
<i>D. funicola</i> *	DTO 333-F1		Dust, outdoors, Denmark		KX976658	KX976858	KX977000
<i>D. fusus</i> *	CBS 372.66 T		Unknown	KM655463	KM655333	KX976859	KX977002
<i>D. indicus</i> *	CGMCC 3.14184 T		Rhizosphere of <i>Panax Notoginseng</i> , Wenshan county, Yunnan Province	KC485005	GU563367	KX976861	JF772453
<i>D. pratensis</i> *	CGMCC 3.14181 T		Soil, Huangnan, Qinghai Province, China	KC485017	GU563372	KX976866	JF772450
<i>D. pseudoerectus</i> *	CBS 252.75 T		Air, Uttar Pradesh, India		NR_147674	KX976869	KX977009
<i>D. pseudofunicola</i> *	CBS 142033 T		Dust, USA		KX976668	KX976870	KX977010
<i>D. ramosissimus</i> *	CGMCC 3.14183 T		Rhizosphere of <i>Panax notoginseng</i> , Wenshan county, Yunnan Province, China	KC485021	GU563371	KX976871	JF772452
<i>D. reflexus</i> *	CBS 157.49 T		Germinating seed, USA	KC485027	HM449051	KX976873	JF772460
<i>D. subfunicola</i> *	CGMCC 3.12892 T		Soil, Shihezi, Xinjiang Autonomous Region, China	KC485014	JX867125	KX976875	JX867122
<i>D. subfunicola</i> *	CGMCC 3.9466		Rhizosphere of <i>Panax notoginseng</i> , Yunnan, China	KC485016	GU563368	KX976876	JF772446
<i>D. subfunicola</i> *	CBS 794.83		Paper, Switzerland		GU563368	KX976876	KX977013
<i>D. subfunicola</i> *	CBS 812.73		Pistol belt, New Guinea		KX976670	KX976877	KX977012
<i>D. variostiolatus</i> *	CBS 179.84		Tarpaulin, New Guinea		NR_147676	KX976879	KX977014
<i>D. variostiolatus</i> *	DTO 319-B9		Dust, Thailand		KX976674	KX976881	KX977016
<i>D. variostiolatus</i> *	DTO 319-A2		Dust, USA		KX976673	KX976880	KX977015
<i>Dichotomopilus</i> sp.	SZMC 26527	C5/LM	Exhaust air filter, school, Vantaa, Finland	MW556671	MW541925	MZ665530	MZ665528
<i>Dichotomopilus</i> sp.	SZMC 26529	Ch1/tu **	Inlet air filter, public building, Espoo, Finland	MT644127	MW541926	MZ665531	MZ665529

Strains isolated during this study are set in bold. * reference strain [2,7], ** described in [5].




Figure 1. Maximum likelihood phylogeny of the examined isolates (set in bold) based on translation elongation factor 1 α . T = ex-type, nT = ex-neotype, eT = ex-epitype. Numbers above branches are bootstrap values. Only values greater than 70% are shown.

2.1. Morphological Features of the Isolated *Chaetomium* Strains

Among the previously described *Chaetomium* species, *Ch. cochliodes* colonies grew rapidly on MEA, OA, and PDA (Figure 2A) reaching 65–70 mm in diameter after 7 days at 25 °C. Hyphae were light beige on MEA, while brownish on OA and PDA, with powdery surface, undulate colony edges and without colored exudates. The strains were unable to produce ascospore-containing ascomata on MEA, while strong dark green ascospore formation was observed after 7 days on OA and PDA. *Ch. interruptum* (Figure 2B) formed white mycelium on all media, brownish exudates diffusing into the media, and did not produce spores during 7 days of culturing at 25 °C. On MEA and OA, it formed regular circular colonies, while on PDA the edges of the colonies grew irregularly. Colony diameters

after 7 days were 40–45, 50–60, and 30–40 mm on MEA, OA, and PDA, respectively. *Ch. globosum* (Figure 2C) colonies overgrew both MEA and OA media in 7 days at 25 °C. On PDA the strains grew slowly, with colony diameters of 30–40 mm after 7 days and a lobate edge. No ascospores were produced on MEA medium, but greenish ascospores were produced on OA and PDA. Colonies ranged from beige (MEA, OA) to brown (PDA) in color, the surface texture was floccose or velvety, and a brownish exudate was produced on all media. *Ch. rectangularare* (Figure 2D) completely overgrew all media in 7 days at 25 °C with white, cottony mycelium and without colored exudates. No ascospores were produced under any of the conditions tested.




Figure 2. Colony morphology of *Chaetomium* strains on different agar media left to right: MEA, OA and PDA after seven days of incubation. (A). *Ch. cochliodes* SZMC 22473, (B). *Ch. interruptum* SZMC 23937, (C). *Ch. globosum* SZMC 23266, (D). *Ch. rectangularare* SZMC 26535.

Ascomata of *Ch. cochliodes* (Figure 3(A1–A5)) were ostiolate, ovoid, greenish olivaceous, with brown wall, *textura intricata*. Terminal hairs were usually around the ostium, light brown or brown, spirally coiled, lateral hairs undulate or loosely coiled, tapering towards the tip. Mature ascospores were brown, limoniform, usually biapiculate at both

Ascomata of *Ch. cochliodes* (Figure 3(A1–A5)) were ostiolate, ovoid, greenish olivaceous, with brown wall, *textura intricata*. Terminal hairs were usually around the ostiolum, light brown or brown, spirally coiled, lateral hairs undulate or loosely coiled, tapering towards the tip. Mature ascospores were brown, limoniform, usually biapiculate at both ends, bilaterally flattened. Ascomata of *Ch. globosum* (Figure 3(B1–B5)) were ostiolate, greenish olivaceous, with brown wall, *textura intricata*. Terminal hairs were light brown or brown, undulate to loosely coiled, lateral hairs brown, flexuous, tapering towards the tips. Mature ascospores were greenish or brown, subglobose or limoniform, bilaterally flattened. Ascomata of *Ch. interruptum* (Figure 3(C1–C5)) were ostiolate, brown, with brown wall, *textura epidermoidea* (tissue of closely interwoven irregularly disposed hyphae without interhyphal spaces, the walls united, usually forming a membranous or epidermis-like tissue). Terminal hairs were brown undulate, lateral hairs brown, flexuous, tapering towards the tips. Mature ascospores were greenish or brown, subglobose or limoniform, bilaterally flattened.




Figure 3. Ascomata of *Chaetomium* strains: (A1–A5): *Chaetomium cochliodes* SZMC 22473, (B1–B5): *Chaetomium globosum* SZMC 23266, (C1–C5): *Chaetomium interruptum* SZMC 23937. 1—ascomata side view; 2—ascomata top view; 3—ascomatal hairs; 4—ascospores; 5—ascomatal wall.

2.2. Phylogeny and Taxonomy

The *tefla*, *ITS*, *rpb2*, and *tub2* dataset consisted of 935, 639, 525, and 571 characters, respectively. The *tefla*, *ITS*, *rpb2*, and *tub2* dataset consisted of 935, 639, 525, and 571 characters, respectively. The indel-based binary dataset was 100 characters long. Isolates SZMC 26527 and SZMC 26529 resolved as members of a new species with high confidence values on the phylograms obtained from both *tefla* (Figure 1) and the other three loci (data not shown). For the final inference the four loci were concatenated and partitioned. Based on the maximum likelihood phylogenetic tree inferred from the concatenated sequences (Figure 4), isolates SZMC 26527 and SZMC 26529 formed a well-supported distinct branch inside the genus *Dichotomopilus* with the closest relatives being *D. funicola*, *D. pseudofunicola*, *D. subfunicola*, and *D. variostiolatus*. This new species is described below as *Dichotomopilus finlandicus* sp. nov.




Figure 4. Maximum likelihood phylogeny of the genus *Dichotomopilus* inferred from the concatenated translation elongation factor 1 α (*efl1*), internal transcribed spacer (ITS), second largest subunit of RNA polymerase II (*rpb2*), and *tub2* (*tub2*) sequences. T = ex-type, nT = ex-neotype, eT = ex-epitype. Numbers above branches are bootstrap values. Only values greater than 70% are shown.

Dichotomopilus finlandicus O. Kedves, S. Kocsubé, and L. Kredics sp. nov. MycoBank accession number: 840621. **Etymology:** Refers to the country of origin. Colonies on PDA (Figure 5(A1)) rapidly growing, about 51–54 mm in diameter after 7 days at 25°C, with a slightly undulate edge, usually with a flobose, white to cream mycelium, irregular connecting growth with colored exudates, and producing grey or black ascogmata in seven days. Colonies on MEA (Figure 5(A2)) rapidly growing, about 55–60 mm in diameter after 2 days at 25°C, with a lobate edge, not forming ascogmata in seven days. Colony color is yellowish-white; the surface texture folded velvety to flobose. Colonies slowly growing on OA (Figure 5(A3)) at 25°C, about 35–38 mm in diameter with a lobate edge, with a velvety surface texture. Culture color white, producing grey or black ascogmata in seven days. Colonies slowly growing on vegetable juice agar (media (Figure 5(A4))) about 31–33 mm in diameter over seven days at 25°C, with a slightly undulate edge, usually with a flobose, white to cream mycelium, without colored exudates, not producing ascogmata in seven days. On DG18 agar media (Figure 5(A5)) regular circular colonies showing weak growth, 12–13 mm in diameter. Colony color orange and white, producing orange exudates. **Ascomata** (Figure 5, B1–C3) superficial, ostiolate, subglobose to ovoid, dark brown, 150–180 μ m high and 110–130 μ m wide ascocarp. Ascumatal wall (Figure 5(D1)) comprising of brown, elongated, or irregular cells (*textura intricata*). Terminal hairs (Figure 5(C1–C3, D1)) usually around the ostiolum, light brown, or olivaceous brown, dichotomously branched 4–6 times, up to 250–320 μ m long, 3–4.5 μ m in diameter at the base, at wide angles and starting primarily from the upper half part, verrucose, regularly at the base, at wide angles and starting primarily from the upper half part, verrucose, septate. Lateral hairs unbranched, seta-like, tapering towards the tip. Asci (Figure 5(D3)) regularly septate. Lateral hairs unbranched, seta-like, tapering towards the tip. Asci (Figure 5(D3)) fasciculate, clavate and long stipitate, stalked, 8 irregularly-arranged ascospores, spore-

bearing portion $18\text{--}19 \times 7\text{--}8 \mu\text{m}$, stalks $6\text{--}11 \mu\text{m}$ long, evanescent. Ascospores: brown, broadly ellipsoid or almond-shaped $5.12\text{--}6.42 (5.88) \times 4.06\text{--}4.82 (4.42) \times 2.31\text{--}3.58 (2.82) \mu\text{m}$ (length \times width \times thickness). Growth temperature: optimum $25\text{--}30^\circ\text{C}$, minimum 15°C , and maximum 38°C . Specimens examined: A piece of inlet air filter ($2 \times 2 \text{ cm}$), public building, Espoo, Finland; Holotype: freeze dried culture specimen in the Szeged Microbiological Collection (SZMC) at the Department of Microbiology, Faculty of Science and Informatics, University of Szeged, Hungary, SZMC 26529; Non-sporulating strain: SZMC 26527 from a school building, Vantaa, Finland.

Pathogens 2021, 10, 1133

11 of 20




Figure 5. *Dichotomopilus finlandicus* SZMC 26529. (A1–A5): colony morphology of seven-day-old cultures on different agar media (A1—MEA, A2—PDA, A3—MEA, A4—PDA, A5—PDA). (B1–B3): morphology of ascomata on different agar media (left to right: OA, MEA, and PDA). (C1–C3): Ascomata C1, C2—MEA, A3—OA, A4—Vegetable juice agar, A5—PDA. (D1–D3): Asci and ascospores (Bars: D1–D3 50 μm). (E1–E6): Scanning electron-microscopic images of: E1,E2: ascomata; E3: terminal ascomatal hairs; E4,E5: ascomatal hairs and ascospores; E6: ascospores.

2.3. Physiological Characterization of the Isolated *Chaetomium* and *Chaetomium*-like Strains

The optimal growth temperature of all isolates was between 25–30 °C (Figure 6). The *Ch. globosum* strains grew at temperatures between 15–40 °C, they were unable to grow at 4 °C, and only three Hungarian *Ch. globosum* isolates (SZMC 22788, SZMC 24508, and SZMC 24938) grew at 40 °C (Figure 6A,B). In addition, four Hungarian isolates of *Ch. globosum* (SZMC 23266, SZMC 24938, SZMC 26845, and SZMC 26857) showed more intensive growth at 30 °C than at 25 °C (Figure 6B). *Ch. cochliodes* SZMC 22473 and SZMC 26528, *Ch. interruptum* SZMC 23937 and *Ch. rectangulare* SZMC 26535 grew at 4 °C but among these strains *Ch. cochliodes* SZMC 26528, *Ch. interruptum* SZMC 23937 and *Ch. rectangulare* SZMC 26535 showed no growth at 37 °C, nor *Ch. cochliodes* SZMC 26528 at 45 °C.

Pathogens 2021, 10, x FOR PEER REVIEW

13 of 20





Figure 6. Temperature effect of growth rate: Colony diameters (cm) of the examined *Chaetomium* and *Chaetomium*-like isolates on PDA after four days measured at various temperatures ranging from 4 °C to 45 °C. (A): *C. globosum* strains with a temperature optimum at 25 °C. (B): *C. globosum* strains with a temperature optimum at 30 °C. (C): strains of other *Chaetomium* species. (D): *D. finlandicus* strains.

Figure 6. Temperature effect of growth rate: Colony diameters (cm) of the examined *Chaetomium* and *Chaetomium*-like isolates on PDA after four days measured at various temperatures ranging from 4 °C to 45 °C. (A): *C. globosum* strains with a temperature optimum at 25 °C. (B): *C. globosum* strains with a temperature optimum at 30 °C. (C): strains of other *Chaetomium* species. (D): *D. finlandicus* strains.



As *Chaetomium* and *Chaetomium*-like species are known as cellulolytic fungi, the polysaccharide (cellulose, hemicellulose, and chitin) degrading ability of the isolated strains was determined. The examined *Chaetomium* and *Dichotomopilus* strains showed various enzyme activities (Figure 7). *Ch. globosum* strains produced the highest amounts of extracellular enzymes, ranging from 20.64 to 71.67 U/mL of cellobiohydrolase, 7.57 to 18.99 U/mL of β -glucosidase, 8.54 to 41.57 U/mL of β -xylosidase, and 16.38 to 45.22 U/mL of β -1,4-N-acetyl-glucosaminidase activity. Two strains of *Ch. globosum* (SZMC 27052 and SZMC 26539) showed the highest cellobiohydrolase enzyme activities, which were almost identical in amount 71.13 ± 2.7 and 71.67 ± 2.3 U/mL (respectively). The smallest amount




Figure 6. Temperature effect on growth rate. Colony diameters (mm) of the examined *Chaetomium* and *D. finlandicus* like isolates on PDA after four days measured at various temperatures ranging from 4 to 45°C. (C) The other enzymes were found with a temperature optimum similar to *Ch. globosum* (21.27 and 7.06 U/mL β -glucosidase, 29.01 and 36.08 U/mL β -xylosidase, 18.29 and 26.89 U/mL β -1,4-N-acetyl-glucosaminidase) (Figure 7).




Figure 7. Extracellular enzyme activities of the examined *Chaetomium* and *Chaetomium*-like strains. Error bars show standard deviation of three replicates.

3. Discussion

The dominant species in this study was *Ch. globosum* in both countries in indoor environments, as also determined in several previous studies [5,7,62]. In both countries, the species *Ch. cochliodes* was found to be also common in indoor environments. *Ch. interruptum* was isolated only from Hungary, while *Ch. rectangulare* and *D. finlandicus* only from Finland. Due to the tendency of application of cellulose-based materials (e.g., wallpapers and drywalls) in modern buildings, cellulose-degrading fungi, such as Chaetomiaceae have an increasing relevance. Most indoor strains were isolated from house dust or surface samples, while isolates from air samples were relatively rare. Similar observations were made by Fogle et al. [63] based on the analysis of samplings performed in 794 buildings in Dallas. Although several theories have emerged to explain this phenomenon, further experiments are needed to clarify the dispersal strategy of these fungi indoors.

In a previous study, Salo et al. [5] tested 42 toxin-producing *Chaetomium* isolates from Finland. In addition to the most common *Ch. globosum*, three other species, *Ch. cochliodes*, *Ch. rectangulare*, and a *Chaetomium*-like species were described for the first time from Finnish buildings. In a study by Vornanen-Winqvist et al. [61], unknown indoor *Chaetomium*-like strains were designated as *Dichotomophilus* sp. The molecular results presented here revealed that the *Chaetomium*-like isolate Ch1/tu (SZMC 26529) in Salo et al. [5] and the

Dichotomopilus sp. isolate C5/LM (SZMC 26527) from Vornanen-Winquist et al. [61] belong to the same new, previously undescribed species of the genus *Dichotomopilus*. Strain Ch1/tu was isolated from an inlet air filter and suggested to originate from the outdoor air [5], while strain C5/LM was isolated from an exhaust air filter. This may indicate that C5/LM had a possible indoor source. On the other hand, the fact that this new species was detected in both inlet and outlet air filters may also suggest that the strains were already incorporated into the filter material during production. Contamination of gypsum wall board with *Chaetomium* strains during production has been described by Andersen et al. [64].

The species *D. finlandicus* described in the recent study could be morphologically and molecularly differentiated from related species, the results of the phylogenetic analyses of the combined dataset of ITS, *tef1α*, *rpb2*, and *tub2* (Figure 4) was 100% bootstrap support. In addition, the phylogenetically closest relative species *D. funicola*, *D. pseudofunicola*, *D. subfunicola*, *D. variostiolatus*, and *D. indicus* are morphologically different from the strain we studied. Based on the morphological properties of these species studied by Wang et al. [7], ascomata, terminal hairs, and the asci were different while the shape and the size of ascospores were similar to *D. finlandicus*, which we describe here as a new species.

The enzymatic activity of the *Chaetomium* and *Chaetomium*-like strains proved to be diverse, and no correlation was found with either the isolation site or the growing substrate. These results are consistent with the findings of Abdel Azeem et al. [22], that enzyme production is isolate-dependent. The authors concluded that enzyme production has no detectable association with ecology, however, although this may be true in the case of plant host specificity, we suggest the ability to produce cellulolytic enzymes as a clear ecological advantage in the case of fungal growth on cellulose-based building materials.

In the rapid screening assays described by Salo et al. [5] and Vornanen-Winquist et al. [65], *Dichotomopilus* strains gave weaker responses than the *Ch. globosum*, *Ch. cochliodes* and *Ch. rectangulare* strains. However, strain Ch1/tu (SZMC 26529), which was designated here as the type strain of the newly described species *D. finlandicus*, inhibited boar sperm motility after 3 d of exposure, indicating that the strain produced a bioactive agent possibly affecting mitochondrial functions, or ion homeostasis [66]. Purification and identification of this substance and characterization of its biological activities will be the subject of further research.

4. Materials and Methods

4.1. Sample Collection and Isolation

Chaetomium and *Chaetomium*-like strains were collected and isolated from schools, flats, and offices in Finland as described previously by Salo et al. [5], as well as from houses, flats, and offices in Hungary (Table 1). To collect fungi from walls, visible colonies, or wet surfaces detected by moisture meter (Greisinger GMI 15) were sampled with sterile swabs. House dust samples were also collected with swabs. Samples were spread directly onto malt extract agar (MEA) supplemented with 2% chloramphenicol, Dichloran - Rose Bengal Agar, or Casitone Agar on site. To collect airborne fungi, air samples of 100 L were collected at 150 cm a.g.l. with 400-hole one-stage Andersen samplers [67] (MAS 100, EMD Millipore, Merck, Darmstadt, Germany; SAS IAQ, International PBI SpA, Milan, Italy; Samp l'Air MK2, AES Chemunex, Bruz, France), at a flow of 100 L/min onto MEA. Between samplings, the devices were sterilized with ethanol (abs.). Incubation of the samples was performed for 5 to 7 days at room temperature. The isolated pure cultures were deposited in the Szeged Microbiology Collection (SZMC, <http://szmc.hu>), Szeged, Hungary.

4.2. Morphological Characterization

The morphology and colony characteristics of the isolates were examined on three different media: 2% (*w/v*) MEA (VWR, Debrecen, Hungary), 3% (*w/v*) oatmeal agar (OA, Merck, Darmstadt, Germany), and potato dextrose agar (PDA, VWR, Debrecen, Hungary), and incubated for seven days in the dark at 25 °C [2,7]. Microscopic studies were performed using light—(Zeiss Primostar, Carl Zeiss, Suzhou, China), stereo—(Zeiss Stemi 305, Carl

Zeiss, Suzhou, China), and scanning electron microscopes. SEM samples were prepared by stabilization in 0.1 M phosphate buffer (pH 7.3) containing 2.5% glutaraldehyde (12 h, 4 °C). The samples were then dehydrated with ethanol-water, gradually increasing the volume ratio of ethanol (50% *v/v*, 60% *v/v*, 70% *v/v*, 80% *v/v*, 90% *v/v*, 95% *v/v*, 100% *v/v*). Finally, the samples were dried (3 h, 30 °C) and coated with gold for microscopic examination. Electron microscopy images were taken with a 10 kV accelerating voltage Hitachi S-4700 Type II FE-SEM microscope, observing secondary electrons with magnitudes of 150×, 250×, 600×, and 2000× [68].

4.3. DNA Extraction, Identification, and Phylogenetic Analysis

Pure cultures of fungi were grown in 2% (*w/v*) MEA for 7 days at room temperature. Fungal genomic DNA was then extracted using the E.Z.N.A.®Fungal DNA Mini Kit (Omega Biotek, Norcross, GA, USA). The extracted genomic DNA was amplified by PCR with the primers listed in Table 2. The PCR mixture (20 µL) contained 2 µL 10× DreamTaq Buffer with 20 mM MgCl₂, 2 µL of 2 mM dNTP mix, 4 µL of each primer (100 µM), 7 µL bidistilled water, 0.1 µL of 5 U/µL DreamTaq DNA Polymerase (Thermo Fischer Scientific, Vilnius, Lithuania) and 1 µL genomic DNA. Amplifications were performed in a Doppio Gradient 2 × 48-well thermal cycler (VWR International, Debrecen, Hungary) according to the amplification cycles shown in Table 2. PCR products were purified using NucleoSpin™ Gel and PCR Clean-up Kit (Macherey-Nagel, Düren, Germany). Sequencing was performed on the sequencing platform of Eurofins Genomics (<http://www.eurofinsgenomics.com>, accessed on 2 September 2021). The resulting sequences were submitted to the GenBank Nucleotide database (ncbi.nlm.nih.gov) under the accession numbers listed in Table 1. In addition to the sequences generated in this study, sequences of reference strains were obtained from the GenBank Nucleotide database (Table 1).

Sequences of the two *Dichotomopilus* isolates were aligned with publicly available sequences of 12 and 11 previously described *Dichotomopilus* and *Chaetomium* species, respectively. Phylogenetic analyses were conducted using four loci (ITS, *tef1α*, *rpb2*, and *tub2*).

Sequences were aligned with Prank v170427 [69]. Alignments of the four loci were concatenated and partitioned. *Tef1α* and *rpb2* sequences were defined as two single partitions, while the *tub2* dataset was partitioned to exons and introns. The ITS dataset was divided to rDNA and ITS1-ITS2 regions. Alignments of *tub2* and ITS datasets contained relative high number of indels, therefore gaps were coded as absence/presence characters by 2matrix v1.0 [70] using the simple indel coding algorithm [71]. The two indel matrices were concatenated and added as a single partition to the dataset. Best fitting model for the phylogenetic inference was selected by using ModelTest-NG v0.1.4 [72], based on the Bayesian information criterion [73], with discrete gamma rate categories. Best fit models for each partition are shown in Table 3. Maximum likelihood analysis was performed using RAXML-NG v0.9.0 [74]. Statistical support of the best ML tree was obtained with 1000 bootstrap replicates.

Table 2. List of the amplified genes, used primers, and PCR conditions

Gene	Primer	PCR Condition
<i>tefla</i>	EF1-728F: CATCGAGAAGTTCGAGAAGG TTEF1-LLErev: AACTTGCAGGCAATGTGG	94 °C 5 min
		94 °C 30 s
		57 °C 30 s
		72 °C 90 s
		40 cycles
		72 °C 7 min
ITS	ITS1: TCCGTAGCTCAACCTCCGG ITS4: TCCCTCCGCTTATGATATGC	94 °C 2 min
		94 °C 30 s
		48 °C 40 s
		72 °C 40 s
		35 cycles
		72 °C 2 min
<i>tub2</i>	BT2a: GGTAAACCAATCGGTGCTGGTTC BT2b: ACCCTCAGTCTAGTCACCCCTGGC	94 °C 2 min
		94 °C 30 s
		55 °C 30 s
		72 °C 45 s
		35 cycles
		72 °C 7 min
<i>rpb2</i>	RPB2-5F_Eur: GAYGAYCGCKGAYCAYTTCGG RPB2-7CR_Eur: CCCATRGCTGCTGTTGCCAT	94 °C 5 min
		94 °C 45 s
		60 °C 45 s
		72 °C 2 min
		94 °C 45 s
		58 °C 45 s
		72 °C 2 min
		94 °C 45 s
		5 cycles
		54 °C 45 s
		72 °C 2 min
		30 cycles
		72 °C 7 min

Table 3. Best-fit models for each partition proposed by ModelTest-NG based on Bayesian information criterion. Sequences were aligned with Prank v170427 [69]. Alignments of the four loci were concatenated and partitioned. *Tefla* and *rpb2* sequences were defined as two single partitions, while the *tub2* dataset was partitioned to exons and introns. The ITS dataset was divided to rDNA and ITS1-ITS2 regions. Alignments of *tub2* and ITS datasets contain relative high number of indels, therefore gaps were coded as absence/presence characters by 2matrix v1.0 [70] using the simple indel coding algorithm [71]. The two indel matrices were concatenated and added as a single partition to the dataset. Best fitting model for phylogenetic inference was selected by using ModelTest-NG v0.1.4 [72], based on the Bayesian information criterion [73], with discrete gamma rate categories. Best fit models for each partition are shown in Table 3. Maximum likelihood analysis was performed using RAxML-NG v0.9.0 [74]. Statistical support of the best ML tree was obtained with 100 bootstrap replicates.

4.4. Enzyme Production

For enzyme activity measurements, fungal strains were grown in cellulose-containing broth (20 g/L mannitol, 10 g/L KH_2PO_4 , 5 g/L NaNO_3 , 2 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 20 g/L cellulose powder) for seven days at 25 °C with shaking (150 rpm) on a MaxQ 8000 Incubated Stackable Shaker (Thermo Fisher Scientific, Waltham, USA). Enzyme activity measurements were performed from culture supernatants with the chromogenic substrates 4-nitrophenyl- β -D-glucopyranoside (β -glucosidase), 4-nitrophenyl- β -D-cellobiose (cellobiohydrolase), 4-nitrophenyl- β -D-xylopyranoside (β -xylosidase), 4-nitrophenyl-N-acetyl- β -D-glucosamine (β -1,4-N-acetyl-glucosaminidase) (Sigma Aldrich, St. Louis, MO, USA). 100 μL of the culture supernatants were pipetted into the wells of a microtiter plate and 100 μL of 3 mM 4-nitrophenyl substrate was added. The mixtures were incubated at

37 °C for 1 hour; thereafter the reactions were stopped by the addition of 100 µL Na₂CO₃ solution (0.1 M). The released 4-nitrophenol was measured on a Spectrostar Nano microtiter plate reader (BMG Labtech, Ortenberg, Germany) at 405 nm. The optical density values obtained were converted to units: 1 unit of enzyme activity was defined as the amount of enzyme required to release 1 µmol of p-nitrophenol per min under the determined reaction conditions. Calibration curve was prepared using standard solutions of 4-nitrophenol of known concentration.

4.5. Temperature Profiling

Optimal growth temperature ranges were determined for all *Chaetomium* and *Chaetomium*-like isolates. PDA plates were inoculated with 7 mm agar plates taken from the edge of seven-day-old colonies. The plates were incubated at 4, 15, 21, 25, 30, 35, 37, 40, and 45 °C, with six replicates each. Colony diameters were measured after four days.

Author Contributions: Conceptualization, O.K. and L.K.; Methodology, O.K., T.B., M.A.A., J.M.S., R.M., H.S., A.K. and Z.K.; Software, S.K. and A.S.; Validation, J.M.S., S.K., A.S., Z.K., C.V. and D.M.; Formal analysis, O.K., S.K. and A.S.; Investigation, O.K., T.B., M.A.A., R.M. and A.K.; Resources, H.S., Z.K., C.V. and D.M.; Data curation, O.K., M.A.A., D.M., S.K. and L.K.; Writing—original draft, O.K., S.K., M.A.A., A.S., A.K., D.M. and L.K.; Writing—review and editing, O.K., M.A.A., J.M.S., H.S., R.M., C.V., Z.K., D.M. and L.K.; Visualization, O.K., S.K. and A.K.; Supervision, O.K., Z.K. and L.K.; Project administration, H.S., D.M. and L.K.; Funding acquisition, O.K., M.A.A., H.S. and R.M. All authors have read and agreed to the published version of the manuscript.

Funding: O.K. was supported by the ÚNKP-20-3—New National Excellence Program of the Ministry for Innovation and Technology from the source of the National Research, Development and Innovation Fund. M.A.A., H.S. and R.M. are grateful to the Academy of Finland (CleanSchool-project, grant no. 330150) for financial support.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Nucleotide sequences were deposited in the GenBank Nucleotide Database (<https://www.ncbi.nlm.nih.gov>), accession numbers are provided in Table 1.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Rodríguez, K.; Stchigel, A.; Guarro, J. Three new species of *Chaetomium* from soil. *Mycologia* **2002**, *94*, 116–126. [CrossRef] [PubMed]
- Wang, X.W.; Wang, X.L.; Liu, F.J.; Zhao, X.M.; Li, J.; Cai, L. Phylogenetic assessment of *Chaetomium indicum* and allied species, with the introduction of three new species and epitypification of *C. funicola* and *C. indicum*. *Mycol. Prog.* **2014**, *13*, 719–732. [CrossRef]
- Došen, I.; Nielsen, K.F.; Clausen, G.; Andersen, B. Potentially harmful secondary metabolites produced by indoor *Chaetomium* species on artificially and naturally contaminated building materials. *Indoor Air* **2017**, *27*, 34–46. [CrossRef] [PubMed]
- Andersen, B.; Frisvad, J.C.; Søndergaard, I.; Rasmussen, I.S.; Larsen, L.S. Associations between fungal species and water-damaged building materials. *Appl. Environ. Microbiol.* **2011**, *77*, 4180–4188. [CrossRef]
- Salo, J.M.; Kedves, O.; Mikkola, R.; Kredics, L.; Andersson, M.A.; Kurnitski, J.; Salonen, H. Detection of *Chaetomium globosum*, *Ch. cochliodes* and *Ch. rectangulare* during the diversity tracking of mycotoxin-producing *Chaetomium*-like isolates obtained in buildings in Finland. *Toxins* **2020**, *12*, 443. [CrossRef]
- Castagnoli, E.; Mikkola, R.; Vornanen-Winquist, C.; Andersson, M.; Kredics, L.; Marik, T.; Kurnitski, J.; Salonen, H. Indoor *Chaetomium*-like isolates; resistance to chemicals, fluorescence and mycotoxin production. *Sisäilmastoseminaari* **2017**, *35*, 227–232.
- Wang, X.W.; Houbaken, J.; Groenewald, J.Z.; Meijer, M.; Andersen, B.; Nielsen, K.F.; Crous, P.W.; Samson, R.A. Diversity and taxonomy of *Chaetomium* and *Chaetomium*-like fungi from indoor environments. *Stud. Mycol.* **2016**, *84*, 145–224. [CrossRef]
- Stiller, M.J.; Rosenthal, S.; Summerbell, R.C.; Pollack, J.; Chan, A. Onychomycosis of the toenails caused by *Chaetomium globosum*. *J. Am. Acad. Dermatol.* **1992**, *26*, 775–776. [CrossRef]
- Aspiroz, C.; Gené, J.; Rezusta, A.; Charlez, L.; Summerbell, R.C. First Spanish case of onychomycosis caused by *Chaetomium globosum*. *Med. Mycol.* **2007**, *45*, 279–282. [CrossRef] [PubMed]
- Hwang, S.M.; Suh, M.K.; Ha, G.Y. Onychomycosis due to nondermatophytic molds. *Ann. Dermatol.* **2012**, *24*, 175–180. [CrossRef] [PubMed]

11. Kim, D.M.; Lee, M.H.; Suh, M.K.; Ha, G.Y.; Kim, H.; Choi, J.S. Onychomycosis caused by *Chaetomium globosum*. *Ann. Dermatol.* **2013**, *25*, 232–236. [\[CrossRef\]](#)
12. Shi, D.; Lu, G.; Mei, H.; de Hoog, G.S.; Zheng, H.; Liang, G.; Shen, Y.; Li, T.; Liu, W. Onychomycosis due to *Chaetomium globosum* with yellowish black discoloration and periungual inflammation. *Med. Mycol. Case Rep.* **2016**, *13*, 12–16. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Hubka, V.; Mencl, K.; Skorepova, M.; Lyskova, P.; Zalabska, E. Phaeohyphomycosis and onychomycosis due to *Chaetomium* spp., including the first report of *Chaetomium brasiliense* infection. *Med. Mycol.* **2011**, *49*, 724–733. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Abbott, S.P.; Sigler, L.; McAleer, R.; McGough, D.A.; Rinaldi, M.G.; Mizell, G. Fatal cerebral mycoses caused by the ascomycete *Chaetomium strumarium*. *J. Clin. Microbiol.* **1995**, *33*, 2692–2698. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Thomas, C.; Mileusnic, D.; Carey, R.B.; Kampert, M.; Anderson, D. Fatal *Chaetomium* cerebritis in a bone marrow transplant patient. *Hum. Pathol.* **1999**, 874–879. [\[CrossRef\]](#)
16. Barron, M.A.; Sutton, D.A.; Veve, R.; Guarro, J.; Rinaldi, M.; Thompson, E.; Cagnoni, P.J.; Moultny, K.; Madinger, N.E. Invasive mycotic infections caused by *Chaetomium perlucidum*, a new agent of cerebral phaeohyphomycosis. *J. Clin. Microbiol.* **2003**, *41*, 5302–5307. [\[CrossRef\]](#)
17. Gao, K.; Liu, X.; Kang, Z.; Mendgen, K. Mycoparasitism of *Rhizoctonia solani* by endophytic *Chaetomium spirale* ND35: Ultrastructure and cytochemistry of the interaction. *J. Phytopathol.* **2005**, *153*, 280–290. [\[CrossRef\]](#)
18. Ding, G.; Song, Y.C.; Chen, J.R.; Xu, C.; Ge, H.M.; Wang, X.T.; Tan, R.X. Chaetoglobosin U, a cytochalasan alkaloid from endophytic *Chaetomium globosum* IFB-E019. *J. Nat. Prod.* **2006**, *69*, 302–304. [\[CrossRef\]](#) [\[PubMed\]](#)
19. Sharma, R.; Kulkarni, G.; Sonawane, M.S.; Shouche, Y.S. A new endophytic species of *Chaetomium* from *Jatropha podagrica*. *Mycotaxon* **2013**, *124*, 117–126. [\[CrossRef\]](#)
20. Kamat, S.; Kumari, M.; Sajna, K.V.; Jayabaskaran, C. Endophytic fungus, *Chaetomium globosum*, associated with marine green alga, a new source of Chrysin. *Sci. Rep.* **2020**, *10*, 18726. [\[CrossRef\]](#)
21. Lakshmikanth, Kamal; Mathur, S.N. Cellulolytic activities of *Chaetomium globosum* on different cellulosic substrates. *World J. Microbiol. Biotechnol.* **1990**, *6*, 23–26. [\[CrossRef\]](#)
22. Abdel-Azeem, A.M.; Gherbawy, Y.A.; Sabry, A.M. Enzyme profiles and genotyping of *Chaetomium globosum* isolates from various substrates. *Plant Biosyst.* **2016**, *150*, 420–428. [\[CrossRef\]](#)
23. Wanmolee, W.; Sornlake, W.; Rattanaphan, N.; Suwannarangsee, S.; Laosiripojana, N.; Champreda, V. Biochemical characterization and synergism of cellulolytic enzyme system from *Chaetomium globosum* on rice straw saccharification. *BMC Biotechnol.* **2016**, *16*, 82. [\[CrossRef\]](#)
24. Zhou, Q.; Jia, J.; Ji, P.; Han, C. A novel application potential of GH6 cellobiohydrolase ctcl6 from thermophilic *Chaetomium thermophilum* for gene cloning, heterologous expression and biological characterization. *Int. J. Agric. Biol.* **2017**, *19*, 355–362. [\[CrossRef\]](#)
25. Li, H.; Liao, Z.B.; Tang, D.; Han, W.B.; Zhang, Q.; Gao, J.M. Polyketides from two *Chaetomium* species and their biological functions. *J. Antibiot.* **2018**, *71*, 677–681. [\[CrossRef\]](#)
26. Wang, Z.; Jia, S.; Cui, J.; Qu, J.; Yue, Y.; Sun, Q.; Zhang, H. Antioxidant activity of a polysaccharide produced by *Chaetomium globosum* CGMCC 6882. *Int. J. Biol. Macromol.* **2019**, *141*, 955–960. [\[CrossRef\]](#)
27. Serena, C.; Ortoneda, M.; Capilla, J.; Pastor, F.J.; Sutton, D.A.; Rinaldi, M.G.; Guarro, J. In vitro activities of new antifungal agents against *Chaetomium* spp. and inoculum standardization. *Antimicrob. Agents Chemother.* **2003**, *47*, 3161–3164. [\[CrossRef\]](#) [\[PubMed\]](#)
28. Jiao, W.; Feng, Y.; Blunt, J.W.; Cole, A.L.J.; Munro, M.H.G. Chaetoglobosins Q, R, and T, three further new metabolites from *Chaetomium globosum*. *J. Nat. Prod.* **2004**, *67*, 1722–1725. [\[CrossRef\]](#)
29. Park, J.H.; Gyung, J.C.; Kyoung, S.J.; He, K.L.; Heung, T.K.; Kwang, Y.C.; Kim, J.C. Antifungal activity against plant pathogenic fungi of chaetoviridins isolated from *Chaetomium globosum*. *FEMS Microbiol. Lett.* **2005**, *252*, 309–313. [\[CrossRef\]](#)
30. Huang, S.; Chen, H.; Li, W.; Zhu, X.; Ding, W.; Li, C. Bioactive chaetoglobosins from the mangrove endophytic fungus *Penicillium chrysogenum*. *Mar. Drugs* **2016**, *14*, 172. [\[CrossRef\]](#)
31. Yan, W.; Cao, L.L.; Zhang, Y.Y.; Zhao, R.; Zhao, S.S.; Khan, B.; Ye, Y.H. New metabolites from endophytic fungus *Chaetomium globosum* CDW7. *Molecules* **2018**, *23*, 2873. [\[CrossRef\]](#)
32. Chovanová, K.; Zámocký, M. Detection of the antibacterial effect of *Chaetomium cochliodes* Palliser CCM F-232 based on agar plugs and unprocessed fungal substances from cultivation media. *Biologia* **2016**, *71*, 1204–1211. [\[CrossRef\]](#)
33. Dissanayake, R.K.; Ratnaweera, P.B.; Williams, D.E.; Wijayarathne, C.D.; Wijesundera, R.L.C.; Andersen, R.J.; de Silva, E.D. Antimicrobial activities of endophytic fungi of the Sri Lankan aquatic plant *Nymphaea nouchali* and chaetoglobosin A and C, produced by the endophytic fungus *Chaetomium globosum*. *Mycology* **2016**, *7*, 1–8. [\[CrossRef\]](#)
34. Gao, W.; He, Y.; Li, F.; Chai, C.; Zhang, J.; Guo, J.; Chen, C.; Wang, J.; Zhu, H.; Hu, Z.; et al. Antibacterial activity against drug-resistant microbial pathogens of cytochalasan alkaloids from the arthropod-associated fungus *Chaetomium globosum* TW1-1. *Bioorg. Chem.* **2019**, *83*, 98–104. [\[CrossRef\]](#)
35. Wang, Z.; Xue, R.; Cui, J.; Wang, J.; Fan, W.; Zhang, H.; Zhan, X. Antibacterial activity of a polysaccharide produced from *Chaetomium globosum* CGMCC 6882. *Int. J. Biol. Macromol.* **2019**, *125*, 376–382. [\[CrossRef\]](#) [\[PubMed\]](#)
36. Wani, N.; Khanday, W.; Tirumale, S. Evaluation of In vitro antioxidant potential of active metabolite constituents of different extracts of *Chaetomium cupreum*-SS02 by spectrophotometric method. *Matrix Sci. Pharma* **2020**, *4*, 50. [\[CrossRef\]](#)
37. Attia, E.; Dashora, K.; Abdel-azeem, A. A potential antimicrobial, extracellular enzymes, and antioxidants resource: Endophytic fungi associated with medicinal plants. *Int. J. Biosci.* **2020**, *17*, 119–132. [\[CrossRef\]](#)

38. Khan, B.; Yan, W.; Wei, S.; Wang, Z.; Zhao, S.; Cao, L.; Rajput, N.A.; Ye, Y. Nematicidal metabolites from endophytic fungus *Chaetomium globosum* YSC5. *FEMS Microbiol. Lett.* **2019**, *366*, fnz169. [\[CrossRef\]](#)
39. Hu, Y.; Zhang, W.; Zhang, P.; Ruan, W.; Zhu, X. Nematicidal activity of chaetoglobosin A produced by *Chaetomium globosum* NK102 against *Meloidogyne incognita*. *J. Agric. Food Chem.* **2013**, *61*, 41–46. [\[CrossRef\]](#)
40. Chen, C.; Tong, Q.; Zhu, H.; Tan, D.; Zhang, J.; Xue, Y.; Yao, G.; Luo, Z.; Wang, J.; Wang, Y.; et al. Nine new cytochalasan alkaloids from *Chaetomium globosum* TW1-1 (Ascomycota, Sordariales). *Sci. Rep.* **2016**, *6*, 18711. [\[CrossRef\]](#)
41. Ruan, B.H.; Yu, Z.F.; Yang, X.Q.; Yang, Y.B.; Hu, M.; Zhang, Z.X.; Zhou, Q.Y.; Zhou, H.; Ding, Z.T. New bioactive compounds from aquatic endophyte *Chaetomium globosum*. *Nat. Prod. Res.* **2018**, *32*, 1050–1055. [\[CrossRef\]](#)
42. Wang, Z.; Chen, P.; Tao, N.; Zhang, H.; Li, R.; Zhan, X.; Wang, F.; Shen, Y. Anticancer activity of polysaccharides produced from glycerol and crude glycerol by an endophytic fungus *Chaetomium globosum* CGMCC 6882 on human lung cancer A549 cells. *Biomolecules* **2018**, *8*, 171. [\[CrossRef\]](#) [\[PubMed\]](#)
43. Senthil Kumar, V.; Kumaresan, S.; Tamizh, M.M.; Hairul Islam, M.I.; Thirugnanasambantham, K. Anticancer potential of NF- κ B targeting apoptotic molecule “flavipin” isolated from endophytic *Chaetomium globosum*. *Phytomedicine* **2019**, *61*, 152830. [\[CrossRef\]](#)
44. Wani, N.; Khanday, W.; Tirumale, S. Biosynthesis of iron oxide nanoparticles using ethyl acetate extract of *Chaetomium cupreum* and their anticancer activity. *Matrix Sci. Pharma* **2020**, *4*, 23. [\[CrossRef\]](#)
45. Wijeratne, E.M.K.; Turbyville, T.J.; Fritz, A.; Whitesell, L.; Gunatilaka, A.A.L. A new dihydroxanthone from a plant-associated strain of the fungus *Chaetomium globosum* demonstrates anticancer activity. *Bioorg. Med. Chem.* **2006**, *14*, 7917–7923. [\[CrossRef\]](#)
46. Ming Ge, H.; Yun Zhang, W.; Ding, G.; Saparpakorn, P.; Chun Song, Y.; Hannongbua, S.; Xiang Tan, R. Chaetoglobins A and B, two unusual alkaloids from endophytic *Chaetomium globosum* culture. *Chem. Commun.* **2008**, *45*, 5978–5980. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Thohinung, S.; Kanokmedhakul, S.; Kanokmedhakul, K.; Kukongviriyapan, V.; Tusksorn, O.; Soyong, K. Cytotoxic 10-(indol-3-yl)-[13]cytochalasans from the fungus *Chaetomium elatum* ChE01. *Arch. Pharm. Res.* **2010**, *33*, 1135–1141. [\[CrossRef\]](#)
48. Li, H.; Xiao, J.; Gao, Y.Q.; Tang, J.J.; Zhang, A.L.; Gao, J.M. Chaetoglobosins from *Chaetomium globosum*, an endophytic fungus in ginkgo biloba, and their phytotoxic and cytotoxic activities. *J. Agric. Food Chem.* **2014**, *62*, 3734–3741. [\[CrossRef\]](#)
49. Zheng, Q.C.; Kong, M.Z.; Zhao, Q.; Chen, G.D.; Tian, H.Y.; Li, X.X.; Guo, L.D.; Li, J.; Zheng, Y.Z.; Gao, H. Chaetoglobosin Y, a new cytochalasan from *Chaetomium globosum*. *Fitoterapia* **2014**, *93*, 126–131. [\[CrossRef\]](#)
50. Hamed Eman, M.H. Anticancer compounds from *Chaetomium globosum*. *Biochem. Anal. Biochem.* **2015**, *4*, 174. [\[CrossRef\]](#)
51. Chen, C.; Shao, Y.; Tao, Y.; Wen, H. Optimization of dynamic microwave-assisted extraction of *Armillaria* polysaccharides using RSM, and their biological activity. *LWT-Food Sci. Technol.* **2015**, *64*, 1263–1269. [\[CrossRef\]](#)
52. Wang, Z.; Liu, X.; Bao, Y.; Wang, X.; Zhai, J.; Zhan, X.; Zhang, H. Characterization and anti-inflammation of a polysaccharide produced by *Chaetomium globosum* CGMCC 6882 on LPS-induced RAW 264.7 cells. *Carbohydr. Polym.* **2021**, *251*, 117129. [\[CrossRef\]](#) [\[PubMed\]](#)
53. Madbouly, A.K.; Abdel-Aziz, M.S.; Abdel-Wahhab, M.A. Biosynthesis of nanosilver using *Chaetomium globosum* and its application to control Fusarium wilt of tomato in the greenhouse. *IET Nanobiotechnol.* **2017**, *11*, 702–708. [\[CrossRef\]](#)
54. El-Mohamedy, R.S.R.; El-Gamal, N.G.; El-Shamy, A.R.; Atalla, S.M.M. Biosynthesis of zinc nanoparticles and its effect on enzymes production by *Bacillus subtilis* and *Pseudomonas fluorescens* using different agricultural wastes. *Int. J. Agric. Technol.* **2018**, *14*, 833–844.
55. Wang, X.W.; Zheng, R.Y. *Chaetomium acropullum* sp. nov. (Chaetomiaceae, Ascomycota), a new psychrotolerant mesophilic species from China. *Nova Hedwigia* **2005**, *80*, 413–417. [\[CrossRef\]](#)
56. Abdel-Azeem, A.M.; Abu-Elsaoud, A.M.; Abo Nahas, H.H.; Abdel-Azeem, M.A.; Balbool, B.A.; Mousa, M.K.; Ali, N.H.; Darwish, A.M.G. Biodiversity and industrial applications of genus *Chaetomium*. In *Industrially Important Fungi for Sustainable Development*; Abdel-Azeem, A.M., Yadav, A.N., Yadav, N., Usmani, Z., Eds.; Springer: Cham, Switzerland, 2021; Volume 1, pp. 147–206.
57. Gatumbi, R.W.; Kung'u, J.N. Notes on species of the genus *Chaetomium* from Kenya. *East. Afr. Agric. For. J.* **1994**, *60*, 79–89. [\[CrossRef\]](#)
58. Udagawa, S.I. A new species of *Chaetomium* from house dust. *Mycoscience* **1997**, *38*, 399–402. [\[CrossRef\]](#)
59. Andersen, B.; Nissen, A.T. Evaluation of media for detection of *Stachybotrys* and *Chaetomium* species associated with water-damaged buildings. *Int. Biodeterior. Biodegrad.* **2000**, *46*, 111–116. [\[CrossRef\]](#)
60. Piecková, E. In vitro toxicity of indoor *Chaetomium kunze* ex fr. *Ann. Agric. Environ. Med.* **2003**, *10*, 9–14.
61. Vornanen-Winqvist, C.; Järvi, K.; Andersson, M.A.; Duchaine, C.; Létourneau, V.; Kedves, O.; Kredics, L.; Mikkola, R.; Kurnitski, J.; Salonen, H. Exposure to indoor air contaminants in school buildings with and without reported indoor air quality problems. *Environ. Int.* **2020**, *141*, 105781. [\[CrossRef\]](#) [\[PubMed\]](#)
62. McMullin, D.R.; Sumarah, M.W.; Miller, J.D. Chaetoglobosins and azaphilones produced by Canadian strains of *Chaetomium globosum* isolated from the indoor environment. *Mycotoxin Res.* **2013**, *29*, 47–54. [\[CrossRef\]](#) [\[PubMed\]](#)
63. Fogle, M.R.; Douglas, D.R.; Jumper, C.A.; Straus, D.C. Growth and mycotoxin production by *Chaetomium globosum*. *Mycopathologia* **2007**, *164*, 49–56. [\[CrossRef\]](#)
64. Andersen, B.; Dosen, I.; Lewinska, A.M.; Nielsen, K.F. Pre-contamination of new gypsum wallboard with potentially harmful fungal species. *Indoor Air* **2017**, *27*, 6–12. [\[CrossRef\]](#)

-
65. Vornanen-Winqvist, C.; Järvi, K.; Toomla, S.; Ahmed, K.; Andersson, M.A.; Mikkola, R.; Marik, T.; Kredics, L.; Salonen, H.; Kurnitski, J. Ventilation positive pressure intervention effect on indoor air quality in a school building with moisture problems. *Int. J. Environ. Res. Public Health* **2018**, *15*, 230. [[CrossRef](#)]
 66. Castagnoli, E.; Salo, J.; Toivonen, M.S.; Marik, T.; Mikkola, R.; Kredics, L.; Vicente-Carrillo, A.; Nagy, S.; Andersson, M.T.; Andersson, M.A.; et al. An evaluation of boar spermatozoa as a biosensor for the detection of sublethal and lethal toxicity. *Toxins* **2018**, *10*, 463. [[CrossRef](#)]
 67. Andersen, A.A. New sampler for the collection, sizing, and enumeration of viable airborne particles. *J. Bacteriol.* **1958**, *76*, 471–484. [[CrossRef](#)]
 68. Kedves, A.; Rónavári, A.; Kónya, Z. Long-term effect of graphene oxide on the aerobic granular sludge wastewater treatment process. *J. Environ. Chem. Eng.* **2021**, *9*, 104853. [[CrossRef](#)]
 69. Löytynoja, A. Phylogeny-aware alignment with PRANK. *Meth. Mol. Biol.* **2014**, *1079*, 155–170. [[CrossRef](#)]
 70. Salinas, N.R.; Little, D.P. 2matrix: A utility for indel coding and phylogenetic matrix concatenation. *Appl. Plant Sci.* **2014**, *2*, 1300083. [[CrossRef](#)] [[PubMed](#)]
 71. Simmons, M.P.; Ochoterena, H. Gaps as characters in sequence-based phylogenetic analysis. *Syst. Biol.* **2000**, *49*, 369–381. [[CrossRef](#)] [[PubMed](#)]
 72. Darriba, D.; Posada, D.; Kozlov, A.M.; Stamatakis, A.; Morel, B.; Flouri, T. ModelTest-NG: A new and scalable tool for the selection of DNA and protein evolutionary models. *Mol. Biol. Evol.* **2020**, *37*, 291–294. [[CrossRef](#)] [[PubMed](#)]
 73. Schwarz, G. Estimating the dimension of a model. *Ann. Stat.* **1978**, *6*, 461–464. [[CrossRef](#)]
 74. Kozlov, A.M.; Darriba, D.; Flouri, T.; Morel, B.; Stamatakis, A. RAxML-NG: A fast, scalable, and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **2019**, *35*, 4453–4455. [[CrossRef](#)] [[PubMed](#)]