
This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Agarwal, Ashok; Finelli, Renata; Durairajanayagam, Damayanthi; Leisegang, Kristian; Henkel, Ralf; Salvio, Gianmaria; Aghamajidi, Azin; Sengupta, Pallav; Crisostomo, Luis; Tsioulou, Petroula A.; Roychoudhury, Shubhadeep; Finocchi, Federica; Darbandi, Mahsa; Mottola, Filomena; Darbandi, Sara; Iovine, Concetta; Santonastaso, Marianna; Zaker, Himasadat; Kesari, Kavindra Kumar; Nomanzadeh, Amir; Gugnani, Nivita; Rambhatla, Amarnath; Duran, Mesut Berkan; Ceyhan, Erman; Kandil, Hussein; Arafa, Mohamed; Saleh, Ramadan; Shah, Rupin; Boitrelle, Florence

Comprehensive Analysis of Global Research on Human Varicocele: A Scientometric Approach

Published in:

World Journal of Men's Health

DOI:

[10.5534/wjmh.210202](https://doi.org/10.5534/wjmh.210202)

Published: 01/10/2022

Document Version













Publisher's PDF, also known as Version of record

Please cite the original version:

Agarwal, A., Finelli, R., Durairajanayagam, D., Leisegang, K., Henkel, R., Salvio, G., Aghamajidi, A., Sengupta, P., Crisostomo, L., Tsioulou, P. A., Roychoudhury, S., Finocchi, F., Darbandi, M., Mottola, F., Darbandi, S., Iovine, C., Santonastaso, M., Zaker, H., Kesari, K. K., ... Boitrelle, F. (2022). Comprehensive Analysis of Global Research on Human Varicocele: A Scientometric Approach. *World Journal of Men's Health*, 40(4), 636-652. <https://doi.org/10.5534/wjmh.210202>



Comprehensive Analysis of Global Research on Human Varicocele: A Scientometric Approach

Ashok Agarwal¹, Renata Finelli¹, Damayanthi Durairajanayagam², Kristian Leisegang³, Ralf Henkel^{1,4,5,6}, Gianmaria Salvio⁷, Azin Aghamajidi⁸, Pallav Sengupta⁹, Luís Crisóstomo^{10,11}, Petroula A. Tsioulou¹², Shubhadeep Roychoudhury¹³, Federica Finocchi⁷, Mahsa Darbandi¹⁴, Filomena Mottola¹⁵, Sara Darbandi¹⁴, Concetta Iovine^{15,16}, Marianna Santonastaso¹⁷, Himasadat Zaker¹⁸, Kavindra Kumar Kesari¹⁹, Amir Nomanzadeh¹⁸, Nivita Gugnani²⁰, Amarnath Rambhatla²¹, Mesut Berkan Duran²², Erman Ceyhan²³, Hussein Kandil²⁴, Mohamed Arafa^{1,25,26}, Ramadan Saleh^{27,28}, Rupin Shah²⁹, Florence Boitrelle^{30,31}

¹American Center for Reproductive Medicine, Cleveland Clinic, Cleveland, OH, USA, ²Department of Physiology, Faculty of Medicine, Universiti Teknologi Mara, Sungai Buloh Campus, Selangor, Malaysia, ³Department of Physiology, School of Natural Medicine, Faculty of Community and Health Sciences, University of the Western Cape, Bellville, South Africa, ⁴LogixX Pharma Ltd., Theale, Berkshire, UK, ⁵Department of Metabolism, Digestion and Reproduction, Imperial College London, London, UK, ⁶Department of Medical Bioscience, University of the Western Cape, Bellville, South Africa, ⁷Department of Endocrinology and Metabolic Diseases, Polytechnic University of Marche, Ancona, Italy, ⁸Department of Immunology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran, ⁹Department of Physiology, Faculty of Medicine, Bioscience and Nursing, MAHSA University, Jenjarom, Selangor, Malaysia, ¹⁰Unidade Multidisciplinar de Investigação Biomédica (UMIB), Instituto de Ciências Biomédicas Abel Salazar (ICBAS), Universidade do Porto, Porto, Portugal, ¹¹Laboratory for Integrative and Translational Research in Population Health (ITR), Universidade do Porto, Porto, Portugal, ¹²Department of Physiology, Group of Physiology of Reproduction and Clinical Embryology, Medical School, National and Kapodistrian University of Athens, Athens, Greece, ¹³Department of Life Science and Bioinformatics, Assam University, Silchar, India, ¹⁴Fetal Health Research Center, Hope Generation Foundation, Tehran, Iran, ¹⁵Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania Luigi Vanvitelli, Caserta, Italy, ¹⁶Gamma Analisi Cliniche S.r.l, Caserta, Italy, ¹⁷Department of Women, Children and General and Specialized Surgery, University of Campania Luigi Vanvitelli, Naples, Italy, ¹⁸Histology and Microscopic Analysis Division, RASTA Specialized Research Institute (RSRI), West Azerbaijan Science and Technology Park (WASTP), Urmia, Iran, ¹⁹Department of Applied Physics, Aalto University, Espoo, Finland, ²⁰BabySoon Fertility and IVF Center, New Delhi, India, ²¹Department of Urology, Vattikuti Urology Institute, Henry Ford Health System, Detroit, MI, USA, ²²Department of Urology, Samsun Training and Research Hospital, Samsun, Turkey, ²³Department of Urology, Baskent University Faculty of Medicine, Ankara, Turkey, ²⁴Fakih IVF Fertility Center, Abu Dhabi, UAE, ²⁵Department of Urology, Hamad Medical Corporation, Doha, Qatar, ²⁶Department of Andrology, Cairo University, Cairo, Egypt, ²⁷Department of Dermatology, Venereology and Andrology, Faculty of Medicine, Sohag University, Sohag, Egypt, ²⁸Ajyal IVF Center, Ajyal Hospital, Sohag, Egypt, ²⁹Division of Andrology, Department of Urology, Lilavati Hospital and Research Centre, Mumbai, India, ³⁰Reproductive Biology, Fertility Preservation, Andrology, CECOS, Poissy Hospital, Poissy, France, ³¹Paris Saclay University, UVSQ, INRAE, BREED, Jouy-en-Josas, France

Purpose: This study provides a comprehensive analysis of research trends on the etiology, mechanisms, potential risk factors, diagnosis, prognosis, surgical and non-surgical treatment of varicocele, and clinical outcomes before and after varicocele repair.

Materials and Methods: Varicocele studies published between 1988 and 2020 were retrieved from the Scopus database on April 5, 2021. Original studies on human varicocele were included, irrespective of language. Retrieved articles were manually screened for inclusion in various sub-categories. Bibliometric data was subjected to scientometric analysis using descriptive statistics. Network, heat and geographic mapping were generated using relevant software.

Received: Sep 27, 2021 **Revised:** Oct 13, 2021 **Accepted:** Oct 25, 2021 **Published online** Jan 25, 2022

Correspondence to: Ashok Agarwal  <https://orcid.org/0000-0003-0585-1026>

Andrology Center and American Center for Reproductive Medicine, Cleveland Clinic, Mail Code X-11, 10681 Carnegie Avenue, Cleveland, OH 44195, USA.

Tel: +1-216-444-9485, **Fax:** +1-216-445-6049, **E-mail:** agarwaa@ccf.org, **Website:** www.Clevelandclinic.org/ReproductiveResearchCenter

Results: In total, 1,943 original human studies on varicocele were published. These were predominantly from the northern hemisphere and developed countries, and published in journals from the United States and Germany. Network map analysis for countries showed several interconnected nodal points, with the USA being the largest, and Agarwal A. from Cleveland Clinic, USA, being a center point of worldwide varicocele research collaborations. Studies of adolescents were underrepresented compared with studies of adults. Studies on diagnostic and prognostic aspects of varicocele were more numerous than studies on varicocele prevalence, mechanistic studies and studies focusing on etiological and risk factors. Varicocele surgery was more investigated than non-surgical approaches. To evaluate the impact of varicocele and its treatment, researchers mainly analyzed basic semen parameters, although markers of seminal oxidative stress are being increasingly investigated in the last decade, while reproductive outcomes such as live birth rate were under-reported in the literature.

Conclusions: This study analyzes the publication trends in original research on human varicocele spanning over the last three decades. Our analysis emphasizes areas for further exploration to better understand varicocele's impact on men's health and male fertility.

Keywords: Analysis, bibliometric; Clinical research; Infertility, male; Publications; Varicocele; Varicocele repair

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Varicocele is defined as an abnormal enlargement and tortuosity of veins in the pampiniform plexus [1]. One in 7 adolescents over 15 years of age is reported to have a varicocele [2]. While this prevalence is the same as that observed in the general population (approximately 15%), it is higher in infertile men [3-5]. In fact, varicocele is considered as a frequent cause of correctable infertility in males [6] as it affects up to 35% to 44% of men with primary infertility [1,3] and 45% to 80% of men with secondary infertility [7,8].

Varicoceles can have a negative impact on spermatogenesis through multiple pathways, thus resulting in a progressive decline in male fertility. The abnormalities of semen quality in infertile men with varicocele are variable ranging from oligozoospermia to complete azoospermia [9]. It is now accepted that oxidative stress (OS) is a central and common pathogenic mediator of testicular damage in varicocele, and that the exposure to scrotal heat, hypoxia, and toxic adrenal and renal metabolites enhances the generation of reactive oxygen species (ROS) [10]. Potential genetic predisposition factors and a decreased ability to fight against the effects of seminal OS have also been described to explain clinical variations of varicocele impact on sperm parameters and fertility [11-18]. Although seminal ROS in physiological amounts are essential for sperm capacitation, acrosome reaction, hyperactivation, and fertilization, increased ROS may lead to OS and result in lipid

peroxidation, protein oxidation and DNA damage [19].

During the last few decades, basic research on the pathophysiological mechanisms involved in varicocele has made it possible to answer certain clinical questions, but still some grey areas remain. For example, while the prevalence of varicocele is higher in infertile men, particularly in men with secondary infertility [1,3,7,8], it has been shown that some men can have a clinical (palpable) varicocele without affecting their sperm parameters or fertility [1,6]. It has been estimated that up to 45% of all males with clinical varicocele have normal semen parameters [1], and approximately 80% of men with varicocele are fertile [20]. Thus, despite advances in the understanding of the mechanisms involved in the pathophysiology of varicocele, it remains difficult to predict clinically which varicocele will be associated with an impairment of sperm parameters and fertility.

Regarding treatment, some questions also remain unanswered. If, according to the guidelines of the American Urological Association (AUA)/American Society for Reproductive Medicine (ASRM) and European Association of Urology (EAU) [21,22], infertility with altered sperm parameters is an indication for varicocele repair, it is difficult to predict which patient will have an improvement in sperm parameters compared to another. Similarly, it is still difficult to predict spontaneous fertility and reproductive outcomes after varicocele repair. This requires longitudinal cohort studies taking into account the grade and laterality of varicocele and the

female partner's fertility status. Randomized controlled studies on varicocele are challenging to carry out due to difficulties encountered in selecting the control arm. On the other hand, animal models are not representative of human physiology because, in animals, varicocele is artificially induced and the exposure to the varicocele is limited in duration compared to that seen in humans [16]. These reasons may explain why some questions remain unresolved and why the gap between basic research and clinical applications is not always bridged.

Scientometrics is a quantitative analysis of written scientific publications through the analysis of bibliometric data (such as authors, institutions, countries, and journals in which studies are published in specific scientific fields). It permits the investigation of developments within a specific field of research as indicated by the thematic pattern of its publications [23]. The generation of network mapping based on publication distribution and keywords is also useful in understanding the dynamics of the field of interest [24]. There are a few scientometric studies that have been conducted with regard to male infertility. These studies have mainly examined the impact of sperm DNA fragmentation (SDF) [25] and OS [26] in male infertility, the role of antioxidants in improving semen parameters [27], the contribution of assisted reproductive technologies (ART) in addressing infertility [28], and the employment of innovative tools such as proteomics in managing male infertility [29].

However, to date, there is no study available evaluating the scientific trends in the context of varicocele. Hence, the aim of this study is to perform a comprehensive analysis of publication trends in human varicocele research. This analysis will detail i) demographics of studies (countries, institutions, journals, and authors), ii) most recurrent keywords used in scientific publications on varicocele, and iii) topics of publications. The topics present in publications in the last 33 years are classified according to the following criteria: i) varicocele in adolescents and adults; ii) prevalence, pathophysiology, etiology, and risk factors of varicocele; iii) diagnosis and prognosis of varicocele; iv) methods of varicocele treatment (surgical/non-surgical interventions); and v) impact of varicocele and varicocele treatment on reproductive outcomes. Analyzing published articles from a scientometric perspective could reveal the current trends and intensity in research interest

on specific areas in human varicocele, which in turn would sketch the wider landscape of male infertility.

MATERIALS AND METHODS

1. Ethics statement

This scientometric study was performed by analyzing data retrieved from the Scopus database (<https://www.elsevier.com/solutions/scopus>) and therefore an approval by the Ethics Committee was not necessary.

2. Data source

Scopus is a bibliographic database containing abstracts and citations for academic journal articles. It covers approximately 21,000 titles from over 5,000 publishers, including 20,000 scientific, technical, medical, and social journals. An advantage of using the Scopus database is the ability to automatically analyze the literature search and generate graphs and tables based on bibliometric metrics (e.g. number of documents by year, author, affiliation, journal, country, type of document, subject area).

3. Data retrieval strategy

Data was collected on April 5, 2021 and included scientific articles from 1988 to 2020 (33 years). Articles published in 2021 were excluded by using the Scopus filter. The keyword "varicoc*" was searched as an entry term in "TITLE-ABSTRACT-KEYWORDS" fields to retrieve all relevant publications. Additional keywords ("man" OR "men" OR "adolescent*" OR "patient*" OR "human") were used to exclusively limit the search to human studies. The query AND NOT TITLE ("review*" OR "meta-analysis") AND (LIMIT-TO (DOCTYPE, "ar")) was used to retrieve only original studies. We used the asterisk "*" after the word to include all variants of the term. All retrieved publications were saved as comma-separated value (CSV) files and subsequently converted to Microsoft Excel files. The extracted studies were manually screened to exclude non-relevant articles. Only original studies on humans were included, while reviews, meta-analyses, and other types of publications (case-reports, editorials, *etc.*) were excluded by using Scopus filters and manual screening.

Afterward, the collected articles were manually validated by independent researchers for inclusion in different subcategories: varicocele in adolescents or adults, prevalence, pathophysiology, etiology and risk factors,

prognostic/diagnostic studies, surgical and non-surgical interventions, impact of varicocele or treatment on semen parameters, seminal OS, sperm DNA damage and/or SDF, testicular histology, testicular inflammation and cytokines, hormones, vasculature and blood flow, reproductive outcomes (pregnancy rate, PR; live birth rate, LBR) in natural conception or ART (Fig. 1). As each article can be included in more than one subgroup, data is reported as the number of publications and percentage out of the total number of publications

identified. Data retrieved were analyzed based on the year of publication, journal, author, affiliation, and country of origin.

4. Network, heat, and geomap analysis

Network and heat maps were generated by using the VOSviewer software (freely available at <https://vosviewer.com>) [30]. The relatedness of each node was calculated by using co-authorship as criteria, while their size reflected the number of publications. Geographic




Fig. 1. Flow diagram representing the scientometric analysis framework of original human studies on varicocele. TESE: testicular sperm extraction, SDF: sperm DNA fragmentation, ICSI: intracytoplasmic sperm injection, LBR: live birth rate, PR: pregnancy rate, IUI: intrauterine insemination, IVF: *in vitro* fertilization.




Fig. 2. Trend of publications in original human studies on varicocele from 1988 to 2020 (Scopus database). In parenthesis, guidelines published on the topic of varicocele [35-40,62,71].

mapping was created by using the Tableau Public software (freely available at <https://public.tableau.com/en-us/s/>).

RESULTS

From 1988 to 2020, 1,943 original human studies on varicocele were identified when the Scopus database was used as a search tool. The publication trend in original human studies on varicocele during these years is shown in Fig. 2.

1. Demographics of publications (countries, institutions, journals & authors)

Italy (n=285), the United States (n=255), and Turkey (n=214) have published the most on varicocele com-

pared to other countries (Fig. 3). These three countries are also the ones that have shown to be most connected with other countries in their research on varicocele (Fig. 4). The top 10 institutions and journals with the largest number of articles are reported in Table 1, 2a, and 2b, respectively. The top 10 institutions publishing original human studies on varicocele are more distributed throughout the continents (Table 1). They include North America (3 institutions in USA and 2 in Canada), Europe (3 different institutions in Italy), and other institutions in the continents of Asia (Iran, Israel, Taiwan), Africa (Egypt) and South America (Brazil).

Comparatively, out of the top 10 journals, 6 originate from the continent of Europe (2 each from the Netherlands and England, and 1 each from Germany and Switzerland), 3 are from North America, while only




Fig. 3. Geographic map illustrating the origin of publications in scientific research on varicocele in original human studies.




Fig. 4. Network map illustrating international research collaborations between countries in original human studies on varicocele.

1 is from Asia (China) (Table 2a). Most of the top 10 journals listed in Table 2a are either in the first (Q1) or second (Q2) quartile rankings for Urology based on the Scimago Journal Ranking (SJR) [31]. Those categorized as Q1 include *European Urology* (#1), *Journal of Urology* (#3), *British Journal of Urology (BJU) International* (#7), whereas those in Q2 comprise *Urology* (#28), *Urologia Internationalis* (#33), *Andrologia* (#46), and *International Urology and Nephrology* (#48). *Fertility and Sterility* is a Q1 journal of the SJR subject category for Reproductive Medicine [32] (#4), while *Zhonghua Nan Ke Xue (National Journal of Andrology)* is a Q4 journal in Medicine (miscellaneous) [33].

The top 10 authors with the highest number of publications in topics related to human studies on varicocele were Mostafa T. (n=29), Agarwal A. (n=25), Goldstein M. (n=24), Zini A. (n=24), Fujisawa M. (n=20), Glassberg K.I. (n=18), Shiraishi K. (n=16), Jarvi K. (n=15), Zampieri N. (n=15), and Tavalae M. (n=14). Network analysis showed Agarwal A. as the author contributing the most on varicocele research through collaboration with multiple different research teams (Fig. 5).

Table 1. Top 10 institutions publishing original human studies on the topic of varicocele

Ranking	Top 10 institutions	Number of publications
1	Cairo University, Egypt	75
2	Sapienza Università di Roma, Italy	36
3	Cleveland Clinic Foundation, USA	30
4	New York Presbyterian Hospital, USA	25
5	Royan Institute, Iran	22
6	University of Toronto, Canada	21
7	Università degli Studi di Verona, Italy	18
8	Mount Sinai Hospital of University of Toronto, Canada	17
9	Università degli Studi di Siena, Italy National Yang-Ming University, Taiwan	16
10	Tel Aviv University, Israel Weill Cornell Medicine, USA Universidade de São Paulo, Brazil Rabin Medical Center, Israel	15

2. Keywords analysis

A network analysis was conducted to identify the most recurrent keywords present in the title of publications. Based on the node sizes, 6 out of 65 words were found to be the most frequently used in the original studies on human varicocele: in order of frequency, these include “laparoscopic varicocelectomy”, “technique”, “experience”, “child”, “varicocele repair”, “adolescent varicocele”, and “improvement” (Fig. 6). This word analysis has limitations because sometimes synonymous keywords decrease the frequency of occurrence of a particular keyword. Hence, based on word associations (associated words belong to the same cluster and are of the same color), it appears that the two most frequently used keyword domains are adolescence (in red, Fig. 6a) and infertility (in green, Fig. 6a). Based on the period of publication, two broad periods were defined (Fig. 6b): 1988–2008 (blue, green, Fig. 6b) and 2008–2020

Table 2a. Top 10 journals publishing original human studies on the topic of varicocele

Ranking	Top 10 journals	Country of origin	Journal impact factor (2021)	SJR quartile (overall)	SJR quartile ^b (subject-based)	Number of publications
1	<i>Journal of Urology</i>	US	7.450	Q1	Q1	132
2	<i>Andrologia</i>	Germany	2.775	Q3	Q2	126
3	<i>Urology</i>	US	2.649	Q3	Q2	124
4	<i>Fertility and Sterility</i>	US	7.329	Q1	Q1	82
5	<i>Zhonghua Nan Ke Xue</i> (<i>National Journal of Andrology</i>)	China	NA	NA	Q4	46
6	<i>BJU International</i> ^a	England	5.588	Q1	Q1	36
7	<i>British Journal of Urology</i> ^a	England	-	-	-	33
8	<i>International Urology and Nephrology</i>	Netherlands	2.370	Q3	Q2	31
9	<i>Urologia Internationalis</i>	Switzerland	2.089	Q3	Q2	30
10	<i>European Urology</i>	Netherlands	20.096	Q1	Q1	28

^aBJU International was formerly known as the British Journal of Urology (1929–1999).

^bAll subject-based SJR quartiles are based on the subject of Urology, except for *Fertility and Sterility* (Reproductive Medicine) and *Zhonghua Nan Ke Xue National Journal of Andrology* (Medicine [miscellaneous]).

SJR: Scimago Journal Ranking (based on Scopus data as of April 2021), NA: not available.

Table 2b. Top 10 journals publishing original human studies on the topic of varicocele

Ranking	Top 10 journals	Semen parameters	Oxidative stress	Sperm DNA fragmentation	Prediction of clinical outcomes	Post-operative recurrence	Biochemical markers	Indication of surgical or medical treatment in varicocele diagnosis	Number of publications
1	<i>Journal of Urology</i>	62 (47.0)	6 (4.5)	3 (2.3)	70 (53.0)	37 (28.0)	12 (9.1)	46 (34.8)	132
2	<i>Andrologia</i>	84 (66.7)	24 (19.0)	11 (8.7)	44 (34.9)	9 (7.1)	37 (29.4)	18 (14.3)	126
3	<i>Urology</i>	61 (49.2)	4 (3.2)	2 (1.6)	60 (48.4)	17 (13.7)	8 (6.5)	37 (29.8)	124
4	<i>Fertility and Sterility</i>	57 (69.5)	8 (9.8)	8 (9.8)	27 (32.9)	5 (6.1)	11 (13.4)	24 (29.3)	82
5	<i>Zhonghua Nan Ke Xue</i> <i>National Journal of Andrology</i>	30 (65.2)	2 (4.3)	5 (10.9)	24 (52.2)	8 (17.4)	3 (6.5)	15 (32.6)	46
6	<i>BJU International</i> ^a	15 (41.7)	4 (11.1)	3 (8.3)	13 (36.1)	7 (19.4)	3 (8.3)	6 (16.7)	36
7	<i>British Journal of Urology</i> ^a	23 (69.7)	0 (0.0)	1 (3.0)	21 (63.6)	11 (33.3)	2 (6.1)	22 (66.7)	33
8	<i>International Urology and Nephrology</i>	11 (35.5)	1 (3.2)	2 (6.5)	14 (45.2)	6 (19.4)	8 (25.8)	11 (35.5)	31
9	<i>Urologia Internationalis</i>	14 (46.7)	0 (0.0)	0 (0.0)	11 (36.7)	6 (20.0)	3 (10.0)	9 (30.0)	30
10	<i>European Urology</i>	13 (46.4)	0 (0.0)	0 (0.0)	11 (39.3)	8 (28.6)	3 (10.7)	7 (25.0)	28

Values are presented as number of publications (%).

^aBJU International was formerly known as the British Journal of Urology (1929–1999).





Fig. 5. Network map illustrating international research collaborations between authors in original human studies on varicocele.

(light green-yellow, Fig. 6b). Keywords related to adolescence, as well as keywords for laparoscopic surgical techniques appear as older words (used before 2008).

Words such as “oxidative stress”, “infertile”, “efficacy”, “expression”, “association”, “body mass index” or those concerning “subinguinal microsurgical techniques”



Topic of publications	Value
Mechanistic studies (n=552)	
Spermatogenesis/spermiogenesis	246 (44.6)
Molecular interactions	106 (19.2)
Pathophysiology	377 (68.3)
Prognostic/diagnostic studies (n=1,285)	
Prediction of clinical outcomes	841 (65.4)
Post-operative recurrence	381 (29.6)
Biochemical markers	226 (17.6)
Indication of surgical or medical treatment in varicocele diagnosis	571 (44.4)
Studies investigating non-surgical and surgical approaches for varicocele treatment (n=1,271)	
Non-surgical approaches	260 (20.5)
Non-surgical pharmaceutical	46 (17.7)
Non-surgical antioxidants	24 (9.2)
Radiological approaches and complementary medicine	199 (76.5)
Surgical approaches for varicocele	1,130 (88.9)

were used in studies published in recent years.

Published reports on varicocele in adolescents included 355 (n=18.3%) articles, with the highest yearly trend

4. Prevalence, pathophysiology, etiology, and risk factors

There were 32 studies on varicocele prevalence (1.6%).

In mechanistic studies (n=552), 377 (68.3%) articles investigated varicocele pathophysiology, 246 (44.6%) spermatogenesis/spermiogenesis, and 106 (19.2%) molecular interactions (Table 3). A total of 222 (11.4%) articles investigated the etiology and risk factors of varicocele. The yearly publication trend was highest in 2014 with 20 published articles.

5. Prognostic and diagnostic studies

All included publications were analyzed based on prognostic or diagnostic studies on varicocele, which included the prediction of clinical outcomes, post-operative recurrence of varicocele, biochemical markers, and indication of surgical or medical treatment in varicocele diagnosis. From a total of 1,285 publications, results indicate that the most reported area was on the prediction of clinical outcomes (n=841, 65.4%), followed by the indication of surgical or medical treatment in varicocele (n=571, 44.4%) and the post-operative recurrence (n=381, 29.6%), whereas biochemical markers were reported less frequently (n=226, 17.6%) (Table 3).

6. Methods of varicocele treatment

Out of 1,943 articles retrieved by Scopus, 1,271 (65.4%) articles discussed various approaches to treat varicocele (Table 3). Surgical approaches were well studied (n=1,130, 88.9% of cases). The non-surgical approaches for varicocele analyzed (n=260, 20.5%) included three different sub-classifications such as pharmaceutical treatment (n=46, 17.7%), treatment with antioxidants (n=24, 9.2%) as well as radiological and complementary approaches (n=199, 76.5%).

7. Impact of varicocele and varicocele treatment

Among studies reporting the impact of varicocele and its treatment on outcomes (n=1,447), semen parameters are the most covered outcomes being mentioned in 915 (63.2%) articles (Fig. 7). The interest in the subject has risen steadily over time, with 2015 being the most productive year. The second most reported outcome has been vasculature and blood flow, being addressed in 314 (21.7%) articles. The interest in seminal OS is comparable to the interest in sperm DNA damage or SDF, reported in 98 (6.8%) and 92 (6.4%) publications, respectively. The interest in both seminal OS and SDF in varicocele publications has increased over time. PR and




Fig. 8. Trend of publications in original human studies investigating the impact of varicocele/varicocele treatment on reproductive outcomes. PR: pregnancy rate, LBR: live birth rate, IVF: *in vitro* fertilization, IUI: intrauterine insemination, ICSI: intracytoplasmic sperm injection, TESE: testicular sperm extraction.




Fig. 7. Trend of publications in original human studies investigating the impact of varicocele/varicocele treatment on several different outcomes. SDF: sperm DNA fragmentation.

natural conception were reported in 275 (19.0%) and 241 (16.7%) articles, respectively (Fig. 8). Interestingly, LBR has been reported in only 21 (1.5%) articles.

DISCUSSION

Based on our review of the literature, this scientometric analysis is the first of its kind focusing on varicocele. Our data showed that the trend of publication of original human studies on varicocele particularly increased during the period 2002 to 2014. Previous scientometric studies showed increasing research in male infertility in general, and hence in varicocele as a common clinical concern and cause of infertility [26,29,34]. Interestingly, the years 2002 to 2014 correspond to the years of publications of guidelines on varicocele. These include the practice guidelines on varicocele and male infertility issued by the EAU in 2002 [35] and 2005 [36], the AUA in 2004 [37], as well as ASRM in 2006 [38], 2008 [39], and 2014 [40]. Additionally, possible factors contributing to the increasing number of publications might be the increasing availability of ultrasonography that has improved the diagnosis of varicocele as well as the implementation of new surgical techniques and the need for comparison between them. The increased trend of original articles on varicocele may be an indication of intensified interest to unfold the dynamics of varicocele and male infertility.

1. Demographics of publications (countries, institutions, journals, authors)

The articles published on varicocele treatment are predominantly from top institutions in the northern hemisphere and developed countries, while only one is located in the southern hemisphere. This may result in a bias in outcomes that may not be generalized to other countries. Excluding the USA and China, research teams from Italy, Turkey, and Japan lead the total number of published papers per year in science and technology [41]. One of the contributing factors for this interest is the existence of specialized journals in these countries, particularly periodicals of national scope. Another significant factor may be the competitiveness between groups working within the same country.

The majority of original human studies on varicocele were published in journals from the USA (*The Journal of Urology* and *Urology*) and Germany (*Andrologia*). These journals belong to either the first or second

quartile of the subject category for Urology, indicating that human studies on varicocele were published in key journals within the field. The publications in the top three journals (*Journal of Urology*, *Andrologia*, and *Urology*) were mainly prognostic/diagnostic studies and those related to surgical techniques. The journal with the highest impact factor (IF) in the top 10 list was the Q1 journal, *European Urology* (IF=20.096 in 2021).

Although network map analysis for countries showed several interconnected nodal points, with the USA being the largest, this analysis also revealed that Agarwal A. from Cleveland Clinic, USA, is a center point of worldwide varicocele research collaborations, connecting three other individually isolated groups of researchers. With a prolific extent of research collaborations, Agarwal A. facilitated a global approach to the conceptualization and management of varicocele, linking the leads in varicocele research from South America (with Hallak J., Bertolla R.P., and Pasqualotto F.F.), Spain (with Gosálvez J. and Amengual M.J.), and Italy (with Gentile V. and Mazzoni G.), as well as many other established scientists (Fig. 5). Previous scientometric studies also highlighted Agarwal A. as the highest contributor in male infertility research, specifically on topics such as SDF and male infertility [25], OS in male fertility [26], the role of antioxidants in improving semen parameters [27], the contribution of ART in addressing infertility [28], as well as employing innovative tools such as proteomics to investigate and manage male infertility [29].

2. Publication topics

The analysis of the recurrent keywords provides insights into the fields towards which the research is evolving [42-44]. Word co-occurrence analysis is a method for detecting major themes in a given research field. It is used to determine the intellectual structure and the main topics of studies by using visual maps [45,46]. This analysis can even provide precise information on the topics of the studies and their evolution over time. According to this network analysis, the word “laparoscopic” has been the most used in the last few decades. As this technique has been progressively replaced by the microsurgical technique, the keywords perfectly reflect the advances in this field. Similarly, the main areas studied are adolescence and fertility, which correspond to the state of clinical practice on varicocele

management. Interestingly, the words concerning adolescence and laparoscopic techniques have been replaced in the last ten years by words concerning fertility, the efficacy of microsurgical techniques, infertility, and the associations of varicocele with clinical factors such as body mass index and OS. This shows a change in the research and clinical practice of varicocele in humans in recent years.

3. Varicocele in adolescents and adults

Studies of varicoceles in adolescents are underrepresented compared with studies of varicoceles in adults (ratio around 1:4). Since the 1990s, the management of varicocele in adolescents has been a dilemma [7,47]. In adolescents, ASRM/AUA recommend repair of varicoceles associated with a persistent reduction in testicular size homolateral to the varicocele or with semen abnormalities. EAU recommends varicocele repair in cases of ipsilateral reduction in testicular volume and evidence of progressive testicular dysfunction (weak recommendation, level of evidence 2a) [22,40]. EAU guidelines also caution against possible overtreatment of varicocele in adolescents [22], noting that many adolescents with varicocele will not experience fertility problems (level of evidence 3), and providing a reminder that in the literature long-term studies of adolescents are still rare. Additionally, ensuring follow-up in adulthood for these patients is a major challenge [48]. Furthermore, adolescent varicoceles are usually diagnosed incidentally, and the presentation of symptomatic cases may differ during pubertal development [49]. Some adolescents may be reluctant to access urologic care even if they are symptomatic and delay genital examination. In addition, collection of semen samples from post-pubertal patients is not always feasible [50] and there are no reference standards for interpreting semen analysis in this population [49]. All of the above-mentioned reasons could explain the difficulties in conducting high-quality studies on adolescent varicocele and hence explain why the management of varicocele in adolescents is still subject to question or caution today. Controversies on this topic may still remain unresolved, as our keyword analysis indicates that the focus of research has recently shifted from adolescent to adult fertility.

4. Prevalence, pathophysiology, etiology, and risk factors

Varicocele is relatively common in men, develop-

ing anytime between puberty and late adulthood, but mostly detected amongst men of reproductive age [51]. Despite this, reports on varicocele prevalence are very rare, accounting for about 1.6% of all studies. This stems from the asymptomatic nature of varicocele and the need for large and well-designed studies with more emphasis on the studied population [3].

Mechanistic studies are relatively numerous, they are mainly interested in the pathophysiology of varicocele and the mechanisms of action on spermatogenesis. Fewer studies were interested in investigating molecular markers. This may be explained considering that mechanistic studies in humans are more difficult than in animals. Studies on risk factors are also underrepresented, accounting for 11.4% of all studies. According to the scientometric analysis of keywords, the associations between varicocele and body mass index or OS seem to be the most studied ones in the last 10 years. To understand why relatively few studies have focused on etiological factors involved in the genesis of varicocele, and risk factors associated with varicocele, it should be noted that long-term cohort studies are difficult to conduct, particularly in adolescents, since follow-up should be extended to the age of fatherhood. This probably explains why etiological and risk factors are poorly studied, although these may be helpful to urologists to act preventively, adapt their follow-up and make decisions about whether or not to treat varicocele [52]. Although the latest ASRM/AUA/EAU guidelines deal well with recommendations for varicocele repair, the notion of risk factors is not addressed [21,22]. According to the keyword mapping, it is possible that these topics will be addressed in the coming years and be the focus of future guidelines.

5. Prognostic and diagnostic studies

Results emerging from the current analysis indicate that studies predicting clinical outcomes, and indications of varicocele repair, accounted for 65.4% and 44.4% of all retrieved publications on prognosis and diagnosis, respectively. Such findings reflect a great interest of researchers towards the identification and definition of predictors of success following varicocele repair. This is particularly true in the context of male infertility where the choice of the proper management plan helps select patients that would benefit from varicocele repair, and avoid performing unnecessary surgery in others. Despite numerous studies on this sub-

ject, it is still impossible to distinguish those patients who will be infertile from those who will be able to conceive spontaneously. It is also impossible to predict which patient will benefit the most from varicocele repair.

According to our analysis, varicocele recurrence is also well studied (29.6% of articles on prognosis and diagnosis). The rate of varicocele recurrence following repair ranges between 0.6% and 28.0%, depending on the chosen technique [53-56]. However, the exact factors that predispose patients to a recurrence are not clear and warrant additional research. On the other hand, biochemical markers were reported in only 17.6% of all retrieved studies on varicocele. These studies focused on understanding the pathophysiological mechanisms underlying the detrimental effects of varicocele on the reproductive health of some men. This relatively low frequency could be explained by the fact that our bibliometric analysis focused on humans, whereas biochemical markers are probably more investigated in animals.

6. Methods of varicocele treatment

Our study has shown higher interest (88.9%) in choosing surgical rather than non-surgical options (20.5%) in the management of varicocele. In 1951, varicocele repair was introduced by Tulloch [57] as a way to treat male infertility and, in 1975, this was further confirmed in 504 infertile patients using the internal spermatic vein high ligation technique [58]. This could be explained by the desire to achieve quick and more reliable therapeutic results [59] and the lack of strong evidence of the role of medical therapy [60]. The publication trend with the preference for surgical treatment is consistent with the recommendations of professional societies [21,40,61]. There is indeed a common consensus regarding the management of varicocele in infertile men among the international societies. However, the guidelines differ in the grade of evidence and in the detailed description of varicocele management. The AUA/ASRM [21] and the EAU [22,62] agree on the surgical management of clinical (palpable) varicocele in non-azoospermic infertile men with impaired sperm parameters associated with otherwise non-explained infertility (moderate recommendation and evidence level B for the AUA/ASRM, strong recommendation and level of evidence 1a for the EAU). There is also a consensus among professional societies that the gold

standard for the treatment of varicocele is surgical repair using microsurgical techniques [21,63]. From our keyword analysis, it is possible to clearly see the shift in the research focus around the world over the last 10 years, where the keyword “laparoscopic” has been replaced by “microsurgical” in publications.

Currently, there is a lack of data on the comparison of the efficacy of antioxidant treatment *versus* surgical repair in the management of infertile men with varicocele. Compared to placebo, antioxidants have been shown to improve semen parameters but not to increase PR in infertile patients with varicocele [64]. However, a recent randomized controlled trial indicated a significant increase in PR in patients treated with antioxidants after varicocelectomy than those treated solely by surgery (29% *vs.* 17.9%, $p=0.029$) [65]. Additionally, a meta-analysis showed that antioxidant therapy can improve semen quality after varicocelectomy [66]. Finally, with the terms “efficacy” and “oxidative stress” becoming more commonly used, it is likely that the role of antioxidants in this setting is becoming more popular.

7. Impacts of varicocele and varicocele repair

Manuscripts that have looked at the impact of varicocele and varicocele repair have used sperm parameters as the primary outcome measured in the majority of cases. Being a simple test, semen analysis has proved to be a useful guide to fertility and is therefore helpful in diagnosis, prognosis and as a gauge of efficacy of varicocele repair [67]. It is also more advantageous than other tests in terms of time, cost and applicability [68], and less financially burdening as a test compared to the more advanced fertility tests, justifying its common utilization in clinical practice. In order to increase the strength of evidence that semen parameters improve after varicocele repair, various studies have tried to group the available data into meta-analyses [69]. The limitation of considering the improvement of semen parameters at follow-up as the primary outcome is that improved semen parameters is not the ultimate goal for the infertile couple. Also, as the guidelines and standard values utilized vary according to the different editions of the World Health Organization (WHO) manual referred to, as a variation in results can be expected when a particular edition is followed *versus* the other [70,71]. Furthermore, conventional semen analysis does not assess the sperm changes taking place in the

female tract prior to fertilization or advanced parameters such as DNA integrity and OS [72].

In the current analysis, PR and natural conception, as well as LBR were less investigated outcomes (19.0%, 16.7%, and 1.5% of all articles, respectively). Interest in this realm has increased over the time with the highest results in 2015, although other outcome variables like testicular histology remained constant. Including the reproductive outcomes in a study requires longer time and more effort than examining the semen parameters. Hence, evaluating and reporting reproductive outcomes in a large study with an adequate sample size is particularly challenging. Also, assisted reproduction outcomes are influenced by several parameters other than semen quality, such as the ovarian response to stimulation protocols, quality of oocytes, experience and technical skills of the embryologist, therefore introducing more confounders in the analysis [73,74]. Natural conception and PR can be assessed at one time without the need for follow-up till the end of pregnancy. On the contrary, LBR is a complex outcome to measure because of the longer follow-up duration. This may hinder a large-scale study and explain why so few studies have reported this outcome over the past decades.

Finally, several factors may explain why studies pertaining to varicocele and their impact on SDF and seminal OS are low in the current literature. There are several tests evaluating SDF which are laborious, require high precision and expensive equipment along with skilled technicians, and mostly they lack in standardization, which raises the difficulty of interpretation [75-78]. There is also a lack of consensus amongst experts about which clinical scenarios these tests should be used in as well as some debate regarding whether varicocele treatment can help improve elevated SDF and OS [79]. Although it is well established that SDF and OS play a fundamental role in varicocele-induced male infertility, the opposite is not true. Elevated SDF or OS values do not necessarily lead to a diagnosis of varicocele [1,80].

8. Limitations

Despite interesting data emerging from the current study, few limitations are noted. First, our study utilized a single bibliographic database (Scopus), and therefore varicocele articles which were not indexed in this database are excluded. Future studies could endeavor to incorporate other databases such as Web of

Science and PubMed. Second, although our study was able to analyze separately the publications concerning adolescents from those concerning adults, it was not able to analyze the trends in the studied parameters among males with advanced age. Finally, an inherent limitation of the scientometric method is the retrospective nature of this analysis.

CONCLUSIONS

This study has provided an in-depth, detailed analysis of the publication trends in original human studies on varicocele, showing a general upward trend over a span of 33 years. The varicocele research landscape appears to have progressed from that of adolescent varicocele and laparoscopic varicocele repair in the early years to male fertility status, clinical efficacy of microsurgical repair, and the association between varicocele and factors such as body mass index or seminal OS in the past decade.

Our results suggest that although some aspects of varicocele are well studied, there are still many facets of varicocele and male infertility that have not been fully elucidated and therefore require greater attention. In fact, the prevalence of varicocele and risk factors remain little investigated. Similarly, the impact of varicocele and its treatment is largely analyzed through basic sperm parameters, but fewer studies have focused on seminal OS and SDF, a gap which might be filled in the coming years. In addition, the scientometric analysis showed that fewer studies looked at reproductive outcome rates of different ARTs. The paternity endpoint is rarely chosen in the current publication on varicocele, even though it would address several important questions about the impact of varicocele and its management on spontaneous or assisted fertility.

Besides well-designed, large-scale clinical studies with long-term follow-up, greater traction on the various aspects of varicocele research could be achieved through global collaborations between clinical and research institutions alike. These types of collaborative studies could help i) clarify the long-standing question of why varicocele impairs fertility only in certain males and what makes them predisposed to these changes, ii) elucidate the fundamental reasons of how and when varicocele-mediated damage develops in these males, and ultimately iii) contribute towards better clinical

practice in managing varicocele throughout the world.

Conflict of Interest

The authors have nothing to disclose.

Funding

None.

Acknowledgements

Authors are thankful to the artists from the Cleveland Clinic's Center for Medical Art & Photography for their help with the illustrations. The study was supported by the American Center for Reproductive Medicine.

Author contribution

Conceptualization: Ashok Agarwal, RF, DD, KL, RH. Writing – original draft: All the authors. Writing – review & editing: All the authors.

REFERENCES

1. Su JS, Farber NJ, Vij SC. Pathophysiology and treatment options of varicocele: an overview. *Andrologia* 2021;53:e13576.
2. Akbay E, Cayan S, Doruk E, Duce MN, Bozlu M. The prevalence of varicocele and varicocele-related testicular atrophy in Turkish children and adolescents. *BJU Int* 2000;86:490-3.
3. Alsaikhan B, Alrabeeah K, Delouya G, Zini A. Epidemiology of varicocele. *Asian J Androl* 2016;18:179-81.
4. Clavijo RI, Carrasquillo R, Ramasamy R. Varicoceles: prevalence and pathogenesis in adult men. *Fertil Steril* 2017;108:364-9.
5. Saypol DC. Varicocele. *J Androl* 1981;2:61-71.
6. Jensen CFS, Østergren P, Dupree JM, Ohl DA, Sønksen J, Fode M. Varicocele and male infertility. *Nat Rev Urol* 2017;14:523-33.
7. Gorelick JJ, Goldstein M. Loss of fertility in men with varicocele. *Fertil Steril* 1993;59:613-6.
8. Jarow JP, Coburn M, Sigman M. Incidence of varicoceles in men with primary and secondary infertility. *Urology* 1996;47:73-6.
9. Sofikitis NV, Miyagawa I, Incze P, Andrighetti S. Detrimental effect of left varicocele on the reproductive capacity of the early haploid male gamete. *J Urol* 1996;156:267-70.
10. Agarwal A, Hamada A, Esteves SC. Insight into oxidative stress in varicocele-associated male infertility: part 1. *Nat Rev Urol* 2012;9:678-90.
11. Ichioka K, Nagahama K, Okubo K, Soda T, Ogawa O, Nishiyama H. Genetic polymorphisms in glutathione S-transferase T1 affect the surgical outcome of varicocelectomies in infertile patients. *Asian J Androl* 2009;11:333-41.
12. Omar SS, Mahfouz W, Dawood W, Abo El-Wafa RAH, Ghazala RA, Zahran AM. Relation of nitric oxide synthase gene (NOS3) polymorphisms to varicocele risk and post-varicocelectomy seminal oxidative stress reduction. *Andrologia* 2020;52:e13525.
13. Gentile V, Nicotra M, Scaravelli G, Antonini G, Ambrosi S, Saccucci P, et al. ACP1 genetic polymorphism and spermatid parameters in men with varicocele. *Andrologia* 2014;46:147-50.
14. Nayeri M, Talebi AR, Heidari MM, Seifati SM, Tabibnejad N. Polymorphisms of sperm protamine genes and CMA3 staining in infertile men with varicocele. *Rev Int Androl* 2020;18:7-13.
15. Allamaneni SS, Naughton CK, Sharma RK, Thomas AJ Jr, Agarwal A. Increased seminal reactive oxygen species levels in patients with varicoceles correlate with varicocele grade but not with testis size. *Fertil Steril* 2004;82:1684-6.
16. Naughton CK, Nangia AK, Agarwal A. Pathophysiology of varicoceles in male infertility. *Hum Reprod Update* 2001;7:473-81.
17. Razi M, Tavalae M, Sarrafzadeh-Rezaei F, Moazamian A, Gharagozloo P, Drevet JR, et al. Varicocele and oxidative stress: new perspectives from animal and human studies. *Andrology* 2021;9:546-58.
18. Agarwal A, Prabakaran S, Allamaneni SS. Relationship between oxidative stress, varicocele and infertility: a meta-analysis. *Reprod Biomed Online* 2006;12:630-3.
19. Baskaran S, Finelli R, Agarwal A, Henkel R. Reactive oxygen species in male reproduction: a boon or a bane? *Andrologia* 2021;53:e13577.
20. Hamada A, Esteves SC, Agarwal A. Insight into oxidative stress in varicocele-associated male infertility: part 2. *Nat Rev Urol* 2013;10:26-37.
21. Schlegel PN, Sigman M, Collura B, De Jonge CJ, Eisenberg ML, Lamb DJ, et al. Diagnosis and treatment of infertility in men: AUA/ASRM guideline part II. *Fertil Steril* 2021;115:62-9.
22. Salonia A, Bettocchi C, Carvalho J, Corona G, Jones TH, Kadioğlu A, et al. Sexual and reproductive health [Internet]. Arnhem: European Association of Urology; c2020 [cited 2021 Oct 15]. Available from: <https://uroweb.org/guideline/sexual-and-reproductive-health/>.

23. Kim MC, Zhu Y. Scientometrics of scientometrics: mapping historical footprint and emerging technologies in scientometrics. In: Jibu M, Osabe Y, editors. *Scientometrics*. London: IntechOpen; 2018.
24. Leydesdorff L, Milojević S. *Scientometrics*. In: Wright JD, editor. *International encyclopedia of the social & behavioral sciences*. 2nd ed. Oxford: Elsevier; 2015;322-7.
25. Baskaran S, Agarwal A, Panner Selvam MK, Finelli R, Robert KA, Iovine C, et al. Tracking research trends and hotspots in sperm DNA fragmentation testing for the evaluation of male infertility: a scientometric analysis. *Reprod Biol Endocrinol* 2019;17:110.
26. Agarwal A, Baskaran S, Panner Selvam MK, Finelli R, Barbăroșie C, Robert KA, et al. Scientific landscape of oxidative stress in male reproductive research: a scientometric study. *Free Radic Biol Med* 2020;156:36-44.
27. Agarwal A, Selvam MKP, Baskaran S, Finelli R, Leisegang K, Barbăroșie C, et al. Highly cited articles in the field of male infertility and antioxidants: a scientometric analysis. *World J Mens Health* 2021;39:760-75.
28. Agarwal A, Panner Selvam MK, Baskaran S, Finelli R, Leisegang K, Barbăroșie C, et al. A scientometric analysis of research publications on male infertility and assisted reproductive technology. *Andrologia* 2021;53:e13842.
29. Agarwal A, Baskaran S, Panner Selvam MK, Barbăroșie C, Master K. Unraveling the footsteps of proteomics in male reproductive research: a scientometric approach. *Antioxid Redox Signal* 2020;32:536-49.
30. Van Eck NJ, Waltman L. Text mining and visualization using VOSviewer. *ISSI Newsl* 2011;7:50-4.
31. Scimago. Journal rankings on urology [Internet]. Madrid: Scimago; c2021 [cited 2021 Oct 15]. Available from: <https://www.scimagojr.com/journalrank.php?category=2748>.
32. Scimago. Journal rankings on reproductive medicine [Internet]. Madrid: Scimago; c2021 [cited 2021 Oct 15]. Available from: <https://www.scimagojr.com/journalrank.php?category=2743>.
33. Scimago. Zhonghua nan ke xue = National journal of andrology [Internet]. Madrid: Scimago; c2021 [cited 2021 Oct 15]. Available from: <https://www.scimagojr.com/journalsearch.php?q=61342&tip=sid&clean=0>.
34. Baskaran S, Agarwal A, Leisegang K, Pushparaj PN, Panner Selvam MK, Henkel R. An in-depth bibliometric analysis and current perspective on male infertility research. *World J Mens Health* 2021;39:302-14.
35. Weidner W, Colpi GM, Hargreave TB, Papp GK, Pomeroy JM, Ghosh C; EAU Working Group on Male Infertility. EAU guidelines on male infertility. *Eur Urol* 2002;42:313-22.
36. Dohle GR, Colpi GM, Hargreave TB, Papp GK, Jungwirth A, Weidner W; EAU Working Group on Male Infertility. EAU guidelines on male infertility. *Eur Urol* 2005;48:703-11.
37. Male Infertility Best Practice Policy Committee of the American Urological Association; Practice Committee of the American Society for Reproductive Medicine. Report on varicocele and infertility. *Fertil Steril* 2004;82 Suppl 1:S142-5.
38. Practice Committee of the American Society for Reproductive Medicine. Report on varicocele and infertility. *Fertil Steril* 2006;86(5 Suppl 1):S93-5.
39. Practice Committee of American Society for Reproductive Medicine. Report on varicocele and infertility. *Fertil Steril* 2008;90(5 Suppl):S247-9.
40. Practice Committee of the American Society for Reproductive Medicine; Society for Male Reproduction and Urology. Report on varicocele and infertility: a committee opinion. *Fertil Steril* 2014;102:1556-60.
41. National Science Foundation. Scientific and technical journal articles [Internet]. Washington, D.C.: World Bank Group; c2021 [cited 2021 Oct 15]. Available from: https://data.worldbank.org/indicator/IP.JRN.ARTC.SC?end=2018&start=2000&view=chart&year_low_desc=true.
42. Sun Y, Li C, Zhao Y, Sun J. Trends and developments in oral health literacy: a scientometric research study (1991-2020). *BDJ Open* 2021;7:13.
43. Briones-Bitar J, Carrión-Mero P, Montalván-Burbano N, Morante-Carballo F. Rockfall research: a bibliometric analysis and future trends. *Geosciences* 2020;10:403.
44. Courtial JP. A cword analysis of scientometrics. *Scientometrics* 1994;31:251-60.
45. Carrión-Mero P, Montalván-Burbano N, Paz-Salas N, Morante-Carballo F. Volcanic geomorphology: a review of worldwide research. *Geosciences* 2020;10:347.
46. Dong D, Chen ML. Publication trends and co-citation mapping of translation studies between 2000 and 2015. *Scientometrics* 2015;105:1111-28.
47. Zundel S, Szavay P, Stanasel I. Management of adolescent varicocele. *Semin Pediatr Surg* 2021;30:151084.
48. Suson KD. Transitional urology for male adolescents: what adult urologists should know. *Curr Urol Rep* 2016;17:71.
49. Chung JM, Lee SD. Current issues in adolescent varicocele: pediatric urological perspectives. *World J Mens Health* 2018;36:123-131.
50. Daudin M, Rives N, Walschaerts M, Drouineaud V, Szyman E, Koscinski I, et al. Sperm cryopreservation in adolescents and young adults with cancer: results of the French national sperm banking network (CECOS). *Fertil Steril* 2015;103:478-86.e1.

51. Canales BK, Zapzalka DM, Ercole CJ, Carey P, Haus E, Aeppli D, et al. Prevalence and effect of varicoceles in an elderly population. *Urology* 2005;66:627-31.
52. Sack BS, Schäfer M, Kurtz MP. The dilemma of adolescent varicoceles: do they really have to be repaired? *Curr Urol Rep* 2017;18:38.
53. Goldstein M, Gilbert BR, Dicker AP, Dwosh J, Gnecco C. Microsurgical inguinal varicocelectomy with delivery of the testis: an artery and lymphatic sparing technique. *J Urol* 1992;148:1808-11.
54. Homonnai ZT, Fainman N, Engelhard Y, Rudberg Z, David MP, Paz G. Varicocelectomy and male fertility: comparison of semen quality and recurrence of varicocele following varicocelectomy by two techniques. *Int J Androl* 1980;3:447-58.
55. Kaufman SL, Kadir S, Barth KH, Smyth JW, Walsh PC, White RI Jr. Mechanisms of recurrent varicocele after balloon occlusion or surgical ligation of the internal spermatic vein. *Radiology* 1983;147:435-40.
56. Marmar JL, Kim Y. Subinguinal microsurgical varicocelectomy: a technical critique and statistical analysis of semen and pregnancy data. *J Urol* 1994;152:1127-32.
57. Tulloch WS. A consideration of sterility factors in the light of subsequent pregnancies. II. Sub fertility in the male. (Tr. Edinburgh Obst. Soc. Session 104). *Edinb Med J* 1951-1952;59:29-34.
58. Dubin L, Amelar RD. Varicocelectomy as therapy in male infertility: a study of 504 cases. *Fertil Steril* 1975;26:217-20.
59. Watanabe M, Nagai A, Kusumi N, Tsuboi H, Nasu Y, Kumon H. Minimal invasiveness and effectivity of subinguinal microscopic varicocelectomy: a comparative study with retroperitoneal high and laparoscopic approaches. *Int J Urol* 2005;12:892-8.
60. Garg H, Kumar R. An update on the role of medical treatment including antioxidant therapy in varicocele. *Asian J Androl* 2016;18:222-8.
61. Jarow JP, Sharlip ID, Belker AM, Lipshultz LI, Sigman M, Thomas AJ, et al.; Male Infertility Best Practice Policy Committee of the American Urological Association Inc. Best practice policies for male infertility. *J Urol* 2002;167:2138-44.
62. Jungwirth A, Giwercman A, Tournaye H, Diemer T, Kopa Z, Dohle G, et al.; European Association of Urology Working Group on Male Infertility. European Association of Urology guidelines on male infertility: the 2012 update. *Eur Urol* 2012;62:324-32.
63. Jungwirth A, Diemer T, Kopa Z, Krausz C, Minhas S, Tournaye H. Male infertility [Internet]. Arnheim: European Association of Urology; c2019 [cited 2021 Oct 15]. Available from: <https://uroweb.org/guideline/male-infertility/>.
64. Pyrgidis N, Sokolakis I, Palapelas V, Tishukov M, Mykoniatis I, Symeonidis EN, et al. The effect of antioxidant supplementation on operated or non-operated varicocele-associated infertility: a systematic review and meta-analysis. *Antioxidants (Basel)* 2021;10:1067.
65. Kızılay F, Altay B. Evaluation of the effects of antioxidant treatment on sperm parameters and pregnancy rates in infertile patients after varicocelectomy: a randomized controlled trial. *Int J Impot Res* 2019;31:424-31.
66. Wang J, Wang T, Ding W, Wu J, Wu G, Wang Y, et al. Efficacy of antioxidant therapy on sperm quality measurements after varicocelectomy: a systematic review and meta-analysis. *Andrologia* 2019;51:e13396.
67. Agarwal A, Sharma R, Harlev A, Esteves SC. Effect of varicocele on semen characteristics according to the new 2010 World Health Organization criteria: a systematic review and meta-analysis. *Asian J Androl* 2016;18:163-70.
68. Flint M, Lampiao F, Agarwal A, du Plessis SS. Sperm assessment: traditional approaches and their indicative value. In: Nagy ZP, Varghese AC, Agarwal A, editors. *Practical manual of in vitro fertilization: advanced methods and novel devices*. New York: Springer; 2012;185-92.
69. Shridharani A, Owen RC, Elkelany OO, Kim ED. The significance of clinical practice guidelines on adult varicocele detection and management. *Asian J Androl* 2016;18:269-75.
70. Ford WC. Comments on the release of the 5th edition of the WHO Laboratory Manual for the Examination and Processing of Human Semen. *Asian J Androl* 2010;12:59-63.
71. World Health Organization (WHO). WHO laboratory manual for the examination and processing of human semen. 5th ed. Geneva: WHO; 2010.
72. Wang C, Swerdloff RS. Limitations of semen analysis as a test of male fertility and anticipated needs from newer tests. *Fertil Steril* 2014;102:1502-7.
73. Gao H, Liu DE, Li Y, Wu X, Tan H. Early prediction of live birth for assisted reproductive technology patients: a convenient and practical prediction model. *Sci Rep* 2021;11:331.
74. Go KJ. 'By the work, one knows the workman': the practice and profession of the embryologist and its translation to quality in the embryology laboratory. *Reprod Biomed Online* 2015;31:449-58.
75. Esteves SC, Zini A, Coward RM, Evenson DP, Gosálvez J, Lewis SEM, et al. Sperm DNA fragmentation testing: summary evidence and clinical practice recommendations. *Andrologia* 2021;53:e13874.
76. Agarwal A, Panner Selvam MK, Baskaran S, Cho CL. Sperm DNA damage and its impact on male reproductive health: a critical review for clinicians, reproductive professionals and

- researchers. *Expert Rev Mol Diagn* 2019;19:443-57.
77. Esteves SC, Agarwal A, Cho CL, Majzoub A. A strengths-weaknesses-opportunities-threats (SWOT) analysis on the clinical utility of sperm DNA fragmentation testing in specific male infertility scenarios. *Transl Androl Urol* 2017;6(Suppl 4):S734-60.
78. Evenson DP, Jost LK, Marshall D, Zinaman MJ, Clegg E, Purvis K, et al. Utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Hum Reprod* 1999;14:1039-49.
79. Agarwal A, Majzoub A, Esteves SC, Ko E, Ramasamy R, Zini A. Clinical utility of sperm DNA fragmentation testing: practice recommendations based on clinical scenarios. *Transl Androl Urol* 2016;5:935-50.
80. Gill K, Kups M, Harasny P, Machalowski T, Grabowska M, Lukaszuk M, et al. The negative impact of varicocele on basic semen parameters, sperm nuclear DNA dispersion and oxidation-reduction potential in semen. *Int J Environ Res Public Health* 2021;18:5977.