
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Clop, A.; Hitruhin, L.; Sengupta, B.

Rotation Bounds for Hölder Continuous Homeomorphisms with Integrable Distortion

Published in:
JOURNAL OF GEOMETRIC ANALYSIS

DOI:
[10.1007/s12220-022-00950-y](https://doi.org/10.1007/s12220-022-00950-y)

Published: 01/08/2022

Document Version
Publisher's PDF, also known as Version of record

Published under the following license:
CC BY

Please cite the original version:
Clop, A., Hitruhin, L., & Sengupta, B. (2022). Rotation Bounds for Hölder Continuous Homeomorphisms with Integrable Distortion. *JOURNAL OF GEOMETRIC ANALYSIS*, 32(8), 1-21. Article 212.
<https://doi.org/10.1007/s12220-022-00950-y>

This material is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.



Rotation Bounds for Hölder Continuous Homeomorphisms with Integrable Distortion

A. Clop¹ · L. Hitruhin² · B. Sengupta³

Received: 6 September 2021 / Accepted: 23 April 2022 / Published online: 25 May 2022
© The Author(s) 2022

Abstract

We obtain sharp rotation bounds for the subclass of homeomorphisms $f : \mathbb{C} \rightarrow \mathbb{C}$ of finite distortion which have distortion function in L^p_{loc} , $p > 1$, and for which a Hölder continuous inverse is available. The interest in this class is partially motivated by examples arising from fluid mechanics. Our rotation bounds hereby presented improve the existing ones, for which the Hölder continuity is not assumed. We also present examples proving sharpness.

Keywords Mappings of finite distortion · Quasiconformal maps · Rotation bounds

1 Introduction

We say that $f : \mathbb{C} \rightarrow \mathbb{C}$ is a mapping of finite distortion if it belongs to the Sobolev space $f \in W^{1,1}_{loc}(\mathbb{C}; \mathbb{C})$, its jacobian determinant $\det(Df) = J(\cdot, f)$ is locally integrable, and there exists a measurable function $\mathbb{K}(\cdot, f) : \mathbb{C} \rightarrow [1, +\infty]$ such that

$$|Df(z)|^2 \leq \mathbb{K}(z, f) \cdot J(z, f)$$

✉ A. Clop
albert.clop@ub.edu

L. Hitruhin
lauri.hitruhin@aalto.fi

B. Sengupta
sengupta@mat.uab.cat

¹ Department of Mathematics and Computer Science, Universitat de Barcelona, 08007 Barcelona, Catalonia, Spain

² Department of Mathematics and Systems Analysis, Aalto University, P.O. Box 11100, 00076 Aalto, Helsinki, Finland

³ Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Catalonia, Spain

at almost every point $z \in \mathbb{C}$. Above, $|Df(z)|$ stands for the operator norm of the differential matrix $Df(z)$ of f at the point z . When $\mathbb{K}(\cdot, f) \in L^\infty$ then f is said to be K -quasiregular, with $K = \|\mathbb{K}(\cdot, f)\|_\infty$ (or K -quasiconformal, if bijective). In the same way quasiregular maps arose as a generalization of holomorphic functions, mappings of finite distortion arose as a generalization of quasiregular maps partially motivated by questions in nonlinear elasticity. The authors address the interested reader to the monograph [11] for quasiregular maps in the plane, and to [1] for a background on mappings of finite distortion.

Recently, there has been a growing interest in understanding the rotational properties of planar homeomorphisms, see [2, 4, 7–10]. Special attention has been devoted to the spiraling rate of these maps. More precisely, given a homeomorphism $f : \mathbb{C} \rightarrow \mathbb{C}$ normalized by $f(0) = 0$ and $f(1) = 1$, one is interested in the growth of $|\arg(f(r))|$ as $r \rightarrow 0$. This growth represents the number of times that the image $f([r, 1])$ winds around the origin as $r \rightarrow 0$. This quantity has been proven to admit several speeds of growth which depend on the class of maps under study. In this way, it was proven in [2] that if f is K -quasiconformal then

$$|\arg(f(r))| \leq \frac{1}{2} \left(K - \frac{1}{K} \right) \log \left(\frac{1}{r} \right) + c_K, \quad \text{for all } 0 < r < 1. \tag{1}$$

In contrast, if the maps under study are homeomorphisms of finite distortion, the situation changes and the order of growth depends on the integrability of the distortion function. Namely, the second named author discovered in [8] that if $e^{\mathbb{K}(\cdot, f)} \in L^p_{loc}$ for some $p > 0$ then

$$|\arg(f(z))| \leq \frac{c}{p} \log^2 \left(\frac{1}{|z|} \right), \quad \text{for small enough } |z|,$$

and moreover, this is sharp up to the value of the constant $c > 0$. In other words, the transition between boundedness and exponential integrability of $\mathbb{K}(\cdot, f)$ results in a larger power of the logarithmic term. Further optimal results were obtained later on in [9], in the case of integrable distortion, that is, when $\mathbb{K}(\cdot, f) \in L^p_{loc}$ for some $p > 1$,

$$|\arg(f(z))| \leq \frac{c}{|z|^{\frac{2}{p}}}, \quad \text{for small enough } |z| \tag{2}$$

or even when $\mathbb{K}(\cdot, f) \in L^1_{loc}$,

$$\lim_{|z| \rightarrow 0} |z|^2 |\arg(f(z))| = 0. \tag{3}$$

The moral here is that more spiraling is allowed at the cost of relaxing the integrability properties of $\mathbb{K}(\cdot, f)$. As explained in [2, 8, 9], the local rotational properties go hand in hand with the local stretching behavior. Especially important for the argument are the estimates for the modulus of continuity of the inverse map.

It turns out mappings of finite distortion also have a role in fluid mechanics. To be precise, let us think of the planar incompressible Euler system of equations in vorticity form,

$$\begin{cases} \frac{d}{dt}\omega + (\mathbf{v} \cdot \nabla)\omega = 0 \\ \operatorname{div}(\mathbf{v}) = 0 \\ \omega(0, \cdot) = \omega_0. \end{cases} \tag{4}$$

Here $\omega = \omega(t, z) : [0, T] \times \mathbb{C} \rightarrow \mathbb{C}$ is the unknown, $\omega_0 \in L^\infty(\mathbb{C}; \mathbb{C})$ is given, and \mathbf{v} is the velocity field. The Biot-Savart law,

$$\mathbf{v} = \frac{i}{2\pi\bar{z}} * \omega$$

makes more precise the relation between \mathbf{v} and ω . As it is well known, Yudovich [15] proved existence and uniqueness of a solution $\omega \in L^\infty([0, T]; L^\infty(\mathbb{C}; \mathbb{C}))$ for any given ω_0 . In particular, the corresponding velocity field \mathbf{v} belongs to the Zygmund class, and therefore, the classical Cauchy–Lipschitz theory guarantees for the ODE

$$\begin{cases} \frac{d}{dt}X(t, z) = \mathbf{v}(t, X(t, z)) \\ X(0, z) = z \end{cases}$$

both existence and uniqueness of a flow map $X : [0, T] \times \mathbb{C} \rightarrow \mathbb{C}$. It was proven in [6] that, for small enough $t > 0$, each of the flow homeomorphisms $X_t = X(t, \cdot) : \mathbb{C} \rightarrow \mathbb{C}$ is indeed a mapping of finite distortion. Moreover, for each small value $t > 0$, there is a number $p(t) > 1$ such that the distortion function $\mathbb{K}(\cdot, X_t)$ belongs to L^p_{loc} whenever $p < p(t)$.

As mappings with L^p distortion, the mappings X_t are a bit special because both X_t and X_t^{-1} are Hölder continuous, as shown in [14], with a Hölder exponent that decays exponentially in time. This is not true in general, and mappings of L^p distortion need not have a Hölder continuous inverse, as shown in [12]. Therefore, it is a question of interest to find out if the Hölder nature of the inverse map results in better rotation bounds. Indeed, even though the bounds obtained in [9] can be applied to X_t , the Hölder continuous nature of X_t^{-1} provides a significant improvement to (2). We describe this improvement in our main Theorem.

Theorem 1 *Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a homeomorphism of finite distortion such that $f(0) = 0$ and $f(1) = 1$, and assume that $\mathbb{K}(\cdot, f) \in L^p_{loc}$ for some $p > 1$. Suppose also that*

$$|f(x) - f(y)| \geq C|x - y|^\alpha, \quad \text{if } |x - y| \text{ is small,}$$

for some $\alpha > 1$. Then

$$|\arg(f(z))| \leq C\sqrt{\alpha}|z|^{-\frac{1}{p}}\log^{\frac{1}{2}}\left(\frac{1}{|z|}\right)$$

whenever $|z|$ is small enough.

In contrast with (2) and (3), the existence of a Hölder continuous inverse allows the power term exponent to be halved, although then the logarithmic term needs to be included. As an application, we can estimate the spiraling rate of X_t .

As an application, we can estimate the spiraling rate of X_t for small times. The rotational behavior of X_t is nowadays studied a lot. For instance, in the case of ω_0 being close to the characteristic function of the unit disk, the article [5] provides bounds for the winding number of most of the trajectories $\{X_t(z)\}_{t>0}$ as $t \rightarrow \infty$. Here, instead, we do not evaluate the rotational behavior at large times, but look instead at spiraling bounds in the space variable for a fixed and small enough time.

Corollary 2 *Given $\omega_0 \in L^\infty(\mathbb{C}; \mathbb{C})$, let \mathbf{v} be the velocity field of Yudovich’s solution to (4), and let X_t be its flow. Then there exists a constant $C > 0$ such that*

$$\left| \arg \left(\frac{X_t(z) - X_t(0)}{X_t(1) - X_t(0)} \right) \right| \leq C \log^{\frac{1}{2}} \left(\frac{1}{|z|} \right) |z|^{-t\|\omega_0\|_\infty} \exp(Ct\|\omega_0\|_\infty)$$

if both $|z|$ and $t > 0$ are small enough.

In particular, if $z = \frac{1}{n}, n = 1, 2, \dots$ and we fix a time t_0 small enough, then the curve $X_{t_0}([\frac{1}{n}, 1])$ cannot wind around $X_{t_0}(0)$ more than a multiple of

$$n^{t_0\|\omega_0\|_\infty} (\log n)^{\frac{1}{2}} e^{Ct_0\|\omega_0\|_\infty}$$

times. Toward the optimality of Theorem 1, we can show the following.

Theorem 3 *Given an increasing, onto homeomorphism $h : [0, +\infty) \rightarrow [0, +\infty)$, and a real number $p > 1$, there exists a homeomorphism $\bar{f} : \mathbb{C} \rightarrow \mathbb{C}$ with the following properties:*

- \bar{f} is a mapping of finite distortion, with $\mathbb{K}(\cdot, \bar{f}) \in L^p_{loc}$.
- $\bar{f}(0) = 0, \bar{f}(1) = 1$.
- If $\alpha > \frac{3p}{p-1}$, then $|\bar{f}(x) - \bar{f}(y)| \geq C|x - y|^\alpha$ whenever $|x - y| < 1$.
- There exists a decreasing sequence $\{r_n\}$, with $r_n \rightarrow 0+$ as $n \rightarrow \infty$, for which

$$|\arg(\bar{f}(r_n))| \geq r_n^{-\frac{1}{p}} \log^{\frac{1}{2}} \left(\frac{1}{r_n} \right) h(r_n).$$

Since h can be chosen to approach 0 at any speed, Theorem 3 shows that the order provided in Theorem 1 is sharp.

The paper is structured as follows. In Sect. 2, we give the basic preliminaries. In Sect. 3, we prove Theorem 1 and Corollary 2. Finally, we prove Theorem 3 in Sect. 4.

2 Preliminaries

A mapping $f : \mathbb{C} \rightarrow \mathbb{C}$ is said to be Hölder continuous, or simply Hölder from above, if there exist constants $C > 0, d > 0$, and $\alpha > 0$ such that for any two points $x, y \in \mathbb{C}$ and $\alpha \in \mathbb{R}^+ \setminus \{0\}$ with $|x - y| < d$, one has

$$|f(x) - f(y)| \leq C|x - y|^\alpha$$

Similarly, we say f is Hölder from below if there are constants $C, \beta > 0$ such that for any two points $x, y \in \mathbb{C}$ and $\beta \in \mathbb{R}^+ \setminus \{0\}$ with $|x - y| < d$, one has

$$|f(x) - f(y)| \geq \bar{C}|x - y|^\beta$$

A mapping $f : \mathbb{C} \rightarrow \mathbb{C}$ is called bi-Hölder if it is both Hölder from above and from below.

Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a mapping of finite distortion and fix a point $z_0 \in \mathbb{C}$. In order to study the pointwise rotation of f at the point z_0 , one usually fixes an argument $\theta \in [0, 2\pi)$, and then looks at how the quantity

$$\arg(f(z_0 + te^{i\theta}) - f(z_0))$$

changes as the parameter t goes from 1 to a small r . This can also be understood as the winding of the path $f([z_0 + re^{i\theta}, z_0 + e^{i\theta}])$ around the point $f(z_0)$. As we are interested in the maximal pointwise spiraling, we need to normalize and then retain the maximum over all directions θ ,

$$\sup_{\theta \in [0, 2\pi)} |\arg(f(z_0 + re^{i\theta}) - f(z_0)) - \arg(f(z_0 + e^{i\theta}) - f(z_0))|. \tag{5}$$

Then, the maximal pointwise rotation is precisely the behavior of the above quantity (5) when $r \rightarrow 0^+$. In this way, we say that the map f spirals at the point z_0 with a rate g , where $g : [0, \infty) \rightarrow [0, \infty)$ is a decreasing continuous function, if

$$\limsup_{r \rightarrow 0^+} \frac{\sup_{\theta \in [0, 2\pi)} |\arg(f(z_0 + re^{i\theta}) - f(z_0)) - \arg(f(z_0 + e^{i\theta}) - f(z_0))|}{g(r)} = C \tag{6}$$

for some constant $0 < C < \infty$. Finding maximal pointwise rotation for a given class of mappings equals finding the maximal spiraling rate for this class. Note that in (6) we must use limit superior as the limit itself might not exist. Furthermore, for a given mapping f there might be many different sequences $r_n \rightarrow 0$ along which it has profoundly different rotational behavior.

Our proof of Theorem 1 relies heavily on the modulus of path families. We give here the main definitions, and address the interested reader to [13] for a closer look at the topic. The image of a line segment I under a continuous mapping is called a path, and we denote by Γ a family of paths. Given a path family Γ , we say that a Borel

measurable function ρ is *admissible* for Γ if any rectifiable $\gamma \in \Gamma$ satisfies

$$\int_{\gamma} \rho(z) dz \geq 1.$$

The *modulus of the path family* Γ is defined by

$$M(\Gamma) = \inf_{\rho \text{ admissible}} \int_{\mathbb{C}} \rho^2(z) dA(z),$$

where $dA(z)$ denotes the Lebesgue measure on \mathbb{C} . As an intuitive rule, the modulus is big if the family Γ has *lots* of short paths, and it is small if the paths are long and there are *not many* of them.

We will also need a weighted version of the modulus. Any measurable, locally integrable function $\omega : \mathbb{C} \rightarrow [0, \infty)$ will be called a *weight function*. In our case, ω will always be the distortion function $\mathbb{K}(\cdot, f)$ of some map f . Then, we define the weighted modulus $M_{\omega}(\Gamma)$ by

$$M_{\omega}(\Gamma) = \inf_{\rho \text{ admissible}} \int_{\mathbb{C}} \rho^2(z) \omega(z) dA(z).$$

Finally, we need the modulus inequality

$$M(f(\Gamma)) \leq M_{\mathbb{K}(\cdot, f)}(\Gamma) \tag{7}$$

which holds for any mapping f of finite distortion for which the distortion $\mathbb{K}(\cdot, f)$ is locally integrable, proven by the second named author in [9].

3 Spiraling Bounds

We will write Theorem 1 in the following, clearly equivalent, form.

Theorem 4 *Let f be a homeomorphism of finite distortion with distortion $\mathbb{K}(\cdot, f) \in L^p(\mathbb{C})$, $p > 1$, normalized by $f(0) = 0$ and $f(1) = 1$. Assume that it satisfies the following condition,*

$$|f(x) - f(y)| \geq C|x - y|^{\alpha}$$

whenever $|x - y|$ is small. Then the winding number $n(z_0)$ of the image of the line segment $\left[z_0, \frac{z_0}{|z_0|} \right]$ around the image of the origin is bounded from above by

$$n(z_0) \leq C\sqrt{\alpha} |z_0|^{-\frac{1}{p}} \log^{\frac{1}{2}} \left(\frac{1}{|z_0|} \right)$$

Proof We would like to prove this theorem using the modulus inequality for homeomorphisms of finite distortion (7) following the presentation in [9]. At first, we would like to estimate the modulus term $M_{\mathbb{K}(\cdot, f)}(\Gamma)$ from above. To this end, let us choose an arbitrary point $z_0 \in \mathbb{C} \setminus \{0\}$ such that $|z_0| < 1$. Without loss of generality, we may assume that z_0 lies on the positive side of the real axis. Next, let us fix the line segments $E = [z_0, 1]$ and $F = (-\infty, 0]$, and Γ be the family of paths connecting a point in E to a point in F . Also, let us fix balls $B_j = B(2^j z_0, 2^j z_0)$, $j \in \{0, 1, \dots, n\}$ and n is the smallest integer such that $2^n z_0 \geq 1$. Define

$$\rho_0(z) = \begin{cases} \frac{2}{r(B_0)} & \text{if } z \in B_0 \\ \frac{2}{r(B_1)} & \text{if } z \in B_1 \setminus B_0 \\ \vdots & \\ \frac{2}{r(B_n)} & \text{if } z \in B_n \setminus B_{n-1} \\ 0 & \text{otherwise} \end{cases}$$

Note that any $z \in E$ belongs to some ball $\frac{1}{2} B_j$ and that $\rho_0(z) \geq \frac{2}{r(B_j)}$, whenever $z \in B_j$. This implies, since $B_j \cap F = \emptyset$ for every j , that $\rho_0(z)$ is admissible with respect to Γ . Hence we can estimate the modulus from above by

$$\begin{aligned} M_{\mathbb{K}(\cdot, f)}(\Gamma) &= \inf_{\rho \text{ admissible}} \int_{\mathbb{C}} \mathbb{K}(\cdot, f) \rho^2(z) \, dA(z) \\ &\leq \int_{\mathbb{C}} \mathbb{K}(\cdot, f) \rho_0^2(z) \, dA(z) \\ &\leq \|\mathbb{K}(\cdot, f)\|_{L^p(B(0,4))} \left(\int_{B(0,4)} \rho_0^{\frac{2p}{p-1}}(z) \, dA(z) \right)^{\frac{p-1}{p}} \\ &\leq c_{f,p} \left(\int_{B(0,4)} \rho_0^{\frac{2p}{p-1}}(z) \, dA(z) \right)^{\frac{p-1}{p}} \end{aligned}$$

Let us now estimate the integral term by using the definition of ρ_0 .

$$\begin{aligned} \int_{B(0,4)} \rho_0^{\frac{2p}{p-1}}(z) \, dA(z) &\leq \sum_{j=0}^n \int_{B_j} \left(\frac{2}{r(B_j)} \right)^{\frac{2p}{p-1}} \, dA(z) \\ &= \sum_{j=0}^n |B_j| \left(\frac{2}{r(B_j)} \right)^{\frac{2p}{p-1}} \\ &= c_p \sum_{j=0}^n \frac{(r(B_j))^2}{(r(B_j))^{\frac{2p}{p-1}}} \\ &= c_p \sum_{j=0}^n \frac{1}{z_0^{\frac{2}{p-1}}} \frac{1}{2^{\frac{2j}{p-1}}} \end{aligned}$$

$$= c \rho z_0^{-\frac{2}{p-1}} \sum_{j=0}^n \frac{1}{2^{\frac{2j}{p-1}}}$$

The series $\sum_{j=0}^n \frac{1}{2^{\frac{2j}{p-1}}}$ converges to a constant depending on p for any fixed $p > 1$. Therefore,

$$M_{\mathbb{K}(\cdot, f)}(\Gamma) \leq c_{f, p} z_0^{-\frac{2}{p}}$$

Next, we would like to estimate the modulus term $M(f(\Gamma))$ from below. Let us start with the definition of $M(f(\Gamma))$ in polar coordinates

$$\begin{aligned} M(f(\Gamma)) &= \inf_{\rho \text{ admissible}} \int_{\mathbb{C}} \rho^2(z) \, dA(z) \\ &= \inf_{\rho \text{ admissible}} \int_0^{2\pi} \int_0^\infty \rho^2(r, \theta) r \, dr \, d\theta \end{aligned}$$

and provide a lower bound for

$$\int_0^\infty \rho^2(r, \theta) r \, dr$$

for an arbitrary direction $\theta \in [0, 2\pi)$ and an arbitrary admissible ρ . To this end, we fix a direction θ and consider the half-line L_θ starting from the origin in the direction θ . Let us choose points $z_0 \leq t_2 < t_0 \leq 1$ so that the image set $f(E)$ winds once around the origin when z moves from a point t_0 to a point t_2 along E and $f(t_0) \in L_\theta$. Since the mapping f is a homeomorphism and the path $f(F)$ contains both the origin and points with big modulus, as F is unbounded, the path $f(F)$ must intersect the line segment $(f(t_2), f(t_0))$ at least once, say at a point $f(t_1)$, with $t_1 \in F$. We can choose t_1 and t_0 such that there are no points from the paths $f(E)$ and $f(F)$ in the line segment $(f(t_1), f(t_0))$, which thus belongs to the path family $f(\Gamma)$. Since the path $f(E)$ cycles around the origin $n(z_0) = \left\lfloor \frac{|\arg(f(z_0)) - \arg(f(1))|}{2\pi} \right\rfloor$ times, we can find at least

$$n(z_0) = \left\lfloor \frac{|\arg(f(z_0)) - \arg(f(1))|}{2\pi} \right\rfloor - 1$$

such disjoint line segments belonging to the path family $f(\Gamma)$. Note that $n(z_0)$ does not depend on the direction θ . Since we are interested in extremal rotation, it can be assumed that $f(E)$ winds around the origin at least once, which makes it clear that $n(z_0)$ is non-negative. Now, the $n(z_0)$ disjoint line segments can be written in the form $(x_j e^{i\theta}, y_j e^{i\theta}) \subset L_\theta$, where $j \in \{1, 2, \dots, n(z_0)\}$ and x_j, y_j are positive real numbers satisfying

$$0 < r_f \leq x_1 < y_1 < \dots < x_{n(z_0)} < y_{n(z_0)} \leq c_f$$

where $c_f = \sup_{z \in E} |f(z)|$ and $r_f = \min_{z \in E} |f(z)|$. Here, neither c_f nor r_f depends on θ or z_0 . So, one could write

$$\int_0^\infty \rho^2(r, \theta) r dr \geq \sum_{j=1}^{n(z_0)} \int_{x_j}^{y_j} \rho^2(r, \theta) r dr.$$

Next, let us consider the Hölder inequality with the functions $f(r) = \rho\sqrt{r}$ and $g(r) = \frac{1}{\sqrt{r}}$, which after squaring both sides gives

$$\int_{x_j}^{y_j} \rho^2(r, \theta) r dr \geq \left(\int_{x_j}^{y_j} \rho(r, \theta) dr \right)^2 \left(\int_{x_j}^{y_j} \frac{1}{r} dr \right)^{-1} \geq \frac{1}{\log\left(\frac{y_j}{x_j}\right)}.$$

The last inequality holds true as ρ is admissible with respect to $f(\Gamma)$ where the line segments $(x_j e^{i\theta}, y_j e^{i\theta})$ belong to the path family $f(\Gamma)$. Therefore,

$$\int_0^\infty \rho^2(r, \theta) r dr \geq \sum_{j=1}^{n(z_0)} \frac{1}{\log\left(\frac{y_j}{x_j}\right)}.$$

It is quite clear from the definition of c_f that

$$\sum_{j=1}^{n(z_0)} \frac{1}{\log\left(\frac{y_j}{x_j}\right)} \geq \sum_{j=1}^{n(z_0)-1} \frac{1}{\log\left(\frac{x_{j+1}}{x_j}\right)} + \frac{1}{\log\left(\frac{c_f}{x_{n(z_0)}}\right)}.$$

Next, let us consider the AM-HM inequality. For every positive real number a_j ,

$$\sum_{j=1}^n a_j \geq \frac{n^2}{\sum_{j=1}^n \frac{1}{a_j}}.$$

At this point, we would like to use AM-HM with the precise choices

$$a_j = \frac{1}{\log\left(\frac{x_{j+1}}{x_j}\right)} \quad \text{if } j \in \{1, 2, \dots, n(z_0) - 1\}, \quad \text{and} \quad a_{n(z_0)} = \frac{1}{\log\left(\frac{c_f}{x_{n(z_0)}}\right)},$$

which give

$$\sum_{j=1}^{n(z_0)} \frac{1}{\log\left(\frac{y_j}{x_j}\right)} \geq \frac{n^2(z_0)}{\log\left(\frac{c_f}{x_1}\right)} \geq \frac{n^2(z_0)}{\log\left(\frac{c_f}{r_f}\right)}.$$

Therefore,

$$\int_0^\infty \rho^2(r, \theta) r dr \geq \frac{n^2(z_0)}{\log\left(\frac{c_f}{r_f}\right)}.$$

The constant c_f can be defined as $\max_{z \in \mathbb{D}} |f(z)|$, which is finite and does not depend on either θ or z_0 , and thus, it is irrelevant at the limit $z_0 \rightarrow 0$. On the other hand, the constant r_f must be estimated using the Hölder modulus of continuity assumption on our mapping f , that is

$$|f(z_0)| \geq C|z_0|^\alpha$$

for sufficiently small z_0 . This combined with the estimate above gives that

$$M(f(\Gamma)) \geq \frac{n^2(z_0)}{C\alpha \log\left(\frac{1}{|z_0|}\right)}$$

Now, using the modulus inequality, we get

$$\frac{n^2(z_0)}{C\alpha \log\left(\frac{1}{|z_0|}\right)} \leq c_{f,p} \left(\frac{1}{|z_0|}\right)^{\frac{2}{p}}$$

which implies the desired estimate. □

Proof of Corollary 2 Corollary 2 follows immediately after noting that one can take $f = X_t$ in Theorem 1. Indeed, from [6, Corollary 3], we know that X_t belongs to $W^{1,p}$ for any $p < \frac{2}{t\|\omega_0\|_\infty}$, provided that $0 < t < \frac{2}{\|\omega_0\|_\infty}$. Since $J(\cdot, X_t) = 1$ due to the incompressibility, it then follows X_t is a homeomorphism with finite distortion, and moreover $\mathbb{K}(\cdot, X_t) \in L^p_{\text{loc}}$ for $p < \frac{1}{t\|\omega_0\|_\infty}$. Especially, if t is so small that $0 < t < \frac{1}{\|\omega_0\|_\infty}$ then one may take $p > 1$. Also, we recall from [14] (see also [3]) that X_t^{-1} is α -Hölder continuous with some exponent $\alpha \geq e^{-ct\|\omega_0\|_\infty}$ for some $c > 0$. Hence, Theorem 1 applies to $f = X_t$ and the claim follows. □

4 Optimality of Spiraling

We will get Theorem 3 in two steps. In the first step, we will construct a map which *only rotates*. This map will already give us the optimal result (in the power scale). In the second step, we will strengthen this up with a second map, that *both rotates and stretches*. This second map is going to be the optimal one.

Given an arbitrary annulus $A = B(0, R) \setminus B(0, r)$, we define the corresponding rotation map as

$$\phi_A(z) = \begin{cases} z & |z| > R \\ z e^{i\alpha \log \frac{|z|}{R}} & r \leq |z| \leq R \\ z e^{i\alpha \log \frac{r}{R}} & |z| < r \end{cases}$$

Here $0 < r < R$, and $\alpha \in \mathbb{R}$. One must note that $\phi_A : \mathbb{C} \rightarrow \mathbb{C}$ is bilipschitz (i.e., both ϕ_A and its inverse are Lipschitz), hence quasiconformal (its quasiconformality constant depends only on α), and moreover, it is conformal outside the annulus A . Note also that ϕ_A leaves fixed all circles centered at 0, since $|\phi_A(te^{i\theta})| = t$ for each $t > 0$ and $\theta \in \mathbb{R}$. Finally, a direct calculation shows for the jacobian determinant that $J(z, \phi_A) = 1$ for each z .

Next, we fix a sequence $\{r_n\}$ such that $0 < r_{n+1} < \frac{r_n}{2e}$ and $r_1 < \frac{1}{e}$. Also, let $R_n = er_n$. These assumptions make sure that $2r_{n+1} < R_{n+1} < \frac{r_n}{2}$. Let us now construct disjoint annuli $A_n = B(0, R_n) \setminus B(0, r_n)$, and set $\{f_n\}_n$ to be a sequence of maps, constructed in an iterative way as follows. For $n = 1$, we set

$$f_1(z) = \phi_{A_1}(z) = \begin{cases} z & |z| > R_1 \\ z e^{i\alpha_1 \log \frac{|z|}{R_1}} & r_1 \leq |z| \leq R_1 \\ z e^{-i\alpha_1} & |z| < r_1 \end{cases}$$

where $\alpha_1 \in \mathbb{R}$, $\alpha_1 \geq 1$, is to be determined later. We then define f_n for $n \geq 2$ as

$$f_n(z) = \phi_{f_{n-1}(A_n)} \circ f_{n-1}(z)$$

again for some values $\alpha_n \in \mathbb{R}$, $\alpha_n \geq 1$, to be determined later. Clearly, each $f_n : \mathbb{C} \rightarrow \mathbb{C}$ is quasiconformal, and conformal outside the annuli A_i , $i = 1, \dots, n$. It is also clear that $f_n(z) = f_{n-1}(z)$ on the unbounded component of $\mathbb{C} \setminus f_{n-1}(A_n)$ (i.e., outside of $B(0, R_n)$). This proves that the sequence f_n is uniformly Cauchy, and hence, it converges to a map f , that is,

$$f = \lim_{n \rightarrow \infty} f_n$$

which is again a homeomorphism by construction. Now, since f_n is quasiconformal for every n and $f_n(z) = f_{n-1}(z)$ everywhere except inside the ball $B(0, R_n)$, where $R_n \rightarrow 0$ as $n \rightarrow \infty$, the limit map f is absolutely continuous on almost every line parallel to the coordinate axes and differentiable almost everywhere.

It is helpful to note that each f_n leaves fixed all circles centered at the origin, so in particular we have $f_n(A_j) = A_j$ for each j , and therefore $\phi_{f_{n-1}(A_n)} = \phi_{A_n}$. Direct

calculation shows that

$$|D\phi_{A_n}(z)| = |\partial\phi_{A_n}(z)| + |\bar{\partial}\phi_{A_n}(z)| = \begin{cases} 1 & |z| > R_n \\ \frac{|2+i\alpha_n|+|\alpha_n|}{2} & r_n \leq |z| \leq R_n \\ 1 & |z| < r_n \end{cases}$$

which allows us to estimate that

$$|\partial f(z)| + |\bar{\partial} f(z)| \leq 2\alpha_n \quad \text{whenever } z \in A_n,$$

and $|Df(z)| \leq 1$ otherwise. Therefore, in order to have $Df(z) \in L^1_{loc}(\mathbb{C})$, it suffices that

$$\sum_n \alpha_n r_n^2 < +\infty. \tag{8}$$

This, together with the absolute continuity, guarantees $f \in W^{1,1}_{loc}(\mathbb{C})$. Also, since f is a homeomorphism, we have that $J_f(z) \in L^1_{loc}(\mathbb{C})$, and in fact $J(z, f) = 1$ at almost every $z \in \mathbb{C}$. Therefore, f is a homeomorphism of finite distortion, with distortion function

$$\mathbb{K}(z, f) = \frac{|Df(z)|^2}{J(z, f)} \leq \begin{cases} 4\alpha_n^2 & z \in A_n, \\ 1 & \text{otherwise.} \end{cases}$$

Especially, in order to have $\mathbb{K}(\cdot, f) \in L^p_{loc}$, it suffices to ensure the convergence of the series

$$\sum_{n=1}^{\infty} |A_n|(4\alpha_n^2)^p \simeq \sum_{n=1}^{\infty} \alpha_n^{2p} r_n^2 \tag{9}$$

which can be done by choosing α_n properly. Note that if (9) holds, then also (8) holds, because our choice of α_n will guarantee $\alpha_n \geq 1$. The last restriction to choose our α_n comes from rotational behavior of f . It is clear from the above construction that $f(0) = 0$, $f(1) = 1$ and

$$|\arg(f(r_n))| \geq \left| \arg\left(\left(\frac{1}{e}\right)^{1+i\alpha_n}\right) \right| = \alpha_n$$

for every r_n . Since we want our map to be optimal for Theorem 1, we may be tempted to choose $\alpha_n = r_n^{-1/p} \log^{1/2}(1/r_n)$. Unfortunately such a choice does not meet the requirement (9). The same problem occurs if we simply choose $\alpha_n = r_n^{-1/p}$. So we choose

$$\alpha_n = h(r_n) r_n^{-1/p}.$$

Here, $h : [0, \infty) \rightarrow [0, \infty)$ is any monotonically decreasing gauge function such that $\lim_{r \rightarrow 0^+} h(r) = 0$. With this choice, (9) is fulfilled if the series

$$\sum_{n=1}^{\infty} h(r_n)^{2p} < +\infty.$$

But this can always be done by simply reducing the already chosen values of r_n , for instance if $h(r_n) < \frac{1}{n^{1/2}}$. Note that this does not provide full optimality for Theorem 1, but it already gives the right order (in the power scale).

We now show that f is Hölder continuous with exponent $1 - \frac{1}{p}$. For this, let us recall that our map f is a limit of iterates of logarithmic spiral maps inside the annuli $A_n = B(0, R_n) \setminus B(0, r_n)$. In particular, as shown in [2], if $\gamma \in \mathbb{R}$ then the basic logarithmic spiral map $g(z) = z|z|^{i\alpha} = ze^{i\gamma \log|z|}$ is L -bilipschitz, for a constant L such that $|\gamma| = L - \frac{1}{L}$. When $|\gamma|$ is large, L is large as well and so one roughly has $|\gamma| \simeq L$. Since our f_n behaves on the annulus A_n as a spiral map with $|\gamma| = \alpha_n$, we deduce the bilipschitz constant of f_n on A_n is

$$L \simeq |\gamma| = \alpha_n = h(r_n) r_n^{-1/p}.$$

Let us now start the proof. To this end, let us consider two arbitrary points x and y in $\mathbb{D} \setminus \{0\}$. We first consider the case where $x, y \in A_n$. In this case, $f(x) = f_n(x)$ and $f(y) = f_n(y)$. Since $r_n > C|x - y|$, we have

$$\begin{aligned} |f(x) - f(y)| &= |f_n(x) - f_n(y)| \lesssim h(r_n) r_n^{-1/p} |x - y| \\ &\leq h(r_n) \left(\frac{C}{|x - y|} \right)^{\frac{1}{p}} |x - y| \\ &\leq C|x - y|^{1 - \frac{1}{p}} \end{aligned}$$

where we have used the bilipschitz nature of f_n on A_n .

We now assume that $x, y \in D_n = B(0, r_n) \setminus B(0, R_{n+1})$. On that set f is of the form $ze^{i\beta}$, where $\beta \in \mathbb{R} \setminus \{0\}$, which is clearly an isometry.

Next, we take $x \in A_n$ and $y \in D_n$. In particular, $|x| \geq |y|$. Then let w be any point on the outer boundary of D_n joining x and y . We have

$$\begin{aligned} |f(x) - f(y)| &\leq |f(x) - f(w)| + |f(w) - f(y)| \\ &\leq C|x - w|^{1 - \frac{1}{p}} + |w - y| \\ &\leq 2C|x - y|^{1 - \frac{1}{p}} \end{aligned}$$

The same happens if $x \in D_{n-1}$ and $y \in A_n$.

So it just remains to see what happens when $x \in A_n = B(0, R_n) \setminus B(0, r_n)$ and $y \in B(0, R_{n+1})$. Let L be the line joining x and y . We divide it into three parts, viz.,

L_1, L_2 and L_3 . L_1 connects x to a point a on the inner boundary of A_n , so that

$$|f(x) - f(a)| = |f_n(x) - f_n(a)| \leq C|x - a|^{1-\frac{1}{p}}$$

Next, L_2 connects a to b , which is the closest point to y where the line L crosses the inner boundary of D_n . From $2R_{n+1} < r_n < \frac{R_n}{2}$ we get that $|f(a)| > 2|f(b)|$. Also, since $a, b \in D_n$ and f is an isometry there, we get

$$|f(b) - f(y)| \leq 2|f(b)| \leq 2|f(a) - f(b)| = 2|a - b|$$

Summarizing

$$\begin{aligned} |f(x) - f(y)| &\leq |f(x) - f(a)| + |f(a) - f(b)| + |f(b) - f(y)| \\ &\leq C|x - a|^{1-\frac{1}{p}} + |a - b| + 2|a - b| \\ &\leq C|x - y|^{1-\frac{1}{p}} \end{aligned}$$

The case $x \in D_n$ and $y \in B(0, r_{n+1})$ can be proved in a similar manner. Therefore, we have covered all the possible cases. Since the set $\mathbb{D} \setminus \{0\}$ is partitioned by separated annuli A_n and D_n , it is clear that we have proved that f is Hölder continuous with exponent $1 - \frac{1}{p}$. At this point, it is worth noting that this regularity could also be proven by means of the Sobolev embedding. Indeed, we proved above that $\mathbb{K}(\cdot, f) \in L^p_{loc}$, and also that the Jacobian determinant is constantly 1. This together implies that $Df \in L^{2p}_{loc}$.

Now we show that also f^{-1} is Hölder continuous. Indeed, let us recall that f is the limit of iterates of logarithmic spiral maps inside the annuli and conformal outside. Now, f^{-1} can be constructed using the same building blocks as f itself, just changing the sign of α_n at each step. This is possible because the inverse of a logarithmic spiral map is the same spiral map, just the direction of rotation is opposite of the original map. Since it is clear that the direction of rotation does not play any role in the proof of Hölder continuity of f , this implies that f^{-1} is also Hölder from above. Thus f is Hölder from below as well.

As we said before, the above example approaches the borderline stated in Theorem 1, but it does not attain full optimality yet. To this end, we have to modify it by adding to our building blocks a stretching factor. This is done by replacing, at each iterate, the logarithmic spiral map $z|z|^{i\alpha} = ze^{i\alpha \log |z|}$ by a complex power $z|z|^{q+i\alpha} = z|z|^q e^{i\alpha \log |z|}$. We now proceed with the details.

So, similarly as in the previous construction, we fix a rapidly decreasing sequence $\{r_n\}$ such that $r_{n+1} < \frac{r_n}{2e}$ and $r_1 < \frac{1}{e}$. Also, let $R_n = er_n$. Given an arbitrary annulus $A = B(0, R) \setminus B(0, r)$, we define the corresponding radial stretching combined with

rotation map as follows:

$$\phi_A(z) = \begin{cases} z & |z| > R \\ z \left| \frac{z}{R} \right|^{q-1} e^{i\alpha \log \frac{|z|}{R}} & r \leq |z| \leq R \\ z \left(\frac{r}{R} \right)^{q-1} e^{i\alpha \log \frac{r}{R}} & |z| < r \end{cases} \tag{10}$$

Note that this time we will have $q \geq 1$. Direct calculation shows that

$$|\partial\phi_A(z)| + |\bar{\partial}\phi_A(z)| = \begin{cases} 1 & |z| > R \\ R^{1-q} |z|^{q-1} \frac{|q+1+i\alpha|+|q-1+i\alpha|}{2} & r \leq |z| \leq R \\ R^{1-q} r^{q-1} & |z| < r \end{cases}$$

and also that

$$J(z, \phi_A) = \begin{cases} 1 & |z| > R \\ q \left(\frac{|z|}{R} \right)^{2(q-1)} & r \leq |z| \leq R \\ \left(\frac{r}{R} \right)^{2(q-1)} & |z| < r \end{cases}$$

whence

$$\mathbb{K}(z, \phi_A) = \begin{cases} 1 & |z| > R \\ \frac{(|q+1+i\alpha|+|q-1+i\alpha|)^2}{4q} & r \leq |z| \leq R \\ 1 & |z| < r \end{cases}$$

In particular, if $2 \leq q + 1 < \alpha$ then one may estimate $\|\mathbb{K}(\cdot, \phi_A)\|_\infty \leq \frac{4\alpha^2}{q}$. Next, let us construct the sequence of maps f_n in an iterative way as follows. For $n = 1$, we set

$$f_1(z) = \phi_{A_1}(z) = \begin{cases} z & |z| < R_1 \\ z \left| \frac{z}{R_1} \right|^{q_1-1} e^{i\alpha_1 \log \frac{|z|}{R_1}} & r_1 \leq |z| \leq R_1 \\ z \left(\frac{1}{e} \right)^{q_1-1} e^{-i\alpha_1} & |z| < r_1 \end{cases}$$

where q_1 and α_1 are to be determined later. Next, assuming we have f_1, \dots, f_{n-1} , we define f_n for $n \geq 2$ as:

$$f_n(z) = \phi_{f_{n-1}(A_n)} \circ f_{n-1}(z)$$

Note that $\phi_{f_{n-1}(A_n)}$ is determined by the inner and outer radii of $\phi_{f_{n-1}(A_n)}$ (which are already available since f_1, \dots, f_{n-1} are known) as well as for the parameters q_n and

α_n , which will be determined later. Clearly, each $f_n : \mathbb{C} \rightarrow \mathbb{C}$ is quasiconformal, and conformal outside the annuli $A_i, i \in \{1, \dots, n\}$. Moreover, one can easily show that

$$\mathbb{K}(\cdot, f_n) = \prod_{j=1}^n \mathbb{K}(\cdot, f_{n-j} \circ \phi_{f_{n-j}(A_{n-j+1})}) = \prod_{j=1}^n \mathbb{K}(\cdot, \phi_{A_{n-j+1}})$$

so that $\mathbb{K}(z, f_n) \leq C \frac{\alpha_j^2}{q_j}$ whenever $z \in A_j, j = 1 \dots n$ while $\mathbb{K}(\cdot, f_n) = 1$ otherwise. In a similar way, we can use that $|D\phi_A(z)| \leq C\alpha$ when $z \in A$ (and $|D\phi_A(z)| \leq 1$ at all other points) to obtain that $|Df_n| \leq C\alpha_j$ on $A_j, j = 1 \dots n$, and $|Df_n| \leq 1$ otherwise.

By construction, we have $f_n(z) = f_{n-1}(z)$ whenever $z \notin B(0, R_n)$. Thus $\{f_n\}_n$ converges uniformly to a map $\bar{f}(z)$, that is,

$$\bar{f} = \lim_{n \rightarrow \infty} f_n$$

which is again a homeomorphism by construction. A similar argument to the one before shows that \bar{f} is absolutely continuous on almost every line parallel to the coordinate axis. For almost every fixed $z_0 \neq 0$, there is a neighborhood of z_0 such that the sequence $\{f_n(z)\}_n$ remains constant for n very large and z in that neighborhood. Therefore, the same happens to the sequences $Df_n(z), J(z, f_n)$ and $\mathbb{K}(z, f_n)$, and so their limits are precisely $D\bar{f}(z), J(z, \bar{f})$ and $\mathbb{K}(z, \bar{f})$. Especially, in order to have $D\bar{f} \in L^1_{loc}$ it suffices that

$$\sum_{n=1}^{\infty} |A_n| \alpha_n < +\infty \tag{11}$$

In case this holds true, then \bar{f} is a homeomorphism in $W^{1,1}_{loc}$, and as a consequence its jacobian determinant $J(\cdot, \bar{f}) \in L^1_{loc}$. Moreover, in order to have $\mathbb{K}(\cdot, \bar{f}) \in L^p_{loc}$ one needs to require that

$$\sum_{n=1}^{\infty} |A_n| \frac{\alpha_n^{2p}}{q_n^p} < \infty \tag{12}$$

Again, as it was the case for f , (12) implies (11) when $q_n^{\frac{p}{2p-1}} < \alpha_n$ and so our parameters α_n and q_n need to be chosen according to (12) as well as the purpose of \bar{f} to be optimal for Theorem 1. For this, again as before, we have $\bar{f}(0) = 0, \bar{f}(1) = 1$ and

$$|\arg(\bar{f}(r_n))| \geq \left| \arg\left(\left(\frac{1}{e}\right)^{q_n+i\alpha_n}\right) \right| = |\alpha_n|$$

which motivates us to choose

$$\alpha_n = h(r_n) \left(\log \frac{1}{r_n} \right)^{1/2} r_n^{-\frac{1}{p}} \quad q_n = \log \frac{1}{r_n},$$

where h is any gauge function such that $h(r) \rightarrow 0$ as $r \rightarrow 0$ and the condition $q_n^{\frac{p}{2p-1}} < \alpha_n$ is satisfied. Indeed, with these choices (12) becomes

$$\sum_n h(r_n)^{2p} < \infty$$

which, as before, may always be granted by choosing smaller r_n , if needed. Having (12) fulfilled, our map \tilde{f} is a mapping of finite distortion with $\mathbb{K}(\cdot, \tilde{f}) \in L^p_{loc}$. Also, the resulting map \tilde{f} attains the optimal rotational behavior stated at Theorem 1 modulo the gauge function h which can be chosen to converge to 0 as slowly as desired.

Therefore, Theorem 3 will be proven if we are able to show that \tilde{f} is Hölder from below. Furthermore, we also show that \tilde{f} is Hölder from above, highlighting regularity of our mappings.

To do this, we first observe that the composition of $z \mapsto ze^{i\alpha \log |z|}$ followed by $z \mapsto z|z|^{q-1}$ is precisely $z \mapsto z|z|^{q-1}e^{i\alpha \log |z|}$. This observation suggests us to decompose $\tilde{f} = g \circ f$, where f is essentially the first example in this section (with different choice of α_n) and g is constructed by building blocks (10) with $\alpha = 0$ at each step. Morally, f leaves fixed all circles centered at 0, and only rotates the annuli A_n , while g conveniently stretches each A_n .

For any $p > 1$, the bi-Hölder nature of f has already been proven when $\alpha_n = h(r_n)r_n^{-1/p}$. Hence we can directly use the same proof there after we estimate

$$h(r_n) \left(\log \frac{1}{r_n} \right)^{1/2} r_n^{-\frac{1}{p}} \leq h(r_n)r_n^{-1/(p-\epsilon)}$$

for all small r_n and $\epsilon = (p - 1)/2$. Therefore, it only remains to show that g is bi-Hölder as well. To this end, we first show that g is Hölder from above using the fundamental theorem of calculus.

Let $x, y \in B(0, 1)$ be given. Without loss of generality let us assume that $|y| \geq |x|$ and let w be the point for which $|w| = |x|$ and $\arg(w) = \arg(y)$. Now

$$|g(x) - g(y)| \leq |g(w) - g(x)| + |g(y) - g(w)|, \tag{13}$$

and we will show that both of these are Hölder. First, since g maps circles centered at the origin radially to similar circles with equal or smaller radius (as $q_n \geq 1$), it is clear that

$$|g(x) - g(w)| \leq |x - w| \leq |x - y|.$$

Let us then concentrate of the second part. First we note, that we can without loss of generality assume that y and w are real numbers as g is a radial mapping. From our

construction, we see that the line segments $[r_n, R_n]$, (R_{n+1}, r_n) and $(R_1, 1]$ partition the line segment $(0, 1]$. Furthermore, from (10), it is clear that the differential is bounded from above by 1 in the segments (R_{n+1}, r_n) and $(R_1, 1]$. On the other hand, in segments $[r_n, R_n]$, we can estimate

$$|g'(t)| \leq \log\left(\frac{1}{r_n}\right) \leq \frac{C}{\sqrt{t}}$$

for any $t \in [r_n, R_n]$ with fixed C that does not depend on n or t . This is so because of our choice of q_n . Combining these two estimates, we have

$$|g'(t)| \leq \frac{C}{\sqrt{t}}$$

for any $t \in (0, 1)$. Thus, we can use fundamental theorem of calculus to estimate

$$\begin{aligned} |g(y) - g(w)| &= \int_w^y |g'(t)| dt \\ &\leq \int_w^y \frac{C}{\sqrt{t}} dt \\ &= 2C(\sqrt{y} - \sqrt{w}) \\ &\leq 2C\sqrt{y - w}. \end{aligned}$$

This proves that also the second part in (13) is Hölder, and thus, we obtain

$$|g(y) - g(x)| \leq |g(y) - g(w)| + |g(w) - g(x)| \leq C\sqrt{|y - w|} + \sqrt{|x - w|} \leq 2C\sqrt{|x - y|},$$

which shows g is Hölder from above.

Let us next prove that g is Hölder from below. To this end, given any two points $x, y \in B(0, 1)$, we again without loss of generality assume that $|y| \geq |x|$ and let w be the point for which $|w| = |x|$ and $\arg(w) = \arg(y)$. Now, as g is a radial homeomorphism, it follows that

$$|g(x) - g(y)| \geq \max\{|g(x) - g(w)|, |g(y) - g(w)|\}$$

Moreover,

$$\max\{|x - w|, |y - w|\} \geq \frac{1}{2}|x - y|$$

Therefore, it is enough to show that both $|g(x) - g(w)|$ and $|g(y) - g(w)|$ satisfy Hölder bounds from below. Note that if $x = 0$ then clearly $w = 0$ and we have only the radial part $|g(y) - g(w)|$.

Let us first check the term $|g(x) - g(w)|$. Since g maps radially circles centered at the origin to similar circles, we see that $|g(x) - g(w)|$ gets contracted the same

amount as the modulus $|g(w)|$ is contracted under g . Now we must consider two possibilities, either $x, w \in A_n$ or $x, w \in D_n$ for some n . Let us first assume $x, w \in A_n = B(0, R_n) \setminus B(0, r_n)$ for some n . Here, we impose an additional assumption that

$$r_n < \left(\frac{1}{e}\right)^{q_{n-1}+q_{n-2}+\dots+q_1-(n-1)}, \tag{14}$$

which we can do as the radii r_n can be assumed to decrease as fast as we want. Then, we can estimate

$$\begin{aligned} |g(x)| &= \left(\frac{1}{e}\right)^{q_{n-1}+q_{n-2}+\dots+q_1-(n-1)} \cdot |x| \left(\frac{|x|}{R_n}\right)^{q_n-1} \\ &\geq r_n \cdot |x| \left(\frac{|x|}{R_n}\right)^{q_n-1} \\ &\geq r_n \cdot |x| \left(\frac{1}{e}\right)^{q_n-1} = e \cdot r_n^2 \cdot |x| \end{aligned}$$

for any $x \in A_n$. Therefore,

$$\begin{aligned} |g(x) - g(w)| &\geq e \cdot r_n^2 \cdot |x - w| \\ &\geq C \cdot |x - w|^3 \end{aligned}$$

since $|x - w| < C \cdot r_n$ for some fixed constant $C > 0$ when $x, w \in A_n$.

Next, let $x, w \in D_n = B(0, r_n) \setminus B(0, R_{n+1})$ for some n . Using (14), we get

$$\begin{aligned} |g(x)| &\geq c \left(\frac{1}{e}\right)^{q_{n-1}+q_{n-2}+\dots+q_1-(n-1)} \cdot r_n \cdot |x| \\ &\geq c \cdot r_n^2 \cdot |x|. \end{aligned}$$

Thus, we can use a similar argument as in the previous case to estimate

$$\begin{aligned} |g(x) - g(w)| &\geq c \cdot r_n^2 \cdot |x - w| \\ &\geq c \cdot |x - w|^3 \end{aligned}$$

since $|x - w| < c \cdot r_n$ for some fixed constant $c > 0$ when $x, w \in D_n$.

Since the set $\mathbb{D} \setminus \{0\}$ is partitioned by separated annuli A_n and D_n we have thus proven that $|g(x) - g(w)|$ satisfies Hölder estimates from below.

Finally, let us prove the Hölder estimates from below for the term $|g(y) - g(w)|$. As the mapping g is radial, we can again assume that y and w are real. We aim to use again the Fundamental Theorem of Calculus, and thus have to estimate the differential from below. Using (14), as well as the fact that $q_n > 1$, we can estimate for any real

number $t \in [r_n, R_n]$ that

$$\begin{aligned} g'(t) &= \left(\frac{1}{e}\right)^{q_{n-1}+q_{n-2}+\dots+q_1-(n-1)} \cdot q_n \cdot \left(\frac{t}{R_n}\right)^{q_n-1} \\ &\geq r_n q_n \cdot \left(\frac{r_n}{R_n}\right)^{q_n-1} \\ &= e q_n r_n^2 \\ &\geq c \cdot t^2 \log \frac{1}{t}. \end{aligned}$$

Next, if $t \in [R_{n+1}, r_n]$, we have

$$\begin{aligned} g'(t) &= \left(\frac{1}{e}\right)^{q_{n-1}+q_{n-2}+\dots+q_1-(n-1)} \cdot \left(\frac{1}{e}\right)^{q_n-1} \\ &\geq e \cdot r_n^2 \\ &\geq c \cdot t^2 \end{aligned}$$

Thus, as before, since $(0, 1)$ is partitioned by the intervals $[r_n, R_n]$, $[R_{n+1}, r_n]$ and $[R_1, 1)$, we end up getting that

$$g'(t) \geq c \cdot t^2$$

for every $t \in (0, 1)$. Now, we use the fundamental theorem of calculus to get

$$\begin{aligned} |g(y) - g(w)| &= \int_w^y g'(t) dt \\ &\geq \int_w^y c \cdot t^2 dt \\ &= C (y^3 - w^3) \\ &\geq C|y - w|^3 \end{aligned}$$

This proves that the second term is Hölder from below as well, which in turn proves that g is Hölder from below. This finishes the proof of Theorem 3.

Acknowledgements A.C. and B.S. are partially supported by projects MTM2016-81703-ERC, MTM2016-75390 (Spanish Government) and 2017SGR395 (Catalan Government). L.H. was partially supported by ICMAT Severo Ochoa project SEV-2015-0554 grant MTM2017-85934-C3-2-P, the ERC grant 307179-GFTIPFD, the ERC grant 834728 Quamap, by the Finnish Academy of Science project 13316965 and by a grant from The Emil Aaltonen Foundation.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence,

and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Astala, K., Iwaniec, T., Martin, G.: Elliptic Partial Differential Equations and Quasiconformal Mappings in the Plane, Princeton Mathematical Series, vol. 48. Princeton University Press, Princeton, NJ (2009)
2. Astala, K., Iwaniec, T., Prause, I., Saksman, E.: Bilipschitz and quasiconformal rotation, stretching and multifractal spectra. *Publ. Math.* **121**(1), 113–154 (2015)
3. Bahouri, H., Chemin, J.Y.: Equations de transport relatives à des champs de vecteurs non-lipschitziens et mécanique des fluides. *Arch. Ration. Mech. Anal.* **127**, 159–181 (1994)
4. Bongers, T.: Stretching and rotation sets of quasiconformal mappings. *Ann. Acad. Sci. Fenn. Math.* **44**, 103–123 (2019)
5. Choi, K., Jeong I.: On the winding number for particle trajectories in a disk-like vortex patch of the Euler equations. <http://arxiv.org/abs/2008.05085v2>
6. Clop, A., Jylhä, H.: Sobolev regular flows of non-Lipschitz vector fields. *J. Differ. Equ.* **266**(8), 4544–4567 (2019)
7. Hitruhin, L.: On multifractal spectrum of quasiconformal mappings. *Ann. Acad. Sci. Fenn. Math.* **41**, 503–522 (2016)
8. Hitruhin, L.: Pointwise rotation for mappings with exponentially integrable distortion. *Proc. Am. Math. Soc.* **144**, 5183–5195 (2016)
9. Hitruhin, L.: Rotational properties of homeomorphisms with integrable distortion. *Conform. Geom. Dyn.* **22**, 78–98 (2018)
10. Hitruhin, L.: Joint rotational and stretching multifractal spectra of mappings with integrable distortion. *Rev. Mat. Iberoam.* **35**(6), 1649–1675 (2019)
11. Iwaniec, T., Martin, G.: Geometric Function Theory and Non-linear Analysis. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York (2001)
12. Koskela, P., Takkinen, J.: Mappings of finite distortion: formation of cusps III. *Acta Math. Sin. English Ser.* **26**, 817–824 (2010)
13. Vuorinen, M.: Conformal Geometry and Quasiregular Mappings, Lecture Notes in Math, vol. 1319. Springer, Berlin (1988)
14. Wolibner, W.: Un théorème d'existence du mouvement plan d'un fluide parfait, homogène, incompressible, pendant un temps infiniment long. *Math. Z.* **37**, 698–726 (1933)
15. Yudovich, V.: Non stationary flow of an ideal and incompressible liquid. *Zh. Vych. Math.* **3**, 1032–1066 (1963)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.