
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.

Teppo, Konsta; Jaakkola, Jussi; Biancari, Fausto; Halminen, Olli; Linna, Miika; Haukka, Jari; Putaala, Jukka; Tiili, Paula; Lehtonen, Ossi; Niemi, Mikko; Mustonen, Pirjo; Kinnunen, Janne; Hartikainen, Juha; Airaksinen, K. E. Juhani; Lehto, Mika

Association of income and educational levels with adherence to direct oral anticoagulant therapy in patients with incident atrial fibrillation

Published in:
Pharmacology Research and Perspectives

DOI:
[10.1002/prp2.961](https://doi.org/10.1002/prp2.961)

Published: 01/06/2022

Document Version
Publisher's PDF, also known as Version of record


Published under the following license:
CC BY-NC

Please cite the original version:
Teppo, K., Jaakkola, J., Biancari, F., Halminen, O., Linna, M., Haukka, J., Putaala, J., Tiili, P., Lehtonen, O., Niemi, M., Mustonen, P., Kinnunen, J., Hartikainen, J., Airaksinen, K. E. J., & Lehto, M. (2022). Association of income and educational levels with adherence to direct oral anticoagulant therapy in patients with incident atrial fibrillation: A Finnish nationwide cohort study. *Pharmacology Research and Perspectives*, 10(3), 1-11. Article e00961. <https://doi.org/10.1002/prp2.961>

This material is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

ORIGINAL ARTICLE

Association of income and educational levels with adherence to direct oral anticoagulant therapy in patients with incident atrial fibrillation: A Finnish nationwide cohort study

Konsta Teppo¹  | Jussi Jaakkola^{1,2} | Fausto Biancari^{3,4} | Olli Halminen⁵ |
 Miika Linna^{6,7} | Jari Haukka⁸ | Jukka Putaala⁹ | Paula Tiili⁸ | Ossi Lehtonen⁶ |
 Mikko Niemi⁸ | Pirjo Mustonen¹⁰ | Janne Kinnunen⁹ | Juha Hartikainen^{7,11} |
 K. E. Juhani Airaksinen^{1,10} | Mika Lehto^{3,8,12}

¹University of Turku, Turku, Finland

²Heart Unit, Satakunta Central Hospital, Pori, Finland

³Heart and Lung Center, Helsinki University Hospital, Helsinki, Finland

⁴Clinica Montevergine, GVM Care & Research, Mercogliano, Italy

⁵Department of Industrial Engineering and Management, Aalto University, Espoo, Finland

⁶Aalto University, Espoo, Finland

⁷University of Eastern Finland, Kuopio, Finland

⁸University of Helsinki, Helsinki, Finland

⁹Neurology, Helsinki University Hospital, and University of Helsinki, Helsinki, Finland

¹⁰Heart Center, Turku University Hospital, Turku, Finland

¹¹Heart Center, Kuopio University Hospital, Kuopio, Finland

¹²Department of Internal Medicine, Lohja Hospital, Lohja, Finland

Correspondence

Konsta Teppo, University of Turku, Turku, 20500, Finland.
 Email: jkitep@utu.fi

Funding information

This work was supported by the Aarne Koskelo Foundation, The Finnish Foundation for Cardiovascular Research, and Helsinki and Uusimaa Hospital District research fund (TYH2019309). *Role of the Funder/Sponsor:* The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Abstract

Low socioeconomic status has been associated with poor outcomes in patients with atrial fibrillation (AF). However, little is known about socioeconomic disparities in adherence to stroke prevention with direct oral anticoagulants (DOACs). We assessed the hypothesis that AF patients with higher income or educational levels have better adherence to DOACs in terms of treatment implementation and persistence. The used nationwide registry-based FinACAF cohort covers all patients with incident AF starting DOACs in Finland during 2011–2018. The implementation analyses included 74 222 (mean age 72.7 ± 10.5 years, 50.8% female) patients, and persistence analyses included 67 503 (mean age 75.3 ± 8.9 years, 53.6% female) patients with indication for permanent anticoagulation (CHA₂DS₂-VASc score >1 in men and >2 in women). Patients were divided into income quartiles and into three categories based on their educational attainment. Therapy

Abbreviations: AF, atrial fibrillation; DOAC, direct oral anticoagulant; FinACAF, Finnish anticoagulation in atrial fibrillation; ICD-10, International Classification of Diseases, Tenth Revision; ISCED, International Standard Classification of Education; MPR, medication possession ratio; OAC, oral anticoagulant; VKA, vitamin-K anticoagulant.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Pharmacology Research & Perspectives* published by John Wiley & Sons Ltd, British Pharmacological Society and American Society for Pharmacology and Experimental Therapeutics.

implementation was measured using the medication possession ratio (MPR), and patients with MPR ≥ 0.90 were defined adherent. Persistence was measured as the incidence of therapy discontinuation, defined as the first 135-day period without DOAC purchases after drug initiation. Patients with higher income or education were consistently more likely adherent to DOACs in the implementation phase (comparing the highest income or educational category to the lowest: adjusted odds ratios 1.18 (1.12–1.25) and 1.21 (1.15–1.27), respectively). No association with income or educational levels was observed on the incidence of therapy discontinuation. In conclusion, we observed that income and educational levels both have independent positive association on the implementation of DOAC therapy but no association on therapy persistence in patients with AF.

KEYWORDS

adherence, atrial fibrillation, direct oral anticoagulants, educational level, income, persistence, socioeconomic status

1 | INTRODUCTION

Atrial fibrillation (AF), the most common cardiac arrhythmia, is associated with a five-fold increased risk of ischemic stroke as well as with stroke recurrence and mortality.^{1,2} Fortunately, the adequate use of oral anticoagulant therapy (OAC) can effectively reduce the risk of ischemic stroke and death.³ Current guidelines recommend the use of direct oral anticoagulants (DOACs) over vitamin-K anticoagulants (VKAs) as the first line anticoagulant due to their superior efficacy and safety profile.⁴ Unlike VKAs, DOACs do not require regular dose monitoring, and the lack of these systematic check-ups has raised concerns about patients' sufficient adherence to DOAC therapy.

Adherence research guidelines recommend dividing medication adherence into three phases: initiation, implementation, and persistence, with implementation referring to how patient's actual dosing corresponds to the prescribed dosing between treatment initiation and discontinuation, and persistence to the length of time between initiation and discontinuation.⁵ All aspects of adherence are crucial for effective stroke prevention with OACs in patients with AF since poor therapy implementation and persistence have been associated with higher mortality and ischemic stroke risk.^{6–8}

Socioeconomic inequalities in health are a major challenge for public health, and their magnitude is affected by differences in health care financing mechanisms.^{9–13} Finland has a public tax-funded health care with universal access, full coverage of public health insurance, and relatively high reimbursement rates of medical treatment.^{14–16} Nevertheless, socioeconomic disparities are observed in Finland in terms of mortality, morbidity, and self-rated health.¹⁷

Previous literature has indicated poor outcomes in patients with AF and low socioeconomic status, and differences in the use of OAC therapy may be underlying these outcome disparities.^{18,19} Indeed, previous studies have suggested that low income and educational levels are associated with lower rate of OAC therapy initiation in patients with AF. However, information on whether socioeconomic factors affect adherence to initiated OAC therapy is limited, especially regarding DOACs.²⁰ The present nationwide cohort study

covering all AF patients in Finland aimed to assess the impact of patients' income and educational levels on the adherence to DOACs focusing on therapy implementation and persistence.

2 | MATERIALS AND METHODS

2.1 | Study population

The Finnish AntiCoagulation in Atrial Fibrillation (FinACAF) Study (ClinicalTrials Identifier: NCT04645537; ENCePP Identifier: EUPAS29845) is a retrospective nationwide cohort study covering all patients with diagnosed AF in Finland during 2004–2018.²¹ Patients were identified from all available national health care registers (hospitalizations and outpatient specialist visits: HILMO; primary health care: AvoHILMO; and National Reimbursement Register upheld by Social Insurance Institute: KELA). The inclusion criterion for the cohort was an International Classification of Diseases, Tenth Revision (ICD-10) diagnosis code I48 (including atrial fibrillation and atrial flutter, together referred as AF) recorded between 2004 and 2018 and FinACAF cohort entry occurred at the date of the first recorded AF diagnosis. The exclusion criteria were permanent migration abroad before December 31st 2018 and age < 18 years at AF diagnosis. Follow-up continued until death or 31st December 2018, whichever occurred first. The current substudy was conducted within a cohort of patients with incident AF between 2007 and 2018, established in previous studies of the FinACAF cohort.^{22,23}

Patients not receiving DOACs during 2011–2018, when DOACs have been approved for stroke prevention in patients with AF, were excluded. Thereafter, we established two separate study cohorts for the analyses of implementation and persistence. In the implementation study cohort, we included only patients with more than one DOAC purchase since true implementation pattern of a long-term preventive therapy cannot be meaningfully determined from only one purchase. In the persistence study cohort, we included patients recommended to receive permanent OAC therapy according to the

contemporary guidelines of the study period, and therefore, females with CHA₂DS₂-VASC score ≤2 and males with CHA₂DS₂-VASC score ≤1 were excluded.^{24,25} In this substudy, follow-up started on the date of the first DOAC purchase. The patient selection process is summarized in the Supplementary Figure 1.

2.2 | Income and educational levels

The patient's highest annual taxable income (in 1000-euro accuracy) during the FinACAF study's observation period 2004–2018 was obtained from the national Tax Register. The annual income was capped to a maximum of 100 000 euros to avoid patients' identifiability due to high incomes. To account for changes in income over time and age, patients were divided into age group and AF diagnosis year specific income quartiles, i.e., each 10-year age group during each cohort entry year was divided into income quartiles using age group and entry year specific cut-points.²⁶ Divisions to income quartiles were performed separately in the implementation and persistence study cohorts.

The patients' highest achieved educational level categorized according to the International Standard Classification of Education (ISCED) was obtained from the Statistics Finland.²⁷ Educational level was divided into three categories: Category 1: ISCED 0–2 (preprimary, primary, and lower secondary education); Category 2: ISCED 3 (Upper secondary or vocational education); Category 3: ISCED 5–8 (tertiary, Bachelor's-level, Master's-level, or doctoral level education). ISCED category 4 does not exist in Finland.

2.3 | Adherence to DOACs

The present substudy focused on the implementation and persistence of initiated DOAC (apixaban, dabigatran, edoxaban, or rivaroxaban) therapy. The commonly used medication possession ratio (MPR) was used to quantify therapy implementation. The MPR of each patient was calculated by dividing the number of days covered with the sum of purchased daily doses during persistent therapy by the number of days between the first and the last DOAC purchase dates added with the days covered with the dose of the last purchase:

$$MPR = \frac{\text{Days covered with the sum of daily doses}}{\text{Days between first and last DOAC purchase plus the days covered with the daily dose from last DOAC purchase}}$$

Medication possession ratio values were capped to a maximum of 1.0 and patients with MPR ≥0.90 were defined adherent, since MPR <0.90 has been associated with reduced efficacy of stroke prevention with DOACs.^{6,28} We assessed treatment implementation during persistent DOAC use, i.e., between treatment initiation and discontinuation, and only DOAC purchases before therapy discontinuation were included in the MPR calculations. Discontinuation was defined as the first 135-day period without DOAC redemptions. The 135-day definition was chosen, since in Finland it is possible to purchase drugs with

reimbursement for a maximum of 90 days and an additional 45-day grace period was allowed. Additionally, sensitivity analyses were performed including all DOAC purchases during entire follow-up instead of only drug purchases before the first discontinuation event.

In the persistence analyses, we determined the incidence of therapy discontinuation, i.e., non-persistence. The date of the first DOAC purchase was the index date and outcome was the first therapy discontinuation event. As stated above, discontinuation event was defined as the first 135-day period without DOAC redemptions and was considered to occur at the end of the 135-day period. Individuals switching to VKA during the 135-day period were censored, and those switching from one DOAC to another during the 135-day period were considered persistent. As a sensitivity analysis, we analyzed the rate of DOAC therapy cessation, considered to occur on the date of the last DOAC purchase in patients with at least six months of follow-up after the last purchase to ensure more definitive termination of stroke prevention. Additionally, patients initiating VKA after the last DOAC purchase were censored in the cessation analysis.

2.4 | Statistical analysis

Statistical analyses were performed with the IBM SPSS Statistics software (version 27.0, SPSS, Inc., Armonk, NY) and R (version 4.0.5, <https://www.R-project.org>). The chi-square test was used to compare differences between proportions, and the one-way analysis of variance to analyze continuous variables. MPR was non-normally distributed, and therefore, mean MPR between socioeconomic groups was compared using the Kruskal–Wallis test. Unadjusted and adjusted odds ratios (ORs) of adherence to DOAC therapy (MPR ≥0.90) for socioeconomic categories were calculated using the binary logistic regression. Poisson regression was used to estimate the crude incidence as well as the unadjusted and adjusted incidence rate ratios (IRRs) of therapy discontinuation for income and educational categories. The observation of discontinuation event may be hindered by mortality occurring during study period, and therefore, the Fine-Gray regression with all-cause death as competing event was used to estimate the unadjusted and adjusted subdistribution hazard ratios (SHRs) of

DOAC therapy discontinuation for income and educational categories. In addition to income and educational levels, the analyses with the Fine-Gray and binary logistic regression models were adjusted for age (categorical variable in 10 year groups), sex, calendar year of DOAC initiation, stroke and bleeding risk factors (heart failure, hypertension, diabetes, prior stroke or transient ischemic attack, vascular disease, prior bleeding, alcohol abuse, renal failure and liver cirrhosis or failure, concomitant use of NSAIDs or antiplatelets), dementia, mental health conditions, dosage of the first

purchased DOAC (once or twice daily), previous use of VKAs and polypharmacy (>5 different medications during the year preceding DOAC initiation), since these factors have been shown to affect medication adherence in previous studies.²⁹⁻³³ The definitions of the comorbidities are displayed in Supplementary Table 1.

3 | RESULTS

Altogether, 74 222 patients (mean age 72.7 (SD 10.5) years, 50.8% female) were included in the implementation study cohort and 67 503 patients (mean age 75.3 (SD 8.9) years, 53.6% female) in the persistence study cohort. In both cohorts, patients with higher income were more likely male, had higher education and lower prevalence of cardiovascular comorbidities, dementia, and alcohol use disorders. Similar trends were observed in patients with higher educational level, and additionally, patients with higher education were younger than patients in the lowest educational category (Table 1). Mean duration of persistent DOAC therapy during follow up was 1.4 (SD 1.1) years in the implementation study cohort and 1.3 (SD 1.1) years in the persistence study cohort.

3.1 | Implementation of DOAC therapy

Overall, mean MPR was 0.89 (SD 0.17) and 49 950 (67.3%) patients were adherent to DOACs (MPR \geq 0.90) during persistent therapy. The mean MPR, proportion of adherent patients as well as the unadjusted and adjusted odds for adherent DOAC use were all consistently higher among patients in higher income and educational levels (Table 2). The findings were reiterated in the sensitivity analyses covering all DOAC purchases during follow-up (Supplementary Table 2).

3.2 | Persistence of DOAC therapy

A total of 11 856 (17.6%) patients discontinued DOAC therapy during follow-up. Persistence of DOAC use reduced substantially over time (Figure 1). The proportion of patients without therapy discontinuation event still in follow-up at one and two years after DOAC initiation were 81.7% and 69.7%, respectively. Inconsistent differences between income and educational categories were observed in the proportions of patients discontinuing DOAC therapy during entire follow-up, with the highest proportion of discontinued therapies in the highest income and educational categories (Table 3). However, no disparities among income and educational groups were observed in the unadjusted or adjusted incidence rates of therapy discontinuation in the Poisson and Fine-Gray regression models, except for the marginally lower discontinuation rate among patients in the 3rd income quartile, when compared to the lowest income quartile (Table 3). Similarly, in the sensitivity analyses, the overall adjusted cessation rate of DOAC therapy did not differ between income or education categories (Supplementary Table 3).

4 | DISCUSSION

This nationwide cohort study based on pharmacy claims data demonstrated that both income and educational level are independently associated with better implementation of initiated DOAC therapy in patients with incident AF. In contrast, no associations with income or education were observed on the persistence of DOAC therapy.

Previous research assessing the relationship of socioeconomic factors and adherence to DOAC therapy in patients with AF is limited and has demonstrated inconsistent results.^{31,34-37} Importantly, most prior works have focused only on a single aspect in medication adherence rather than encompassing both implementation and persistence dimensions of drug utilization. Additionally, these studies have been prone to possible selection, information, and confounding biases owing to use of area-based socioeconomic data, patient samples from only a single level of care, and lack of controlling for possible mortality differences and other confounding factors. Therefore, the results of this large nationwide study covering all patients with AF in Finland considerably increase our understanding of the impact of income and educational attainment on the overall DOAC therapy adherence.

Patients in the highest income and educational categories had 18%–21% higher adjusted odds of sufficient adherence to DOACs (MPR \geq 0.90) in the therapy implementation phase than patients in the lowest categories, a finding in line with previous reports associating higher socioeconomic status with better therapy implementation in other chronic conditions, although studies on DOAC use in patients with AF have been inconclusive.³⁷⁻⁴⁰ On the other hand, no meaningful difference in persistence of DOAC therapy was observed among income and educational levels, in concordance with a number of observations of similar medication persistence in different socioeconomic categories among patients with and without AF, while reports on worse medication persistence among patients with low socioeconomic status can also be found from previous literature.^{34,41-45} However, the large heterogeneity in adherence measures and definitions as well as in the used socioeconomic variables hampers the generalizability and comparability of the results from previous studies. Lower utilization of OAC therapy has been proposed as an underlying mechanism in the observed worse outcomes in patients with AF and low socioeconomic status.¹⁸ Indeed, our findings suggest that inferior implementation of DOAC therapy may contribute to the higher risks among these patients, while differences in treatment persistence are unlikely to play a substantial role in the outcome disparities.

The observed socioeconomic disparities in treatment implementation are likely multifactorial. Importantly, although the costs of DOACs have been largely reimbursed to patients with AF at risk of stroke since 2012 in Finland, DOACs are still significantly more expensive than VKAs, likely hindering their use in patients with low income. Lower levels of health literacy and trust between patients and clinicians may impair understanding of the importance of stroke prevention among patients with lower income or educational background. Additionally, the higher prevalence of mental health

TABLE 1 Descriptive characteristics of the of the study cohorts

| | Income quartiles | | | | p-value | Educational categories | | | p-value |
|--|----------------------------|-------------------|-------------------|-----------------------------|---------------|-----------------------------|-------------------|-----------------------------|---------|
| | 1st (lowest) n = 18 455 | 2nd n = 18 703 | 3rd n = 18 533 | 4th (highest) n = 18 531 | | s1st (lowest) n = 34 576 | 2nd n = 21 559 | 3rd (highest) n = 18 087 | |
| Implementation study cohort | | | | | | | | | |
| Mean annual income (thousands of euros) | 3.9 (5.0) | 13.9 (8.4) | 25.7 (10.8) | 58.7 (24.7) | <.001 | 16.0 (18.6) | 24.1 (21.4) | 45.6 (28.1) | <.001 |
| Demographics | | | | | | | | | |
| Mean age, years | 73.3 (10.4) | 73.0 (10.5) | 72.4 (10.5) | 72.4 (10.4) | <.001 | 76.2 (9.4) | 69.3 (10.6) | 70.2 (10.2) | <.001 |
| Female sex | 12 257 (66.4) | 10 902 (58.3) | 8 604 (46.4) | 5 944 (32.1) | <.001 | 19 593 (56.7) | 10 523 (48.8) | 7 591 (42.0) | <.001 |
| Educational categories | | | | | | | | | |
| 1st | 11 573 (62.7) | 10 417 (55.7) | 7 976 (43.0) | <.001 | N/A | N/A | N/A | N/A | <.001 |
| 2nd | 5767 (31.2) | 6119 (32.7) | 5788 (31.2) | 3885 (21.0) | N/A | N/A | N/A | N/A | <.001 |
| 3rd | 1115 (6.0) | 2167 (11.6) | 4769 (25.7) | 10 036 (54.2) | N/A | N/A | N/A | N/A | <.001 |
| Income quartiles | | | | | | | | | |
| 1st | N/A | N/A | N/A | N/A | 11 573 (33.5) | 5 767 (26.7) | 1 115 (6.2) | 1 115 (6.2) | <.001 |
| 2nd | N/A | N/A | N/A | N/A | 10 417 (30.1) | 6 119 (28.4) | 2 167 (12.0) | 2 167 (12.0) | <.001 |
| 3rd | N/A | N/A | N/A | N/A | 7 976 (23.1) | 5 788 (26.8) | 4 769 (26.4) | 4 769 (26.4) | <.001 |
| 4th | N/A | N/A | N/A | N/A | 4 610 (13.3) | 3 885 (18.0) | 10 036 (55.5) | 10 036 (55.5) | <.001 |
| Comorbidities and risk scores | | | | | | | | | |
| Abnormal liver function | 55 (0.3) | 45 (0.2) | 47 (0.3) | 43 (0.2) | .60 | 80 (0.2) | 60 (0.3) | 50 (0.3) | .46 |
| Abnormal renal function | 546 (3.0) | 531 (2.8) | 483 (2.6) | 470 (2.5) | .04 | 1 135 (3.3) | 504 (2.3) | 391 (2.2) | <.001 |
| Alcohol use disorder | 1008 (5.5) | 559 (3.0) | 500 (2.7) | 419 (2.3) | <.001 | 1088 (3.1) | 879 (4.1) | 519 (2.9) | <.001 |
| Any vascular disease | 5260 (28.5) | 5072 (27.1) | 4699 (25.4) | 4434 (23.9) | <.001 | 10 593 (30.6) | 5241 (24.3) | 3631 (20.1) | <.001 |
| Dementia | 722 (3.9) | 557 (3.0) | 494 (2.7) | 435 (2.3) | <.001 | 1479 (4.3) | 428 (2.0) | 301 (1.7) | <.001 |
| Diabetes | 5064 (27.4) | 4575 (24.5) | 4226 (22.8) | 3827 (20.7) | <.001 | 8985 (26.9) | 5157 (23.9) | 3550 (19.6) | <.001 |
| Heart failure | 2978 (16.1) | 2425 (13.0) | 2051 (11.1) | 1749 (9.4) | <.001 | 5291 (15.3) | 2477 (11.5) | 1435 (7.9) | <.001 |
| Hypertension | 15 039 (81.5) | 15 030 (80.4) | 14 646 (79.0) | 14 189 (76.6) | <.001 | 28 436 (82.2) | 16 806 (78.0) | 13 662 (75.5) | <.001 |
| Prior bleeding | 1773 (9.6) | 1757 (9.4) | 1813 (9.8) | 1739 (9.4) | .51 | 3583 (10.4) | 1979 (9.2) | 1520 (8.4) | <.001 |
| Prior ischemic stroke or TIA | 3079 (16.7) | 2817 (15.1) | 2728 (14.7) | 2584 (13.9) | <.001 | 5718 (16.5) | 3109 (14.4) | 2381 (13.2) | <.001 |
| Prior myocardial infarction | 1537 (8.3) | 1447 (7.7) | 1398 (7.5) | 1242 (6.7) | <.001 | 3034 (8.8) | 1528 (7.1) | 1062 (5.9) | <.001 |
| Psychiatric disorder | 3516 (19.1) | 2527 (13.5) | 2117 (11.4) | 1715 (9.3) | <.001 | 4757 (13.8) | 3150 (14.6) | 1968 (10.9) | <.001 |

(Continues)

TABLE 1 (Continued)

| | Income quartiles | | | | p-value | Educational categories | | | | p-value |
|--|----------------------------|-------------------|-------------------|-----------------------------|---------|----------------------------|-------------------|-----------------------------|-------|---------|
| | 1st (lowest) n = 18 455 | 2nd n = 18 703 | 3rd n = 18 533 | 4th (highest) n = 18 531 | | 1st (lowest) n = 34 576 | 2nd n = 21 559 | 3rd (highest) n = 18 087 | | |
| CHA ₂ DS ₂ -VAsC score | 3.8 (1.7) | 3.6 (1.7) | 3.3 (1.7) | 3.1 (1.7) | <.001 | 3.9 (1.7) | 3.1 (1.7) | 2.9 (1.6) | <.001 | |
| Modified HAS-BLED score | 2.7 (1.0) | 2.6 (0.9) | 2.6 (1.0) | 2.6 (1.9) | <.001 | 2.8 (0.9) | 2.5 (1.0) | 2.5 (1.0) | <.001 | |
| Persistence study cohort | | | | | | | | | | |
| Income quartiles | | | | | | | | | | |
| Mean annual income (thousands of euros) | 2.8 (3.6) | 11.6 (6.7) | 22.8 (9.3) | 55.3 (24.3) | <.001 | 14.9 (17.8) | 21.7 (20.9) | 43.5 (28.0) | <.001 | |
| Demographics | | | | | | | | | | |
| Mean age, years | 75.8 (8.9) | 75.6 (8.9) | 75.0 (8.9) | 75.0 (8.9) | <.001 | 77.7 (8.3) | 72.5 (9.2) | 73.4 (8.5) | <.001 | |
| Female sex | 11 739 (69.5) | 10 209 (60.4) | 8315 (49.4) | 5897 (35.0) | <.001 | 20 082 (58.5) | 9494 (52.3) | 6584 (43.7) | <.001 | |
| Educational categories | | | | | | | | | | |
| 1st | 11 279 (66.8) | 10 278 (60.8) | 8141 (48.3) | 4609 (27.3) | <.001 | N/A | N/A | N/A | <.001 | |
| 2nd | 4763 (28.2) | 4986 (29.5) | 4940 (29.3) | 3451 (20.5) | | N/A | N/A | N/A | | |
| 3rd | 840 (5.0) | 1651 (9.8) | 3764 (22.3) | 8801 (52.2) | | N/A | N/A | N/A | | |
| Income quartiles | | | | | | | | | | |
| 1st | N/A | N/A | N/A | N/A | | 11 279 (32.9) | 4763 (26.3) | 840 (5.6) | | |
| 2nd | N/A | N/A | N/A | N/A | | 10 278 (30.0) | 4986 (27.5) | 1651 (11.0) | | |
| 3rd | N/A | N/A | N/A | N/A | | 8141 (23.7) | 4940 (27.2) | 3764 (25.0) | | |
| 4th | N/A | N/A | N/A | N/A | | 4609 (13.4) | 3451 (19.0) | 8801 (58.5) | | |
| Comorbidities and risk scores | | | | | | | | | | |
| Abnormal liver function | 50 (0.3) | 43 (0.3) | 45 (0.3) | 41 (0.2) | .80 | 81 (0.2) | 52 (0.3) | 46 (0.3) | .31 | |
| Abnormal renal function | 564 (3.3) | 575 (3.4) | 536 (3.2) | 523 (3.1) | .38 | 1 237 (3.6) | 545 (3.0) | 416 (2.8) | <.001 | |
| Alcohol use disorder | 757 (4.5) | 502 (3.0) | 448 (2.7) | 379 (2.2) | <.001 | 963 (2.8) | 677 (3.7) | 446 (3.0) | <.001 | |
| Any vascular disease | 5513 (32.7) | 5333 (31.5) | 5144 (30.5) | 4801 (28.5) | <.001 | 11 496 (33.5) | 5495 (30.3) | 3800 (25.2) | <.001 | |
| Dementia | 783 (4.6) | 610 (3.6) | 540 (3.2) | 499 (3.0) | <.001 | 1636 (4.8) | 457 (2.5) | 339 (2.3) | <.001 | |
| Diabetes | 5256 (31.1) | 4910 (29.0) | 4697 (27.9) | 4254 (25.2) | <.001 | 9816 (28.6) | 5521 (30.4) | 3780 (25.1) | <.001 | |

TABLE 1 (Continued)

| | Income quartiles | | | | Educational categories | | | | p-value |
|--|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------|---------|
| | 1st (lowest) | 2nd | 3rd | 4th (highest) | 1st (lowest) | 2nd | 3rd (highest) | p-value | |
| Heart failure | n = 16 882 3080 (18.2) | n = 16 915 2610 (15.4) | n = 16 845 2229 (13.2) | n = 16 861 1934 (11.5) | n = 34 307 5768 (16.8) | n = 18 140 2580 (14.2) | n = 15 056 1505 (10.0) | <.001 | |
| Hypertension | 14 669 (86.9) | 14 677 (86.8) | 14 509 (86.1) | 14 176 (84.1) | 29 634 (86.4) | 15 706 (86.6) | 12 691 (84.3) | <.001 | |
| Prior bleeding | 1770 (10.5) | 1757 (10.4) | 1844 (10.9) | 1802 (10.7) | 3779 (11.0) | 1918 (10.6) | 1476 (9.8) | <.001 | |
| Prior ischemic stroke or TIA | 3253 (19.3) | 3032 (17.9) | 3035 (18.0) | 2903 (17.2) | 6256 (18.2) | 3373 (18.6) | 2594 (17.2) | .004 | |
| Prior myocardial infarction | 1646 (9.8) | 1563 (9.2) | 1536 (9.1) | 1383 (8.2) | 3357 (9.8) | 1640 (9.0) | 1131 (7.5) | <.001 | |
| Psychiatric disorder | 3043 (18.0) | 2397 (14.2) | 2024 (12.0) | 1679 (10.0) | 4696 (13.7) | 2704 (14.9) | 1743 (11.6) | <.001 | |
| CHA ₂ DS ₂ -VASc score | 4.2 (1.4) | 4.0 (1.4) | 3.8 (1.4) | 3.6 (1.4) | 4.2 (1.5) | 3.7 (1.4) | 3.5 (1.3) | <.001 | |
| Modified HAS-BLED score | 2.9 (0.9) | 2.9 (0.8) | 2.8 (0.8) | 2.8 (0.8) | 2.9 (0.8) | 2.8 (0.9) | 2.8 (0.8) | <.001 | |

Note: Values denote n (%) or mean (standard deviation).

Abbreviations: CHA₂DS₂-VASc, congestive heart failure, hypertension, age ≥75 years, diabetes, history of stroke or TIA, vascular disease, age 65–74 years, sex category (female); modified HAS-BLED score, hypertension, abnormal renal or liver function, prior stroke, bleeding history, age >65 years, alcohol abuse, concomitant antiplatelet/NSAIDs (no labile INR, max score 8); TIA, transient ischemic attack.

conditions, alcohol use disorders and dementia among patients with lower socioeconomic status may affect medication behavior.^{32,33} Yet, these same factors seem not to impact DOAC therapy implementation and persistence to the same extent, since meaningful disparities in medication persistence were not observed.

The main limitations of our study are related to the observational nature of the used administrative data. Hence, the findings reflect associations and not necessarily causation, residual confounding of unmeasured factors cannot be excluded, and information bias may be present due to inaccurate recording of diagnoses and other data. Additionally, some of the used adjusting variables may also have a role as mediators instead of only confounders, especially in the association between educational attainment and adherence. Moreover, our results rely on pharmacy claims and the proportion of drugs truly taken is unknown, and since we lacked data on DOAC prescriptions, we were unable to assess the primary non-adherence to prescribed therapy. Furthermore, clinically indicated treatment gaps are not accounted for in our data, possibly causing downward bias on our adherence estimates. Similarly, we lacked information on the actual patient-level reasons for therapy discontinuation. Moreover, a gold standard to define medication adherence is lacking and there are numerous methods to quantify therapy implementation and persistence, which may influence the results considerably.⁴⁶ Nevertheless, our aim was to compare socioeconomic differences in adherence rather than to calculate absolute adherence estimates, and therefore the results are possibly not materially affected by the chosen methodologies. Indeed, the results of the sensitivity analyses were uniform with the results of the main analyses. Likewise, differences in the used socioeconomic variables, for example household vs. individual income, may lead to varying results in health inequality research.²⁶ However, in our study, income and educational levels both had a similar association on the adherence estimates.

Particular strengths of our study are the large sample size and the comprehensive nature of the used nationwide data, covering all patients with AF in Finland from all levels of care and their individual socioeconomic data as well as all redeemed DOAC prescriptions since DOACs are not sold over the counter without prescription. In addition to the previously reported lower initiation of OAC therapy among AF patients with low socioeconomic status, our results highlight important socioeconomic disparities in the implementation phase of initiated DOAC therapy, and emphasize the need for efforts to ensure adequate quality of stroke prevention for all patients with AF at risk of ischemic stroke.²⁰

In conclusion, in this nationwide observational study based on pharmacy claims data, higher income and educational attainment were both independently associated with better implementation of DOAC therapy in patients with incident AF. However, no meaningful disparities in persistence of DOAC therapy was observed among income or educational levels.

DISCLOSURES

Konsta Teppo: none. Jussi Jaakkola: none. Fausto Biancari: none Olli Halminen: none. Jukka Putaala: Dr. Putaala reports personal fees from

TABLE 2 Adherence to direct oral anticoagulant (DOAC) therapy according to income and educational levels

| | Mean MPR | Proportion of adherent patients (MPR \geq 0.90) | Unadjusted OR (95% CI) | Adjusted OR (95% CI) |
|-------------------------------|----------|---|------------------------|----------------------|
| Income quartiles | | | | |
| 1st (lowest) | 0.878 | 64.8% | (Reference) | (Reference) |
| 2nd | 0.882 | 66.3% | 1.07 (1.03–1.12) | 1.06 (1.01–1.11) |
| 3rd | 0.890 | 68.1% | 1.16 (1.11–1.21) | 1.09 (1.04–1.14) |
| 4th (highest) | 0.897 | 70.0% | 1.27 (1.21–1.32) | 1.18 (1.12–1.25) |
| Educational categories | | | | |
| 1st (lowest) | 0.864 | 61.0% | (Reference) | (Reference) |
| 2nd | 0.904 | 72.0% | 1.64 (1.58–1.70) | 1.12 (1.07–1.16) |
| 3rd (highest) | 0.911 | 73.8% | 1.80 (1.73–1.87) | 1.21 (1.15–1.27) |

Note: ORs estimated with binary logistic regression with the following variables included in adjusted analyses: age, sex, calendar year, heart failure, hypertension, diabetes, prior stroke or TIA, vascular disease, prior bleeding, alcohol abuse, renal failure, liver cirrhosis or failure, concomitant use of NSAIDs or antiplatelets, dementia, psychiatric disorder, DOAC dosing, previous VKA use, polypharmacy, income quartiles and education categories. Abbreviations: CI, confidence interval; MPR, medication possession ratio; OR, odds ratio.

* $p < .001$.




FIGURE 1 Cumulative incidence curve of DOAC therapy discontinuation according to income and educational levels

Boehringer-Ingelheim, personal fees and other from Bayer, grants and personal fees from BMS-Pfizer, personal fees from Portola, other from Amgen, personal fees from Herantis Pharma, personal fees from Terve Media, other from Vital Signum, personal fees from Abbott, outside the submitted work. Pirjo Mustonen: Consultant: Roche, BMS-Pfizer-alliance, Novartis Finland, Boehringer Ingelheim, MSD Finland. Jari Haukka: Consultant: Research Janssen R&D; Speaker: Bayer Finland. Miika Linna: Speaker: BMS-Pfizer-alliance, Bayer, Boehringer-Ingelheim. Juha Hartikainen: Research grants: The Finnish Foundation for Cardiovascular Research, EU Horizon

2020, EU FP7. Advisory Board Member: BMS-Pfizer-alliance, Novo Nordisk, Amgen. Speaker: Cardiome, Bayer. K. E. Juhani Airaksinen: Research grants: The Finnish Foundation for Cardiovascular Research; Speaker: Bayer, Pfizer and Boehringer-Ingelheim. Member in the advisory boards: Bayer, Pfizer and AstraZeneca. Mika Lehto: Consultant: BMS-Pfizer-alliance, Bayer, Boehringer-Ingelheim, and MSD; Speaker: BMS-Pfizer-alliance, Bayer, Boehringer Ingelheim, MSD, Terve Media and Orion Pharma. Research grants: Aarne Koskelo Foundation, The Finnish Foundation for Cardiovascular Research, and Helsinki and Uusimaa Hospital District research fund,

TABLE 3 Incidence of direct oral anticoagulant (DOAC) therapy discontinuation according to income and educational levels

| | Events | Patient years | Incidence (per patient year) | Unadjusted IRR | Adjusted IRR | Unadjusted SHR | Adjusted SHR |
|----------------------|---------------|---------------|------------------------------|------------------|------------------|------------------|------------------|
| Income quartile | | | | | | | |
| 1st (lowest) | 2 940 (17.4%) | 20 863 | 0.14 (0.14–0.15) | (Reference) | (Reference) | (Reference) | (Reference) |
| 2nd | 2 855 (16.9%) | 21 487 | 0.13 (0.13–0.14) | 0.94 (0.90–0.99) | 0.96 (0.91–1.01) | 0.96 (0.91–1.01) | 0.96 (0.91–1.01) |
| 3rd | 2 832 (16.8%) | 21 898 | 0.13 (0.13–0.13) | 0.92 (0.87–0.97) | 0.93 (0.88–0.98) | 0.94 (0.90–0.99) | 0.94 (0.89–0.99) |
| 4th (highest) | 3 229 (19.2%) | 22 566 | 0.14 (0.14–0.15) | 1.02 (0.97–1.07) | 1.01 (0.95–1.07) | 1.06 (1.01–1.11) | 1.01 (0.95–1.07) |
| Educational category | | | | | | | |
| 1st (lowest) | 5 962 (17.4%) | 42 790 | 0.14 (0.14–0.14) | (Reference) | (Reference) | (Reference) | (Reference) |
| 2nd | 3 109 (17.1%) | 23 225 | 0.13 (0.13–0.14) | 0.96 (0.92–1.00) | 1.00 (0.95–1.04) | 0.99 (0.95–1.03) | 1.01 (0.96–1.05) |
| 3rd (highest) | 2 785 (18.5%) | 20 798 | 0.13 (0.13–0.14) | 0.96 (0.92–1.01) | 0.99 (0.94–1.04) | 1.01 (0.97–1.06) | 1.01 (0.96–1.07) |

Note: IRRs estimated by Poisson regression and SHRs by Fine-Gray regression with all-cause death as competing event. Adjusted analyses included the following variables: age, sex, calendar year, heart failure, hypertension, diabetes, prior stroke or TIA, vascular disease, prior bleeding, alcohol abuse, renal failure, liver cirrhosis or failure, concomitant use of NSAIDs or antiplatelets, dementia, psychiatric disorder, DOAC dosing, previous VKA use, polypharmacy, income quartiles and education categories.

Abbreviations: IRR, incidence rate ratio; SHR, subdistribution hazard ratio; 95% confidence intervals in parenthesis.

* $p < .001$.

Boehringer-Ingelheim. Paula Tiili: none. Mikko Niemi: none. Ossi Lehtonen: none.

AUTHOR CONTRIBUTIONS

Dr Teppo had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Concept and design: Teppo, Jaakkola, Putaala, Mustonen, Haukka, Airaksinen, Lehto, Acquisition, analysis, or interpretation of data: All authors. Drafting of the manuscript: Teppo, Critical revision of the manuscript for important intellectual content: All authors. Statistical analysis: Teppo, Jaakkola, Obtained funding: Lehto, Administrative, technical, or material support: Jaakkola, Halminen, Haukka, Supervision: Jaakkola, Putaala, Mustonen, Haukka, Airaksinen, Lehto.

ETHICS STATEMENT

The study protocol was approved by the Ethics Committee of the Medical Faculty of Helsinki University, Helsinki, Finland (nr. 15/2017) and granted research permission from the Helsinki University Hospital (HUS/46/2018). Respective permissions were obtained from the Finnish register holders (KELA 138/522/2018; THL 2101/5.05.00/2018; Population Register Centre VRK/1291/2019-3; Statistics Finland TK-53-1713-18/u1281; and Tax Register VH/874/07.01.03/2019). The patients' identification numbers were pseudonymized, and the research group received individualized, but unidentifiable data. Informed consent was waived due to the retrospective registry nature of the study. The study conforms to the Declaration of Helsinki as revised in 2002.

DATA AVAILABILITY STATEMENT

Because of the sensitive nature of the data collected for this study, requests to access the dataset from qualified researchers trained in human subject confidentiality protocols may be sent to the Finnish national register holders (KELA, Finnish Institute for Health and Welfare, Population Register Center and Tax Register).

ORCID

Konsta Teppo  <https://orcid.org/0000-0002-4460-0994>

REFERENCES

1. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the framingham study. *Stroke*. 1991;22(8):983-988. doi:10.1161/01.STR.22.8.983
2. Dai H, Zhang Q, Much AA, et al. Global, regional, and national prevalence, incidence, mortality, and risk factors for atrial fibrillation, 1990–2017: results from the Global Burden of Disease Study 2017. *Eur Heart J Q Care Clin Outc*. 2021;7(6):574-582. doi:10.1093/ehjqc/co/qcaa061
3. Hart RG, Pearce LA, Meta-analysis AMI. Antithrombotic therapy to prevent stroke in patients who have nonvalvular atrial fibrillation. *Ann Intern Med*. 2007;146(12):857. doi:10.7326/0003-4819-146-12-200706190-00007
4. Hindricks G, Potpara T, Dagres N, et al. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J*. 2021;42(5):373-498. doi:10.1093/eurheartj/ehaa612

5. De Geest S, Zullig LL, Dunbar-Jacob J, et al. ESPACOMP medication adherence reporting guideline (EMERGE). *Ann Intern Med.* 2018;169(1):30. doi:10.7326/M18-0543
6. Komen JJ, Heerdink ER, Klungel OH, et al. Long-term persistence and adherence with non-vitamin K oral anticoagulants in patients with atrial fibrillation and their associations with stroke risk. *Eur Heart J Cardiovasc Pharmacother.* 2021;7(F1):f72-f80. doi:10.1093/ehjcvp/pvaa017
7. Hurtado-Navarro I, García-Sempere A, Rodríguez-Bernal C, Santa-Ana-Tellez Y, Peiró S, Sanfélix-Gimeno G. Estimating adherence based on prescription or dispensation information: impact on thresholds and outcomes. A real-world study with atrial fibrillation patients treated with oral anticoagulants in Spain. *Front Pharmacol.* 2018;9: doi:10.3389/fphar.2018.01353
8. Ozaki AF, Choi AS, Le QT, et al. Real-world adherence and persistence to direct oral anticoagulants in patients with atrial fibrillation. *System Rev Meta-Anal Circul Cardiovasc Q Outcom.* 2020;13(3). doi:10.1161/CIRCOUTCOMES.119.005969
9. Marmot M. Health equity in England: the marmot review 10 years on. *BMJ.* 2020;368:m693. doi:10.1136/bmj.m693
10. Bor J, Cohen GH, Galea S. Population health in an era of rising income inequality: USA, 1980–2015. *The Lancet.* 2017;389(10077):1475-1490. doi:10.1016/S0140-6736(17)30571-8
11. Lueckmann SL, Hoebel J, Roick J, et al. Socioeconomic inequalities in primary-care and specialist physician visits: a systematic review. *Int J for Equity Health.* 2021;20(1): doi:10.1186/s12939-020-01375-1
12. Myint CY, Pavlova M, Thein KNN, Groot W. A systematic review of the health-financing mechanisms in the association of south-east asian nations countries and the people's Republic of China: lessons for the move towards universal health coverage. *PLoS One.* 2019;14(6):e0217278. doi:10.1371/journal.pone.0217278
13. Lagarde M, Palmer N. The impact of health financing strategies on access to health services in low and middle income countries. *Cochrane Database Syst Rev.* 2018;2018(4): doi:10.1002/14651858.CD006092.pub2
14. Kringos D, Boerma W, Hutchinson A, Saltman RB. Building primary care in a changing Europe. European Observatory on Health Systems and Policies; (Observatory Studies Series 38). 2015.
15. Medicines reimbursement policies in Europe. Published online October 5, 2018; 2018. ISBN: 978 92 890 5336 5. Accessed June 6, 2021. <https://www.euro.who.int/en/publications/abstracts/medicines-reimbursement-policies-in-europe>
16. Laugesen K, Ludvigsson JF, Schmidt M, et al. Nordic health registry-based research: a review of health care systems and key registries. *Clin Epidemiol.* 2021;13:533-554. doi:10.2147/CLEP.S314959
17. Karvonen S, Martelin T, Kestilä L, Junna L. Tulotason mukaiset terveyserot ovat edelleen suuria. THL: Published online; 2017.
18. Lunde ED, Nielsen PB, Riahi S, et al. Associations between socioeconomic status, atrial fibrillation, and outcomes: a systematic review. *Expert Rev Cardiovasc Ther.* 2018;16(11):857-873. doi:10.1080/14779072.2018.1533118
19. Ravvaz K, Weissert JA, Jahangir A, Ruff CT. Evaluating the effects of socioeconomic status on stroke and bleeding risk scores and clinical events in patients on oral anticoagulant for new onset atrial fibrillation. *PLoS One.* 2021;16(3):e0248134. doi:10.1371/journal.pone.0248134
20. Lunde ED, Joensen AM, Fonager K, et al. Socioeconomic inequality in oral anticoagulation therapy initiation in patients with atrial fibrillation with high risk of stroke: A register-based observational study. *BMJ Open.* 2021;11(5):e048839. doi:10.1136/bmjopen-2021-048839
21. Lehto M, Halminen O, Mustonen P, et al. The nationwide Finnish anticoagulation in atrial fibrillation (FinACAF): study rationale, design, and patient characteristics. *Eur J Epidemiol.* 2022;37(1):95-102. doi:10.1007/s10654-021-00812-x
22. Jaakkola J, Teppo K, Biancari F, et al. The effect of mental health conditions on the use of oral anticoagulation therapy in patients with atrial fibrillation: the FinACAF study. *Eur Heart J Q Care Clin Outcomes.* 2021. doi:10.1093/ehjqcco/qcab077
23. Teppo K, Jaakkola J, Airaksinen KEJ, et al. Mental health conditions and adherence to direct oral anticoagulants in patients with incident atrial fibrillation: a nationwide cohort study. *Gen Hosp Psychiatry.* 2022;74:88-93. doi:10.1016/j.genhosppsych.2021.12.012
24. Kirchhof P, Benussi S, Kotecha D, et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J.* 2016;37(38):2893-2962. doi:10.1093/eurheartj/ehw210
25. Camm AJ, Lip GYH, De Caterina R, et al. 2012 focused update of the ESC Guidelines for the management of atrial fibrillation. *Eur Heart J.* 2012;33(21):2719-2747. doi:10.1093/eurheartj/ehs253
26. Shi J, Tarkiainen L, Martikainen P, van Raalte A. The impact of income definitions on mortality inequalities. *SSM - Populat Health.* 2021;15:100915. doi:10.1016/j.ssmph.2021.100915
27. UNESCO Institute for Statistics. International Standard Classification of Education. <http://uis.unesco.org/sites/default/files/documents/international-standard-classification-of-education-isced-2011-en.pdf>
28. Kim D, Yang P-S, Jang E, et al. The optimal drug adherence to maximize the efficacy and safety of non-vitamin K antagonist oral anticoagulant in real-world atrial fibrillation patients. *Europace.* 2020;22(4):547-557. doi:10.1093/europace/euz273
29. Lowres N, Giskes K, Hespe C, Freedman B. Reducing stroke risk in atrial fibrillation: adherence to guidelines has improved, but patient persistence with anticoagulant therapy remains suboptimal. *Korean Circ J.* 2019;49(10):883. doi:10.4070/kcj.2019.0234
30. Rodríguez-Bernal CL, Peiró S, Hurtado I, García-Sempere A, Sanfélix-Gimeno G. Primary nonadherence to oral anticoagulants in patients with atrial fibrillation: real-world data from a population-based cohort. *J Manag Care Spec Pharm.* 2018;24(5):440-448. doi:10.18553/jmcp.2018.24.5.440
31. Banerjee A, Benedetto V, Gichuru P, et al. Adherence and persistence to direct oral anticoagulants in atrial fibrillation: a population-based study. *Heart.* 2020;106(2):119-126. doi:10.1136/heartjnl-2019-315307
32. Stoehr GP, Lu S-Y, Lavery L, et al. Factors associated with adherence to medication regimens in older primary care patients: the steel valley seniors survey. *Am J Geriat Pharmacother.* 2008;6(5):255-263. doi:10.1016/j.amjopharm.2008.11.001
33. Teppo K, Jaakkola J, Airaksinen KJ, et al. Mental health conditions and non-persistence of direct oral anticoagulant use in patients with incident atrial fibrillation: a nationwide cohort study. *JAHA.*
34. Liu C, Du X, Jiang C, et al. Long-term persistence with newly-initiated warfarin or non-VKA oral anticoagulant (NOAC) in patients with non-valvular atrial fibrillation: Insights from the prospective China-AF registry. *Med Sci Monit.* 2019;25:2649-2657. doi:10.12659/MSM.915875
35. Toorop MMA, Chen Q, Tichelaar VYIG, Cannegieter SC, Lijfering WM. Predictors, time course, and outcomes of persistence patterns in oral anticoagulation for non-valvular atrial fibrillation: a Dutch Nationwide Cohort Study. *Eur Heart J.* 2021;42(40):4126-4137. doi:10.1093/eurheartj/ehab421
36. Zielinski GD, Rein N, Teichert M, et al. Persistence of oral anticoagulant treatment for atrial fibrillation in the Netherlands: a surveillance study. *Research and Practice. Thromb Haemost.* 2020;4(1):141-153. doi:10.1002/rth2.12261
37. Maura G, Pariente A, Alla F, Billionnet C. Adherence with direct oral anticoagulants in nonvalvular atrial fibrillation new users and associated factors: a French nationwide cohort study. *Pharmacoeconom Drug Saf.* 2017;26(11):1367-1377. doi:10.1002/pds.4268
38. Jaam M, Ibrahim MIM, Kheir N, Awaisu A. Factors associated with medication adherence among patients with diabetes in the Middle

- East and North Africa region: A systematic mixed studies review. *Diabetes Res Clin Pract.* 2017;129: doi:10.1016/j.diabres.2017.04.015
39. Oosterom-Calo R, van Ballegooijen AJ, Terwee CB, et al. Determinants of adherence to heart failure medication: a systematic literature review. *Heart Fail Rev.* 2013;18(4):409-427. doi:10.1007/s10741-012-9321-3
40. Kilgore K, Pulungan Z, Teigland C, Parente A. The impact of demographic and socio-economic factors on medication adherence. *Value Health.* 2016;19(3):A289. doi:10.1016/j.jval.2016.03.751
41. Aylward BS, Rausch JR, Modi AC. An examination of 1-year adherence and persistence rates to antiepileptic medication in children with newly diagnosed epilepsy. *J Pediatr Psychol.* 2015;40(1):66-74. doi:10.1093/jpepsy/jsu010
42. Xu J, Prvu Bettger J, Pan Y, Liu L, Li Z, Wang Y. Association of socioeconomic status with medication persistence and adherence after an acute ischemic stroke. *European Stroke Journal.* 2018;3(1 Supplement 1).
43. King W, Lacey A, White J, Farewell D, Dunstan F, Fone D. Socioeconomic inequality in medication persistence in primary and secondary prevention of coronary heart disease – a population-wide electronic cohort study. *PLoS One.* 2018;13(3):e019408. doi:10.1371/journal.pone.0194081
44. Evans C, Marrie RA, Zhu F, et al. Adherence and persistence to drug therapies for multiple sclerosis: A population-based study. 2016.05.006
45. Malo S, Aguilar-Palacio I, Feja C, Lallana MJ, Armesto J, Rabanaque MJ. Effect of patient and treatment factors on persistence with antihypertensive treatment: A population-based study. *PLoS One.* 2021;16(1):e0245610. doi:10.1371/journal.pone.0245610
46. Arnet I, Kooij MJ, Messerli M, Hersberger KE, Heerdink ER, Bouvy M. Proposal of standardization to assess adherence with medication records: methodology matters. *Ann Pharmacother.* 2016;50(5):360-368. doi:10.1177/1060028016634106

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Teppo K, Jaakkola J, Biancari F, et al. Association of income and educational levels with adherence to direct oral anticoagulant therapy in patients with incident atrial fibrillation: A Finnish nationwide cohort study. *Pharmacol Res Perspect.* 2022;10:e00961. doi:[10.1002/prp2.961](https://doi.org/10.1002/prp2.961)